


What about Θ_0^c ?

Θ ($nd^2 \log n$) seen by many experiments (and by ZEUS)



: E N I T L O

Introduction Procedure Results Summary

DESY seminar, March 12, 2004



for the ZEUS Collaboration

Leonid Gladilin (MSU)



Charm pentadark search

We report a search for Θ_0^c signal in $M(D_{*-}d) (+ \text{c.c.})$ spectra

This decay mode can be dominant (Karliner-Lipkin, hep-ph/0401072)

If $M(\Theta_0^c) < M(D_{*+}) + M(d) = 2948 \text{ MeV}$, Θ_0^c can decay to $D_{*-}d (+ \text{c.c.})$

Such Θ_0^c would decay to $D_{*-}d (+ \text{c.c.})$

Cheung (hep-ph/0308176): $M(\Theta_0^c) = 2938 - 2997 \text{ MeV}$

$T(\Theta_0^c) \sim 21 \text{ MeV}$

Karliner-Lipkin (hep-ph/0307343): $M(\Theta_0^c) = 2985 \pm 50 \text{ MeV}$

can decay weakly to Θ_{+-}

Such Θ_0^c would be too light to decay to D mesons

Wu-Ma (hep-ph/0402244): $m(\Theta_0^c) = (4M(\Theta_0^c) + 2M(\Theta_0^*)) / 6 = 2704 \text{ MeV}$

Jaffe-Wilczek (hep-ph/0307341): $M(\Theta_0^c) = 2710 \text{ MeV}$

Predictions:

Introduction

Procedure: $D_{*+} \rightarrow D_0 \pi^+ \rightarrow (K^- \pi^s_+) \pi^s_+$ reconstruction


DATA 1995-2000 (126.5 pb $^{-1}$)
 $P_T(D_{*+}) > 1.35 \text{ GeV}, |\eta(D_{*+})| < 1.6$

Candidates with $144 < \Delta M < 147 \text{ MeV}$
(yellow band) were used. In this range

$$N(D_{*+}) = 42730 \pm 350$$

after background subtraction:

Summary of cuts:
 $P_T(K) > 0.45 \text{ GeV}, P_T(\pi) > 0.45 \text{ GeV}$
 $P_T(\pi^s) > 0.10 \text{ GeV}$
 $P_T(D_{*+}) / E_{\text{out} 10^\circ} > 0.12$
 $1.83 < M(K\pi) < 1.90 \text{ GeV}$ (wider for high $P_T(D_{*+})$)



tuned in the ZEUS non-charm pentadiquark analysis
In addition, require lower/upper limit from the proton dE/dx band


2) select protons with $P > 2 \text{ GeV}$

1) select protons with $P < 1.35 \text{ GeV}$ and require $dE/dx < 1.3$

two strategies:

π than a bulk of π , K
can/should be faster
protons from $b\bar{q}$ decays
for low- P protons
it is effective mostly
 dE/dx can be tried

$P_T(d) < 0.15 \text{ GeV}$





Procedure: selection of π candidates

$$0.3/D^2 + 0.8 < dE/dx < 1.0/D^2 + 1.2$$

dE/dx for p/\bar{p} candidates



$V_0 \leftarrow p\bar{t}$ from sec.vert.



Procedure: proton dE/dx band

Pitifully, no signal observed ...

$$M(D^*_d) = \Delta M_{\text{ext}} + M(D^*_{+})^{\text{PDG}} = M(K\pi\pi s) - M(K\pi\pi s) + M(D^*_{+})^{\text{PDG}}$$



Measured $M(D^*_d)$ spectra


Charmonium fragmentation universality requires $f(c \rightarrow \Theta_c^0)$ to be the same in ep , γp , pp and other interactions. Still it is useful to check DIS alone because it permits cleaner selection (smaller W^{ep} \iff smaller multiplicities) than in inclusive case but ~ 4.5 times smaller signal is cleaner.

$N(D^{*\pm}) = 9697 \pm 145$

$E_e > 8 \text{ GeV}, Q^2 > 1 \text{ GeV}^2$

$P_T(D^{*\pm}) > 1.35 \text{ GeV}, |\eta(D^{*\pm})| < 1.6$

DATA 1995-2000 (126.5 pb^{-1})





Charmonium fragmentation universality requires $f(c \rightarrow \Theta_c^0)$ to be the same in ep , γp , pp and other interactions. Still it is useful to check DIS alone because it permits cleaner selection (smaller W^{ep} \iff smaller multiplicities) than in inclusive case but ~ 4.5 times smaller signal is cleaner.

Procedure: $D^{*\pm}$ in DIS with $Q^2 > 1 \text{ GeV}^2$

again, nothing to fit ...

$$M(D^*_s p) = \Delta M_{\text{ext}} + M(D^*_{s+})^{\text{PDG}} = M(K\pi\pi s p) - M(K\pi\pi s) + M(D^*_{s+})^{\text{PDG}}$$



Measured $M(D_s^* p)$ spectra in DIS with $Q^2 > 1 \text{ GeV}^2$

Signal did not show up

selecting DIS with $Q^2 < 1 \text{ GeV}^2$ (was shown) or $Q^2 > 15 \text{ GeV}^2$
 varying dE/dx requirements for low- P selection
 no dE/dx requirements for high- P selection
 require in addition $\cos\Theta_*(d) < -0.7$, where $\Theta_*(d)$ is the angle
 between p direction in \vec{q} r.f. and \vec{q} direction in the lab
 studying/removing reflections from $D^{**} \leftarrow D_{\pm} \pi_{\pm}$
 removing the cut on $P_T(D^*)/E_{\text{out } 10^\circ}$; using $z(D^*) < 0.2$ instead
 making all cuts „as close as possible to H1 selection“

Systematic studies

high- P selection : 0.4% from $N(D_{\pm}^*)$

low- P selection : 0.3% from $N(D_{\pm}^*)$

$$\frac{N_{\text{rec}}(\text{all } d)}{N_{\text{rec}}(P(d) < 2 \text{ GeV})} \sim 40\%$$

$$\% \sim 30 \frac{N_{\text{rec}}(\text{all } d)}{N_{\text{rec}}(P(d) > 1.35 \text{ GeV}, dE/dx(d) < 1.3)}$$

$$\% \sim 1 \frac{N_{\text{rec}}(D_{\pm}^*)}{N_{\text{rec}}(\Theta_0^c \rightarrow D_{-}^* \rightarrow d + \text{c.c.})}$$



Naïve estimation of expected signals (inspired by HI observations):

$$f(c \rightarrow \Theta_0^c) \times B(\Theta_0^c \rightarrow D_{-}^* \rightarrow d)$$

we are not yet ready with the upper limit on

Naïve estimation of expected signals



so large signals are excluded



Naïve signal expectations

so large signals are certainly not here

sensitivity is smaller



Naive signal expectations in DIS with $Q^2 > 1 \text{ GeV}^2$


Using all HERA-I data (126.5 pb^{-1}),
the ZEUS collaboration does not see
any resonance structure
in $M(D^*_s p)$ spectra
The ZEUS data constrain
the uncorrected fraction of D_s^\pm mesons
originating from Θ_c^0 decays
to be below 1%

Summary


$$N(V_\pm) = 1390 \pm 380$$

$$N(D_\pm) = 9600 \pm 620$$

no dE/dx selection (high- P_T range)



no lifetime tagging (pre-MVD data)



$$\mathcal{L}_{int} = 65.5 \text{ pb}^{-1}$$

Backup: $D_\pm \rightarrow K^\pm \pi^\pm \pi^\mp$ and $V_\pm \rightarrow K^\pm p \bar{p}$

of charm fragmentation

HERA measurements confirms universality

We use correct normalisation for PQCD predictions

charm fragmentation fractions are universal

$$f(c \rightarrow D^*_+) = 0.263 \pm 0.019^{+0.056}_{-0.042} \pm 0.022 \pm 0.021$$

$$f(c \rightarrow D^*_+) = 0.223 \pm 0.009^{+0.003}_{-0.005}$$

$$0.076 \pm 0.007$$

$$f(c \rightarrow D^+_+) = 0.076 \pm 0.020^{+0.017}_{-0.001}$$

$$0.156 \pm 0.043^{+0.036}_{-0.035} \pm 0.046$$

$$f(c \rightarrow D_s^+) = 0.107 \pm 0.009 \pm 0.005$$

$$0.549 \pm 0.023$$

$$f(c \rightarrow D_0) = 0.557 \pm 0.019^{+0.005}_{-0.013}$$

$$0.202 \pm 0.020^{+0.045}_{-0.033} \pm 0.021$$

$$f(c \rightarrow D_+) = 0.249 \pm 0.014^{+0.004}_{-0.008}$$

e^+e^- data
Combined

$P_T(D, V_c) > 3.8 \text{ GeV}, |\eta(D, V_c)| < 1.6$

H1 prel. (DIS)

Backup: fragmentation fractions

(ZEUS prel.)

$$f(c \rightarrow D^*_{s+}) \cdot B^{D^*_{s+} \rightarrow D^* \pi^+ \pi^-} < 0.7\% \quad (95\% \text{ C.L.})$$

Using world average for $f(c \rightarrow D^*_{s+})$:

after backgr. subtraction: “ $N(D^*_{s+}) = 91 \pm 75$

covers both predictions and DELPHI’s observation

Search window: $2.59 < \Delta M_{ext} + M(D^*_{s+}) < 2.67 \text{ GeV}$

$$\Delta M_{ext} = M(K\pi\pi S\pi\pi) - M(K\pi\pi)$$

\Rightarrow ZEUS search



CLEO and OPAL did not confirm



$$T < 15 \text{ MeV}$$

Observed by DELPHI (~ 50): $M = 2637 \text{ MeV}$

$$D^*_{s+} \rightarrow D^* \pi^+ \pi^-$$

Backup: search for radially excited D^*_{s+} meson





expected for $j_p = 1^-, 2^+$

does not contradict to $R = -1$


ZEUS : consistent with $R = 0$, i.e. $j_p^+ = j_+$

CLEO ($D_+^{s1} \rightarrow D^*_0 K_+$): $R = -0.23_{-0.40}^{+0.32}$

$$R = -0.53 \pm 0.32(\text{stat.})_{+0.05}^{-0.14}(\text{syst.}) \quad (\text{ZEUS prel.})$$

Fit to a form : $1 + R \cos^2 \alpha$

Helicity angle α : between K_0^s and π^s in $D^{*\pm}$ r.f.




$$M(D_{s1}^+) = 2534.2 \pm 0.6 \pm 0.5 \text{ MeV} \quad (\sim M_{\text{PDG}})$$

$$N(D_+^{s_1}) = 62.3 \pm 9.3$$

$$\nabla M_{ext} = M(K - M(s) - M(S))$$

$$D_{\pm}^{s1}(2536) \rightarrow D^{\ast\pm} K_0^s, \quad K_0^s \rightarrow \pi^+ \pi^-$$

Backup: charm-strange $D_{\pm}^{s1}(2536)$ meson



- 1) the same amounts of excited D mesons in e^+e^- and ep data
- 2) situation with $f(c \rightarrow D_0^*)$ is not clear
- 3) $f(c \rightarrow D_{+}^{s1})$ is twice as large as the expectation :
 $\gamma_s \times f(c \rightarrow D_0^1) \approx 0.3 \times 2\% = 0.6\%$ Why ?

| | $f(c \rightarrow D_0^1) [\%]$ | $f(c \rightarrow D_0^*) [\%]$ | $f(c \rightarrow D_{+}^{s1}) [\%]$ |
|--------------|--|--|--|
| ZEUS (prel.) | $1.46 \pm 0.18^{+0.33}_{-0.27} \pm 0.06$ | $2.00 \pm 0.58^{+1.40}_{-0.48} \pm 0.41$ | $1.24 \pm 0.18^{+0.08}_{-0.06} \pm 0.14$ |
| CLEO | 1.8 ± 0.3 | 1.9 ± 0.3 | |
| OPAL | 2.1 ± 0.8 | 5.2 ± 2.6 | $1.6 \pm 0.4 \pm 0.3$ |
| ALEPH | 1.6 ± 0.5 | 4.7 ± 1.0 | $0.94 \pm 0.22 \pm 0.07$ |
| DELPHI | 1.9 ± 0.4 | 4.7 ± 1.3 | |

Using world average for $f(c \rightarrow D_{+}^*)$:

Backup: fragmentation fractions for excited D mesons

$D^{*\pm}$ in PHP : reconstructed $D^{*\pm}$ in tagged/untagged PHP events

Inclusive PHP : dijet events

$D^{*\pm}$ in DIS : reconstructed $D^{*\pm}$ in DIS events (low Q^2)

Inclusive DIS : almost offine selection

Third level trigger:

Tagged PHP : 44/35m taggers, CAL energies and SLT tracks

Untagged PHP : CAL energies and SLT tracks (high-W)

DIS : scattered electron and CAL energies

Second level trigger:

Tagged PHP : 44m and 35m taggers, CTD-CAL and CTD-FLT

Untagged PHP : CTD-CAL and CTD-FLT

DIS : scattered electron (and CTD-FLT)

CTD-FLT: “tracks” looking to the nominal interaction point

CAL-FLT: regional energy sums

First level trigger:

Backup: trigger selection