

## Full Simulation Study of

$$W^+W^- \rightarrow t\bar{t}$$


at TESLA

Juan Alcaraz  
CIEMAT, Madrid

Ester Ruiz Morales  
UAM, Madrid

7 th ECFA/DESY Workshop  
for a Linear  $e^+e^-$  Collider  
DESY, 22-25 September 2000


## Motivation




$W^+W^- \rightarrow t\bar{t}$   
at High Energy  
 $e^+e^-$  Colliders

Great physical interest in models of strong-interaction  
Electroweak Symmetry Breaking

- High Energy  $W_L \equiv \pi$ -GB from the symmetry breaking interactions. In S.I. models:  $W_L$  amplitudes enhanced, may have new resonances.
- Depending on models: can be the same resonances as in  $VV$ -scattering, or new ones from the dynamics generating the top quark mass.



Test Higgs Mechanism



Test top-mass generation

- Scalar or Vector resonance exchange could be distinguished by top quark polarization analysis.
- At LHC,  $WW \rightarrow t\bar{t}$  hidden under the huge QCD background.

Observable at planned  $e^+e^-$  Linear Colliders ?

# Strong - interaction

Electroweak Symmetry Breaking  $\equiv$

The Higgs Sector responsible for EW symmetry breaking is formed by a new set of particles with new strong interactions with a typical scale  $\sim 1 \text{ TeV}$ .  
( $\Rightarrow$  there is no fundamental / light Higgs boson)

- The strong interactions could form resonances  $\equiv$  bound states of the new set of particles.
- "SM heavy Higgs" means in this context: scalar bound state with the same couplings to  $W$  and  $t$  as a fundamental Higgs.  
(this will happen, in general, if the strong-interactions that form the scalar resonance are responsible of all EW symmetry breaking)

## Previous Studies

- Theoretical Study: (Sitges, ERM and M. Peskin)
  - $WW \rightarrow t\bar{t}$  in different S.I. Models:
    1. SM: Heavy Higgs gives  $LL$  and  $RR$   $t\bar{t}$  pairs.
    2. Technicolor:  $T_\rho$  couples to  $LR$  and  $RL$  final states.  $T_\rho$  exchange as in  $VV$ -scattering.
    3. Topcolor: New scalar (Top-Higgs) exchange. Not relevant in  $VV$ -scattering.
  - Signals at 1.5 TeV LC: In several models analyzed, the new resonances give cross section enhancements of similar size to the SM Higgs boson signals.
  - $t\bar{t}$  polarization analysis based on experimental assumptions.
- SM Simulation: (Snowmass, T. Barklow)  
1.5 TeV Collider
- SM Simulation: (JLC Study, Tsukamoto)  
Early nineties, old collider parameters.

## Our analysis

- Realistic simulation study for TESLA:  $\sqrt{s} = 1$  TeV,  $\mathcal{L} = 1$  ab<sup>-1</sup>.
- Full calculation including reducible and irreducible backgrounds. Need reliable Event Generators.
- Include ISR and Beamstrahlung (CIRCE).
- Realistic Detector Simulation (SIMDET, V3.02).
- Detailed event reconstruction.

# Event Generators 1

- We found some difficulties:

Generator	Advantages	Shortcomings
CompHep	<ul style="list-style-type: none"> <li>- All SM Diagrams</li> <li>- Fast</li> <li>- Reliable</li> </ul>	<ul style="list-style-type: none"> <li>- Hard to include: Beamstrahlung</li> <li>- New models</li> </ul>
Pandora	<ul style="list-style-type: none"> <li>- Full helicity</li> <li>- Top decays</li> <li>- Easy to modify <i>models</i></li> </ul>	<ul style="list-style-type: none"> <li>- Only Fusion Diagrams</li> <li>- Effective-W approx with <math>p_T</math> functions</li> </ul>
NextCalibur	<ul style="list-style-type: none"> <li>- All <math>ee \rightarrow 4f</math> SM diagrams</li> <li>- separate <math>t\bar{t}</math> helicities</li> </ul>	<ul style="list-style-type: none"> <li>- Too slow</li> <li>- Improvements needed</li> </ul>

- Main Problem:

At  $\sqrt{s} = 1.5 \text{ TeV}$ ,  $WW$ -fusion + Effective-W approximation with helicity- and  $p_T$ - dependent  $W$  distributions OK, but

At  $\sqrt{s} = 1 \text{ TeV}$ , there are negative interference effects among irreducible backgrounds and non-fusion diagrams cannot be neglected

⇒ Need full  $ee \rightarrow 4f$  generator for studies at TESLA

- NextCalibur = Excalibur  $ee \rightarrow 4f$  program + Higgs diagrams + "Massive" fermions + ISR improved.



We had to implement:

- Weighter → Event Generator.
- Run on Linux and other platforms.
- Include Beamstrahlung (CIRCE).
- Control on Higgs mass, and final top polarizations.
- Implement helicity-dependent top decay kinematics (Pandora).

## Event Generators 2

$$\sqrt{s} = 1.5 \text{ TeV}, \quad m_H = 800 \text{ GeV}$$



(Generator Level, no ISR, no Beamstrahlung)



## Event Generators 3

$$\sqrt{s} = 1 \text{ TeV}, \quad m_H = 800 \text{ GeV}$$

(Generator Level, no ISR, no Beamstrahlung)



## SM Signal and Backgrounds


- **Signal:**  $e^+e^- \rightarrow \nu\bar{\nu}t\bar{t}$  Helicity Cross Sections (fb):

$m_H$ (GeV)	$\sigma(\text{LL}) = \sigma(\text{RR})$	$\sigma(\text{LR})$	$\sigma(\text{RL})$	Total
100	0.113	0.22	0.21	0.66
500	1.65	0.22	0.21	3.75
800	0.33	0.22	0.21	1.11

(Generator Level, with IRS and Beamstrahlung)

- Events looks like  $e^+e^- \rightarrow t\bar{t}$  but:

- Large missing T and L momentum from two  $\nu$ 's.  $\sim \frac{M_W}{2}$
- Lower visible mass, in general.



- **Huge but manageable backgrounds:**

$$\sigma(ee \rightarrow q\bar{q}) = 5400 \text{ fb} \quad \rightarrow (5.4 \text{ Mevents!})$$

$$\sigma(ee \rightarrow W^+W^-) = 3700 \text{ fb} \quad \rightarrow (3.7 \text{ Mevents!})$$

- **Critical backgrounds:**

$$\sigma(ee \rightarrow t\bar{t}) = 243 \text{ fb}$$

$$\sigma(ee \rightarrow eett\bar{t}) = 17 \text{ fb}$$

- Backgrounds generated with **PYTHIA**.

In this analysis, we have only used 6 jet events

(4.000)


(60.000)

(200.000)


(20.000)

# Preselection Cuts 1

- $M_{\text{Miss}} \geq 100 \text{ GeV}$ : Cuts  $ee \rightarrow t\bar{t}$ . (ISR + Beaus.)




- $E_T \geq 50 \text{ GeV}$ : Cuts  $ee \rightarrow eett$ . ( $\gamma\gamma$ -fusion)



## Preselection Cuts 2

- **No lepton:** force event into 5 jets and reject it if find lepton with  $E_{Lept} \geq 0.7E_{Jet}$ .
- **Force six jet event:** Durham Algorithm, require  $Y_{Cut}^{(5-6)} \geq 0.0005$ .  
Cuts  $ee \rightarrow q\bar{q}, WW$ .



After Preselection cuts  
 still huge background from misreconstructed  $ee \rightarrow t\bar{t}$

## Jet Association for the best $\chi^2$

- Find the combination giving best





$$\chi_{\min}^2 = \frac{(M_{W1} - M_W)^2}{(5\text{GeV})^2} + \frac{(M_{W2} - M_W)^2}{(5\text{GeV})^2} + \frac{(M_{t1} - M_W)^2}{(10\text{GeV})^2} + \frac{(M_{t2} - M_W)^2}{(10\text{GeV})^2}$$

and keep the event only if this combination is within  $5\sigma$  of the expected mass values

$$|M_{W_i} - M_W| \leq 5\sigma \approx 25\text{GeV}; \quad |M_{t_i} - M_W| \leq 5\sigma \approx 50\text{GeV}$$

This rejects misreconstructed  $ee \rightarrow tt$  events.

- Reconstructed  $t$  and  $W$  masses:






Final acceptance for signal events  $\approx 10\%$ .

(OK with Tsukamoto)


## Final Cuts

- $\text{Thrust} \leq 0.9$ ,  $C_{\text{par}} \geq 0.5$ : Select spherical events.
- $\text{Major} \geq 0.3$ : Select non-planar events.
- Moderate  $b$ -tagging: To further reject  $ee \rightarrow qq, WW$ .  
Based on consistency with primary vertex.




Negligible  $qq$  and  $WW$   
backgrounds after  
these cuts


# Higgs Signals




$m_H = 400$  GeV  
Signal = 693  
Bckgrd = 1218




# Higgs Signals




$m_H = 500$  GeV  
Signal = 450  
Bckgrd = 1218



# Higgs Signals



$m_H = 800$  GeV  
Signal = 54  
Bckgrd = 1218



## Higgs Signal Significance


• **Signal:**  $X(m_H) \equiv \sigma_{\text{seen}}(m_H) - \sigma_{\text{expected}}(m_H = 100)$ .

**Background:**  $\sigma_{\text{expected}}(m_H = 100)$ .

$m_H$ (GeV)	375	400	500	600	800
N. Signal Events	342	543	351	157	42
N. Backg. Events	520				


(Number of Events with  $M_{\text{vis}} + M_{\text{miss}} \geq 900$  GeV.)

• Fits to  $X(m_H)$  using:  $M_{\text{vis}}$ : 10 bins (300-800) GeV.  
 $M_{\text{vis}} + M_{\text{miss}}$ : 10 bins (600-1000) GeV.



**$H \rightarrow t\bar{t}$  Signal Observable with Significance  $\geq 5\sigma$   
for  $m_H$  in 350-650 GeV range at TESLA.**

# Comparison with Higgs Signal from VV-Scattering



Higgs Signal Significance

$m_H$ (GeV)	TESLA (1 TeV, $1 \text{ ab}^{-1}$ ) $H \rightarrow t\bar{t} \rightarrow (6j)$	ATLAS (14 TeV, $100 \text{ fb}^{-1}$ ) $H \rightarrow VV \rightarrow (\text{all})$
375	16	35
400	20	40
500	13	40
600	6	40
800	2	12

## Top quark Yukawa Measurements

- At LHC:  $t\bar{t}H$ :  $\frac{\delta Y_t}{Y_t}(\text{stat}) \approx 10\%$  for  $m_H \leq 120$  GeV. (Atlas TDR)

- At  $e^+e^-$  LC:


$e^+e^- \rightarrow t\bar{t}H(H \rightarrow b\bar{b})$ :  $\frac{\delta Y_t}{Y_t} \approx 10\%$  for  $m_H \leq 120$  GeV.

(Baer, Dawson, Reina), (Juste, Merino)

$e^+e^- \rightarrow t\bar{t}Z$ :  $\frac{\delta Y_t}{Y_t} \approx 10\%$ , for  $m_H \approx 2m_t$  GeV.

(Fujii) (old JLC setting)

$e^+e^- \rightarrow \nu\bar{\nu}t\bar{t}$ : At TESLA,  $\sqrt{s} = 1\text{TeV}$ ,  $\mathcal{L} = 1\text{ab}^{-1}$  we find



$m_H$ (GeV)	$\Delta Y_t/Y_t$ (%)
375	2.7
400	2.2
500	3.9
600	7.8
800	30.5

Possible to measure  $Y_t$  with  $\frac{\delta Y_t}{Y_t} \leq 10\%$ ,  
for  $m_H$  in 350-650 GeV range at TESLA.

## Conclusions

- $W^+W^- \rightarrow t\bar{t}$  interesting at TESLA, if there is no light Higgs boson.
- A  $WW \rightarrow H \rightarrow t\bar{t}$  signal will be observable in TESLA if  $\sqrt{s} = 1$  TeV,  $\mathcal{L} = 1$  ab $^{-1}$  and  $m_H \leq 650$  GeV.
- The Higgs-top Yukawa coupling will be measured with a precision better than 10%, if  $\sqrt{s} = 1$  TeV,  $\mathcal{L} = 1$  ab $^{-1}$  and  $M_H \leq 650$  GeV.
- Unique way to measure the Top Yukawa coupling to a heavy Higgs boson (at LHC,  $H \rightarrow t\bar{t}$  is not observable for  $m_H > 120$  GeV.)
- Similar results can be expected in other scenarios of strong electroweak symmetry breaking (technicolor).
- Extensions of this analysis for TESLA still under study:
  - $\sqrt{s} = 800$  GeV.
  - Include semileptonic  $t\bar{t}$  decays.
  - Top polarization studies.