



LCD-LCWS

NLC – The Next Linear Collider Project



Presentation at the LCWS 2005 Meeting

*The Importance of Positron Polarization and the
Deleterious Effects of Beam/Bremmstrahlung on
the Measurement of Supersymmetric Particle
Masses and other Parameters*

March 2005



LCD-LCWS

NLC – The Next Linear Collider Project



THE POSITRON POLARIZATION GROUP

CERN-PH-TH/2005-036, DCPT-04-100, IPPP-04-50

G. Moortgat-Pick, et. al.



LCD-LCWS

NLC – The Next Linear Collider Project



THE COLORADO GROUP

Lisa Boyle, Shenjian Chen, Bradford Dobos, Keith Drake,

Chris Geraci, Jack Gill, Jason Gray, Andrew Hahn,

Kyle Miller, Martin Nagel, Uriel Nauenberg,


Matthew Phillips, Joseph Proulx,

Will Ruddick, Jesse Smock, Jinlong Zhang



ACTIVITIES

- ◆ *The Importance of Positron Polarization in Determining SUSY Masses and other Parameters.*
- ◆ *Simulation of Supersymmetry. New method to overcome the negative effects of beamstrahlung and bremsstrahlung.*





LCD-LCWS



NLC – The Next Linear Collider Project

Positron Polarization Helps





LCD-LCWS




NLC – The Next Linear Collider Project


electron Left Pol 80%


electron Right Pol 80%

$P(e^+) \text{ vs. } \sigma \text{ For } P(e^-)=80\% \text{ L}$



$P(e^+) \text{ vs. } \sigma \text{ For } P(e^-)=80\% \text{ R}$









LCD-ALCPG



NLC – The Next Linear Collider Project

Electron, Positron Energy Spectrum from $\tilde{e}^+ \tilde{e}^- \rightarrow \text{all } e e$ *e⁻ Spect. e⁻ 80%R e⁺ Spect.* *e⁻ Spect. e⁻ 80%R e⁺ Spect.*





LCD-ALCPG




NLC – The Next Linear Collider Project

Muon Energy Spectrum from $e^+ e^- \rightarrow \tilde{\mu}^+ \tilde{\mu}^-$


e^- 80% R e^+ 80% L

e^- 80% L e^+ 80% R

e^- 80%R e^+ 80%L to $\mu^- \mu^+$ with $W^+ W^+$ Background (750 GeV)



e^- 80%L e^+ 80%R to $\mu^- \mu^+$ with $W^+ W^+$ Background (750 GeV)





LCD-LCWS

NLC – The Next Linear Collider Project



These selectron, smuon signals with various electron and positron polarization are clear evidence for supersymmetry.!!!

These energy distributions can not be produced with Standard Model processes.

Need positron polarization to observe dramatic energy distribution shape variations.




LCD-LCWS



NLC – The Next Linear Collider Project

Measurement of the sfermion Mixing Angle ($\theta_{\tilde{f}}$)

Varying the electron and positron polarizations





LCD-LCWS

NLC – The Next Linear Collider Project



*This is one case where removal of the
2 γ process is crucial*

The leptons from stau decays are soft.



LCD-LCWS






NLC – The Next Linear Collider Project

$$E_{cm} = 500 \text{ GeV}$$

$$\sigma(e^+ e^- \rightarrow \tilde{t}_1 \tilde{t}_1) \text{ fb}$$

$$M_{\tilde{t}_1} = 200 \text{ GeV}$$







LCD-LCWS




NLC – The Next Linear Collider Project





Measuring the sfermion mixing angle

The sfermion production goes via γ , Z exchange in the s-channel and the coupling constants depend on the sfermion mixing angle (θ_i). The cross section can be enhanced by varying the positron polarization and the sensitivity on the mixing angle can be determined more readily and measured.



LCD-LCWS



NLC – The Next Linear Collider Project

Study of Chargino and Neutralino Production with Positron Polarization.



LCD-LCWS




NLC – The Next Linear Collider Project

$E_{cm} = 1 \text{ TeV}$


$\sigma(\tilde{\chi}_i^+ \tilde{\chi}_j^-) \text{ fb}$

SPS1a

$P_{e^-} = -0.9$



$P_{e^-} = +0.9$





LCD-LCWS





NLC – The Next Linear Collider Project

$E_{cm} = 1 \text{ TeV}$

$\sigma(\tilde{\chi}_1 \tilde{\chi}_j) \text{ fb}$

SPS1a





LCD-LCWS



NLC – The Next Linear Collider Project

$$\sigma(\tilde{\chi}_1 \tilde{\chi}_2) \times B.R.(\tilde{\chi}_2 \rightarrow \tilde{l}_R l_1) \times B.R.(\tilde{l}_R \rightarrow l_2 \tilde{\chi}_1^0) \text{ fb}$$

$$\begin{matrix} \rightarrow & & \rightarrow & & \rightarrow \\ \rightarrow & & \rightarrow & & \rightarrow \end{matrix}$$

Define $T = \{P(e^-) \times P(l_2)\} - P(l_1)$

$$\frac{\sigma(T > 0) - \sigma(T < 0)}{\sigma(T > 0) + \sigma(T < 0)}$$

$$A(T) = \frac{\sigma(T > 0) - \sigma(T < 0)}{\sigma(T > 0) + \sigma(T < 0)}$$

$$A_{CP} = \text{Diff. in Pol. of } \tau$$



LCD-LCWS

NLC – The Next Linear Collider Project




Determined the values of σ and $A(T)$ as a function of electron and positron polarization for the following parameters

$$M_0 = 100 \text{ GeV} \quad \tan(\beta) = 10$$

$$M_2 = 400 \text{ GeV} \quad \varphi(M_1) = 0.2\pi$$

$$|A_\tau| = 250 \text{ GeV} \quad \varphi(A_\tau) = \varphi(\mu) = 0$$

$$|\mu| = 240 \text{ GeV}$$





LCD-LCWS




NLC – The Next Linear Collider Project

$$\sigma(\tilde{\chi}_1^0 \tilde{\chi}_1^0 l_1 l_2) \text{ fb}$$

A_T in %







LCD-LCWS



NLC – The Next Linear Collider Project

$\sigma (\tilde{\chi}_1^0 \tilde{\tau}_1^+ \tau^-) fb$

$A_{CP} \%$







LCD-LCWS



NLC – The Next Linear Collider Project

$$\sigma(\tilde{\chi}_1^0 \tilde{\tau}_1^+ \tau^-) \text{ fb}$$

$$A_{CP} \%$$






LCD-LCWS



NLC – The Next Linear Collider Project

$$\tilde{\chi}_1^0 \tilde{\chi}_2^0, \tilde{\chi}_2^0 \rightarrow \tilde{\chi}_1^0 l^+ l^- \quad \chi_3^0 \chi_2^0, \chi_2^0 \rightarrow \chi_1^0 \bar{l}^+ l^-$$





LCD-LCWS

NLC – The Next Linear Collider Project



WORD OF CAUTION

All neutralino signals than end in 2 leptons without a mass constraint (like Z^0) are overwhelmed by selectron and sneutrino production channels because of t -channels and by 2 γ channels specially if the leptons come from τ decays.

Needs careful simulation. Playing with positron polarization should help.



LCD-LCWS

NLC – The Next Linear Collider Project



The Effect of Beam-Bremmstrahlung





LCD-LCWS



NLC – The Next Linear Collider Project

CM Energy Distribution with Beamstrahlung





Simulation of Selectron Production Case Study

- *Consider Case SPS1 , $M_{1/2} = 250 \text{ GeV}$, $M_0 = 100 \text{ GeV}$.*
- *Mass of $e_R = 143.11 \text{ GeV}$, Mass of $e_L = 204.6 \text{ GeV}$, Mass of $\chi_1^0 = 95.47$*
- *Compare Fits with Beam and Bremsstrahlung and without.*
- *We use the $e^+ - e^-$ Energy Spectra Subtraction Technique to remove Standard Model Background.*




LCD-LCWS




NLC – The Next Linear Collider Project


Selectron Production

$e^+ - e^-$ Energy Spectra





Chi-Square Fits for the SPS1 Snowmass Point



$M_{1/2} = 250$ GeV
 $M_0 = 100$ GeV
 $\tan(\beta)$ fixed at 10




LCD-LCWS




NLC – The Next Linear Collider Project


Resultant Fits to Energy Edges

No Bremm



Bremm





LCD-LCWS

NLC – The Next Linear Collider Project



New Method to Determine Masses


- ◆ *Compare Energy Spectrum to those Generated with different parameters encompassing the correct one.*
- ◆ *Do a Chi Square Fit to the Spectra Comparison.*
- ◆ *Choose the minimum and determine the masses.*



LCD-LCWS



NLC – The Next Linear Collider Project







LCD-LCWS



NLC – The Next Linear Collider Project

$M_{1/2}$ vs M_0 curves for $M_{sel L}$ values

$M_{1/2}$ vs M_0 curves for $M_{sel R}$ values






LCD-LCWS



NLC – The Next Linear Collider Project

$M_{1/2}$ vs M_0 curves for $M(\chi_1^0)$



No dependence on $\tan(\beta)$



LCD-ALCPG



NLC – The Next Linear Collider Project

Simulation of Selectron, Smuon Production Case Study

- *Consider Case SPS3 , $M_{1/2} = 400 \text{ GeV}$, $M_0 = 90 \text{ GeV}$.*
- *Mass of $e_R = 179.1 \text{ GeV}$, Mass of $e_L = 292.5 \text{ GeV}$, Mass of $\chi^0_1 = 158.2 \text{ GeV}$.*
- *Compare Fits with Beam and Bremsstrahlung and without.*
- *For selectrons we use the $e^+ - e^-$ Energy Spectra Subtraction Technique to remove Standard Model Background.*




LCD-LCWS



NLC – The Next Linear Collider Project

Chi Square Fit for the SPS3 Snowmass Point




$$M_{1/2}(expec.) = 400 \text{ GeV}$$

$$M_{1/2}(fit) = 400.22^{+0.19}_{-0.54} \text{ GeV}$$

M_0 fixed at 90 GeV

$\tan(\beta)$ fixed at 10



LCD-ALCPG




NLC – The Next Linear Collider Project

Muon Energy Spectrum from $e^+ e^- \rightarrow \tilde{\mu}^+ \tilde{\mu}^-$


e^- 80% R e^+ 80% L

e^- 80% L e^+ 80% R

e^- 80%R e^+ 80%L to $\mu^- \mu^+$ with $W^+ W^+$ Background (750 GeV)



e^- 80%L e^+ 80%R to $\mu^- \mu^+$ with $W^+ W^+$ Background (750 GeV)







LCD-LCWS



NLC – The Next Linear Collider Project







LCD_LCWS



NLC – The Next Linear Collider Project

SPS3 Point; $E_{cm}=750$ GeV ; $M_{1/2}=400$ GeV, $M_0=90$ GeV





LCD-LCWS



NLC – The Next Linear Collider Project

Resultant Masses from Fits

	<i>Input Masses</i>	<i>Mass fit E.P.</i>	<i>Mas Fit ChiS.</i>
$\tilde{\mu}_R$	179.1	171.3	179.0
$\tilde{\mu}_L$	292.5	287.4	292.0



LCD-LCWS



NLC – The Next Linear Collider Project

Study of Sneutrino Production

A Very Interesting Case

SPS6 Point

$$E_{cm} = 750 \text{ GeV}$$

$$M_0 = 150 \text{ GeV}, M_{1/2} = 300 \text{ GeV}$$

$$A_0 = 0 \text{ GeV}, \tan(\beta) = 10$$

$$M_{\tilde{\nu}} = 243.8 \text{ GeV}; M_{\tilde{\chi}_1^+} = 222.4 \text{ GeV}$$

~ ~ ~ ~

$$\nu_e \rightarrow \tilde{\chi}_1^+ e^-; \tilde{\chi}_1^+ \rightarrow \tilde{\chi}_1^0 W^+; W^+ \rightarrow \text{hadrons}$$




LCD-LCWS



NLC – The Next Linear Collider Project

Energy Spectrum of Hadronic Jets after Hadronic Mass (W) Cut







LCD-LCWS



NLC – The Next Linear Collider Project

No Beam/Bremmstrahl

Beam/Bremmstrahl










LCD-LCWS



NLC – The Next Linear Collider Project

Sneutrino Mass Dependence on Parameters









LCD-LCWS



NLC – The Next Linear Collider Project

Chargino Mass Dependence on Parameters








LCD-LCWS



NLC – The Next Linear Collider Project

Chi-Square fits to the Electron Energy Distribution





LCD-LCWS

NLC – The Next Linear Collider Project



Resultant Masses from Fits

<i>Input Mass</i>	<i>Before Strahl</i>	<i>After Strahl</i>	<i>ChiS Fit</i>	
	<i>End Point</i>	<i>End Point</i>		
ν_e	243.8	243.6	248.9	243.5
χ_1^+	222.4	222 .1	227.4	222 .0



LCD-LCWS

NLC – The Next Linear Collider Project



CONCLUSION

The slepton, sneutrino signals are easy to observe and easy to measure with positron polarization if the 2 photon process is tagged with excellent efficiency.

The masses depend on all the parameters of the SUGRA model and hence we can determine consistency of M_0 and $M_{1/2}$ with high accuracy ($\sim 0.2\%$) and determine A_0 and $\tan(\beta)$.



LCD-LCWS

NLC – The Next Linear Collider Project



Neutralino Production Study

$$e^+ e^- \rightarrow \tilde{\chi}_2^0 \tilde{\chi}_2^0$$

$$\tilde{\chi}_2^0 \rightarrow Z^0 + \tilde{\chi}_1^0$$

$$Z^0 \rightarrow l^+ l^- \text{ one decay}$$

$$Z^0 \rightarrow q q \text{ other decay}$$




LCD-LCWS



NLC – The Next Linear Collider Project

Energy Distribution of the Z







LCD-LCWS



NLC – The Next Linear Collider Project

Dependence of $\tilde{\chi}_2^0$ Mass on $\tan(\beta)$






LCD-LCWS



NLC – The Next Linear Collider Project



LCD-LCWS



NLC – The Next Linear Collider Project



LCD-LCWS



NLC – The Next Linear Collider Project