

# Polarized $e^-$ and $e^+$ at the ILC

## Summary of the Polarization Report

*Gudrid Moortgat-Pick (CERN)  
for the POWER group*

LCWS05

Stanford, 22/03/2005

### 1. 'Polarization talks' at LCWS05

### 2. Polarization report $\sim$ finished!

- The physics case for polarized  $e^-$  and  $e^+$  beams
- Machine overview
- Polarimetry overview

### 3. Further news and events

## Very active 'POWER' group: Polarization related talks at LCWS05

- 'Physics':

- \* 'The importance of  $e^+$  polarization': U. Nauenberg@Susy
- \* 'Review of physics with polarized beams': N. Paver@New Physics

- 'Machine':

- \* 'Polarized  $e^+$  sources at the ILC': V. Bharadwaj@Accelerator
- \* 'Enhanced Fabry Perot resonators': A. Variola@Accelerator
- \* 'Status of experiment E166 at SLAC': R. Pöschl@Accelerator
- \* 'Beam dynamic simulation': W. Gai@Accelerator
- \* 'Polarized  $e^+$  Generation at KEK-ATF': T. Omori @Accelerator
- \* + many questions and discussion

- 'Polarimetry':

- \* 'Polarimetry for 2 IR's and downstream designs': K. Moffeit@MDI
- \* 'Upstream Compton Polarimetry': N. Meyners@MDI

⇒ Only few examples here, for details, please, look at the talks directly and also at the 'POWER report' ! ([www.ippp.dur.ac.uk/~gudrid/power/report.pdf](http://www.ippp.dur.ac.uk/~gudrid/power/report.pdf))

# Physics at the $e^+e^-$ Linear Collider

## \* Discovery of New Physics (NP)

- Large potential for direct searches
- Impressive potential also for indirect searches!

## \* Unraveling the structure of NP

- precise determination of underlying dynamics and parameters
- model distinction through model-independent searches

## \* High precision measurements

- tests of the SM with unprecedented precision
- even smallest hints of NP could be observed

⇒ **Beam polarization = decisive tool for direct and indirect searches!**

**'State of the art':**

Polarized  $e^-$  beam at SLAC: SLC  $\sim 75\%$   
E158  $\sim 90\%$   
at Nagoya, KEK:  $\sim 90\%$

**new results** show that  $P(e^-) \sim 90\%$  can be expected at ILC!

⇒ won't such high  $P(e^-)$  suffice?

## Polarization report - 'POWER Write-Up'

- **The Physics case for having both beams polarized:**  
140 pages, ~ 70 authors, ~ 30 institutes  
→ incl. 80 pages physics, 20 pages machine, 20 pages polarimetry  
→ will be submitted to Physics Reports and to hep-ph (very soon!)  
→ <http://www.ippp.dur.ac.uk/~gudrid/power/report.pdf>
- **News from physics** with polarized beams in Susy, SM, other NP!  
→ focus on use of  $P_{e^+}$  compared to  $P_{e^-}$  only
- **Machine overview** about **polarized  $e^+$  source**  
and **polarization measurements**
- **Thanks a lot to all authors!**  
**Very special thanks to**  
**S. Riemann,**  
**P. Osland and N. Paver,**  
**R. Pitthan, P. Schüler, J. Sheppard, M. Woods!**

Final Draft

# Revealing fundamental interactions: the rôle of polarized positrons and electrons at the Linear Collider\*

G. MOORTGAT-PICK<sup>1,10</sup>, T. ABE<sup>13</sup>, G. ALEXANDER<sup>23</sup>, B. ANANTHANARAYAN<sup>2</sup>, A.A. BABICH<sup>3</sup>,  
V. BHARADWAJ<sup>6</sup>, D. BARBER<sup>7</sup>, A. BARTL<sup>4</sup>, A. BRACHMANN<sup>6</sup>, S. CHEN<sup>13</sup>, J. CLARKE<sup>5</sup>,  
J.E. CLENDENIN<sup>6</sup>, J. DANTON<sup>21</sup>, K. DESCH<sup>29</sup>, M. DIEHL<sup>7</sup>, B. DOBOS<sup>13</sup>, T. DORLAND<sup>13</sup>,  
H. EBERL<sup>8</sup>, J.R. ELLIS<sup>1</sup>, K. FLOETTMANN<sup>7</sup>, H. FRAAS<sup>9</sup>, F. FRANCO-SOLLOVA<sup>7</sup>, F. FRANKE<sup>9</sup>,  
A. FREITAS<sup>31</sup>, J. GOODSON<sup>13</sup>, J. GRAY<sup>13</sup>, A. HAN<sup>13</sup>, S. HEINEMEYER<sup>1</sup>, S. HESSELBACH<sup>4,9</sup>,  
T. HIROSE<sup>28</sup>, K. HOHENWARTER-SODEK<sup>4</sup>, J. KALINOWSKI<sup>24</sup>, T. KERNREITER<sup>4</sup>, O. KITTEL<sup>9</sup>,  
S. KRAML<sup>1</sup>, W. MAJEROTTO<sup>8</sup>, A. MARTINEZ<sup>13</sup>, H.-U. MARTYN<sup>30</sup>, A. MIKHAILICHENKO<sup>25</sup>,  
W. MENGES<sup>7</sup>, K. MOENIG<sup>7</sup>, K. MOFFEIT<sup>6</sup>, S. MORETTI<sup>11</sup>, O. NACHTMANN<sup>12</sup>, F. NAGEL<sup>12</sup>,  
T. NAKANISHI<sup>15</sup>, U. NAUENBERG<sup>13</sup>, T. OMORI<sup>15</sup>, P. OSLAND<sup>1,14</sup>, A.A. PANKOV<sup>3</sup>, N. PAVER<sup>16</sup>,  
R. PITTHAN<sup>6</sup>, W. POROD<sup>17,27</sup>, J. PROULX<sup>13</sup>, P. RICHARDSON<sup>10</sup>, S. RIEMANN<sup>7</sup>, S.D. RINDANI<sup>18</sup>,  
T.G. RIZZO<sup>6</sup>, P. SCHÜLER<sup>7</sup>, C. SCHWANENBERGER<sup>19</sup>, D. SCOTT<sup>5</sup>, J. SHEPPARD<sup>6</sup>, R.K. SINGH<sup>2</sup>,  
A. STAHL<sup>22</sup>, H. STEINER<sup>26</sup>, A. WAGNER<sup>9</sup>, G. WEIGLEIN<sup>10</sup>, M. WOODS<sup>6</sup>, P. ZERWAS<sup>7</sup>,  
F. ZOMER<sup>20</sup>

# Outline of the report

## a) Introduction:

- ★ possible general dependences on beam polarization w.r.t. kind of interaction
- ★ definitions and gain in accuracy for  $A_{LR}$  measurement with  $P_{e^+}$

## b) Open questions of the SM: top, Higgs, GigaZ

- ★  $t$  and  $H$  couplings and properties
- ★ application of Blondel scheme for high precision tests

## c) Searches for New Physics: Susy, CI, ED, LQ, new CP-violation,...

- ★ parameter determination (many!), CP-violating effects, background supp.
- ★ model-independent approaches in direct and indirect searches

## d) Summary of the Physics cases

- ★ qualitative and quantitative improvement factors listed in short summaries
- ★ summary tables for longitudinally and transversely polarized beams

## e) Technical aspects:

- ★ history of polarized  $e^-$  at SLC; polarized  $e^-$  source design for ILC
- ★ polarized  $e^+$ : undulator-based schemes, comments on laser-based scheme
- ★ polarization measurement via up-/downstream polarimetry, annihilation data


# Some physics examples – quick rush through

## Supersymmetry: Test of quantum numbers


Association of chiral electrons to scalar partners  $e_{L,R}^- \leftrightarrow \tilde{e}_{L,R}^-$  and  $e_{L,R}^+ \leftrightarrow \tilde{e}_{R,L}^+$

1. separation of scattering (direct SM→SUSY vertex) ↔ annihilation channel

2. test of 'chirality': **only**  $\tilde{e}_R^- \tilde{e}_L^+$  may survive at  $P_{e^-} > 0$  and  $P_{e^+} > 0$ !



high  $P_{e^-} > 0$ !



high  $P_{e^-} > 0$  and  $P_{e^+} > 0$ !


⇒ Even high  $P_{e^-}$  not sufficient but  $P_{e^+}$  needed!

# Some physics examples – quick rush through

## Supersymmetry: Test of Yukawa couplings


Test of SU(2), U(1) gauge couplings  $\equiv$  SUSY Yukawa couplings

1. separation of the pairs  $\tilde{e}_R^- \tilde{e}_R^+$  and  $\tilde{e}_R^- \tilde{e}_L^+$
2. 'variation' of Yukawa couplings accepted within experimental uncertainty



$e^+$  Yukawa couplings: only  $P_{e^-}$

$\Rightarrow$  SU(2), U(1) Yukawa coupling 'not' measurable



$P_{e^-}$  and  $P_{e^+}$

$\Rightarrow \Delta$  SU(2)  $\sim$  80%,  $\Delta$  U(1)  $\sim$  2.5%

*Freitas*

$\Rightarrow$  Even high  $P_{e^-}$  not sufficient but  $P_{e^+}$  needed!




# Some physics examples – quick rush through

## Searches for CP-violation, e.g. in SUSY (many new phases)

### 'Construction' of T-odd asymmetries with angular correlations


1. small phases  $\rightarrow$  small asymmetries expected

$$\sigma(e^+e^- \rightarrow \tilde{\chi}_1^0 \tilde{\chi}_2^0 \rightarrow 2\tilde{\chi}_1^0 \ell^- \ell^+)$$



$$A_T(e^+e^- \rightarrow \tilde{\chi}_1^0 \tilde{\chi}_2^0 \rightarrow 2\tilde{\chi}_1^0 \ell^- \ell^+)$$

$\phi_\mu = 0, \text{ small } \phi_{M_1}$



- with longitudinally pol. beams:

$$P_{e^+} \text{ needed for better measurability of asymmetry } \sim |A_T| \sqrt{\mathcal{L} \times \sigma}$$

- with transversely polarized beams: direct access to CP-odd asymmetries

$\Rightarrow$  Both  $e^-$  and  $e^+$  beams have to be polarized!

# Some physics examples – quick rush through

## Transversely polarized beams also for indirect searches:

- Who guaranties that we will ever reach the new heavy scale?
    - indirect searches important!
    - however, strong model-dependence!
  - With transversely polarized beams:
    - exploit **azimuthal asymmetries** also for indirect searches!
    - a) **distinction** between SM and different models of large extra dimensions *Rizzo*
    - b) access to new CP-violating kind of interactions in  $t\bar{t}, \gamma Z, W^+W^-$  *Nagel*  
*Rindani*
      - **unique** access to  $\Re$  parts of CP-sensitive couplings!
- ⇒ Transversely polarized beams are very effective also for indirect seaches w/wo CP-violation
- in principle both  $e^-$  and  $e^+$  beams polarized required!

# Some physics examples – quick rush through

Which effects are possible?  $|M|^2 \sim \bar{v}(\lambda_{e^+})\Gamma u(\lambda_{e^-})\bar{u}(\lambda'_{e^-})\Gamma^\dagger v(\lambda'_{e^+})$

Interaction structure		Longitudinal		Transverse	
$\Gamma$	$\Gamma^\dagger$	Bilinear	Linear	Bilinear	Linear
<b>S</b>	<b>S</b>	$\sim P_{e^-}P_{e^+}$	–	$\sim P_{e^-}^T P_{e^+}^T$	–
<b>P</b>	<b>S</b>	–	$\sim P_{e^\pm}$	$\sim P_{e^-}^T P_{e^+}^T$	–
<b>V,A</b>	<b>S</b>	–	–	–	$\sim P_{e^\pm}^T$
<b>T</b>	<b>S</b>	$\sim P_{e^-}P_{e^+}$	$\sim P_{e^\pm}$	$\sim P_{e^-}^T P_{e^+}^T$	
<b>P</b>	<b>P</b>	$\sim P_{e^-}P_{e^+}$	–	$\sim P_{e^-}^T P_{e^+}^T$	–
<b>V,A</b>	<b>P</b>	$\sim P_{e^-}P_{e^+}$	$\sim P_{e^\pm}$	$\sim P_{e^-}^T P_{e^+}^T$	$\sim P_{e^\pm}^T$
<b>T</b>	<b>P</b>	$\sim P_{e^-}P_{e^+}$	$\sim P_{e^\pm}$	$\sim P_{e^-}^T P_{e^+}^T$	
<b>V,A</b>	<b>V,A</b>	$\sim P_{e^-}P_{e^+}$	$\sim P_{e^\pm}$	$\sim P_{e^-}^T P_{e^+}^T$	–
<b>T</b>	<b>V,A</b>	–	–	–	$\sim P_{e^\pm}^T$
<b>T</b>	<b>T</b>	$\sim P_{e^-}P_{e^+}$	$\sim P_{e^\pm}$	$\sim P_{e^-}^T P_{e^+}^T$	–

$P, S = (\text{pseudo})\text{scalar}$

$A, V = (\text{axial})\text{vector}$

$T = \text{tensor}$

⇒ impact of beam polarization depends on kind of interaction(s)

- with  $P_{e^-}$  and  $P_{e^+}$  much higher flexibility with regard to NP candidates for direct as well as indirect searches!

# Technical aspects of polarizing $e^-$ and $e^+$ at the ILC

- Polarized electron source: strained photocathode technology

⇒  $P(e^-) \approx 90\%$  expected

- Positron sources under discussion for the ILC:

- \* Conventional source

⇒ no polarization

- \* Undulator-based source

⇒ polarized, > 150 GeV electrons needed

- Other schemes? Laser-based source

- Status of polarimetry

⇒ Compton polarimetry well advanced

- ★ up-, as well as downstream ILC designs under work

- ★ high accuracy expected  $\Delta P/P \sim 0.25\%$ !

better precision requires  $P_{e^+}$ !

## Results of the report

- clear physics case for polarized  $e^-$  and  $e^+$ 
  - ★ many  $\equiv (n + 1)$  examples from **different** physics scenarios!
  - ★  $P_{e^+}$  always advantageous, independent of direction of NP
  - ★ more observables, higher flexibility, better statistics+systematics

$\Rightarrow P_{e^+}$  crucial preparation for 'being prepared for the Unexpected'!
- ILC scheme designs well on track for  $P_{e^-} = \pm 90\%$  and  $P_{e^+} = \pm 60\%$ 
  - + **kicker system** for pulse-train-to-pulse-train  $\rightarrow$  two IR's with  $P_{e^\pm}$
  - $\Rightarrow$  important also for switching  $\pm P$
- possible to provide polarized beams without loss of peak luminosity and without any critical impact on commissioning!
  - ★ long undulator, additional polarized  $e^-$  source for commissioning,...

$\Rightarrow$  details have to be decided before the final ILC design!
- designs for up-/and downstream polarimetry for the ILC
  - ★ accuracy of  $\Delta P_{e^\pm}/P_{e^\pm} \sim 0.25\%$

$\Rightarrow$  Report should be seen as contemporary status report!  
*still studies ongoing, new ideas+examples coming up*  
however, the final source design should be decided soon!

# Future events related to 'Beam Polarization at the ILC'

- 'Workshop on Positron Sources for the ILC', Daresbury, April, 11-13!

- ★ Program: Polarized and unpolarized  $e^+$  source
- ★ Drive beams, target issues, polarimetry, capture issues, operational aspects
- ★ webpage: [www.astec.ac.uk/id\\_mag/workshop.htm](http://www.astec.ac.uk/id_mag/workshop.htm)

- ongoing experiment at KEK (since 2001)

- ★ polarized  $e^+$  from laser-compton scheme with  $P_{e^+} \sim 80\%$  measured via transmission polarimetry

- First run of the project 'E166' at SLAC in May, 2005

⇒ first results hopefully soon!

- Helical undulator prototypes for ILC parameters in 2005

- ★ Test of a SC helical undulator at Rutherford Lab already started
- ★ Construction of a PPM helical undulator at Daresbury Lab started

- Snowmass in August 2005

Polarization session needed: detailed design comparison between conventional and undulator-based scheme, also news for laser-based scheme planned

## Summary: polarized $e^-$ and $e^+$ at the ILC

- With  $P_{e^+}$  **only** gains, independent in which direction NP points
  - ★ key additional observables for unraveling the underlying physics: kind of interaction, particle properties, parameter determination, . . .
  - ★ significant improvement for model-independent approaches in direct as well as indirect searches for NP
  - ★ Analyzing NP might be challenging → best of all tools needed!
- ⇒ **full potential of the ILC could only be realized with  $P_{e^-}$  and  $P_{e^+}$ !**
- **undulator-based scheme is feasible and its design well advanced**
  - ★ **some prototypes** for ILC design already under construction
- **To-do list:** detailed designs for both conventional and undulator-based source? revisit reliability issues? cost comparison? start with conventional? polarized source as upgrade? (expect  $\sim 3\%$  increment in ILC project cost to have both conventional and undulator-based sources)

⇒ **Please, think about these questions now, i.e. in 2005, . . . !**  
 **$P_{e^-}$  only together with  $P_{e^+}$  provide a unique tool for the ILC:**

- ★ high potential of applications
- ★ and preparation for the 'Unexpected!'