

Spin Dynamics and Polarization
Cockcroft Institute, June 5 and 6, 2006
Gudrid Moortgat-Pick

Transformation: Two 2-component Dirac equations in one 4-component Dirac equation

To avoid too much matrix algebra, we show the equality of the two Dirac equations, page 19, with the four-dim denotation Dirac equation, also page 19, in this way:

$$|\sigma, p, 1 \rangle = \left[\frac{p_0}{m} - \frac{\vec{p}\vec{\sigma}}{m} \right] |\sigma, p, 2 \rangle \quad (1)$$

write as matrix for both solution of $\sigma = \pm \frac{1}{2}$:

$$\begin{pmatrix} |+\frac{1}{2}, p, 1 \rangle \\ |-\frac{1}{2}, p, 1 \rangle \end{pmatrix} = \frac{1}{m} \begin{pmatrix} (p_0 - p_3) & -(p_1 - ip_2) \\ -(p_1 + ip_2) & (p_0 + p_3) \end{pmatrix} \begin{pmatrix} |+\frac{1}{2}, p, 2 \rangle \\ |-\frac{1}{2}, p, 2 \rangle \end{pmatrix} \quad (2)$$

\Rightarrow

$$|+\frac{1}{2}, p, 1 \rangle = \frac{1}{m}(p_0 - p_3)|+\frac{1}{2}, p, 2 \rangle + \frac{1}{m}(-p_1 + ip_2)|-\frac{1}{2}, p, 2 \rangle \quad (3)$$

$$|-\frac{1}{2}, p, 1 \rangle = \frac{1}{m}(-p_1 - ip_2)|+\frac{1}{2}, p, 2 \rangle + \frac{1}{m}(p_0 + p_3)|-\frac{1}{2}, p, 2 \rangle \quad (4)$$

Analogous procedure with the second 2-component Dirac equation:

$$|\sigma, p, 2 \rangle = \left[\frac{p_0}{m} + \frac{\vec{p}\vec{\sigma}}{m} \right] |\sigma, p, 1 \rangle \quad (5)$$

Write it also explicitly as matrix:

$$\begin{pmatrix} |+\frac{1}{2}, p, 2 \rangle \\ |-\frac{1}{2}, p, 2 \rangle \end{pmatrix} = \frac{1}{m} \begin{pmatrix} (p_0 + p_3) & +(p_1 - ip_2) \\ (p_1 + ip_2) & (p_0 - p_3) \end{pmatrix} \begin{pmatrix} |+\frac{1}{2}, p, 1 \rangle \\ |-\frac{1}{2}, p, 1 \rangle \end{pmatrix} \quad (6)$$

\Rightarrow

$$|+\frac{1}{2}, p, 2 \rangle = \frac{1}{m}(p_0 + p_3)|+\frac{1}{2}, p, 1 \rangle + \frac{1}{m}(p_1 - ip_2)|-\frac{1}{2}, p, 1 \rangle \quad (7)$$

$$|-\frac{1}{2}, p, 2 \rangle = \frac{1}{m}(p_1 + ip_2)|+\frac{1}{2}, p, 1 \rangle + \frac{1}{m}(p_0 - p_3)|-\frac{1}{2}, p, 1 \rangle \quad (8)$$

Now, deriving the 4-dim denotation, page 19. First, the definition of the used γ matrices:

$$\gamma_0 = \begin{pmatrix} 0 & I \\ I & 0 \end{pmatrix}, \quad \gamma_i = \begin{pmatrix} 0 & -\sigma_j \\ +\sigma_j & 0 \end{pmatrix}, \quad j = 1, 2, 3, \quad (9)$$

where I denotes the 2×2 unit-matrix and σ^j the Pauli-matrices, page 8.

Keeping track of the order of the indices, eqn.(1) and (5) become:

$$(\gamma_\mu^\dagger p_\mu - m)_{\alpha'\alpha} |\sigma, p, \alpha'\rangle = 0 \quad (10)$$

More explicitly:

$$\left\{ \begin{array}{l} \left(\begin{array}{cccc} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{array} \right)^\dagger p_0 + \left(\begin{array}{cccc} 0 & 0 & 0 & -1 \\ 0 & 0 & -1 & 0 \\ 0 & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{array} \right)^\dagger p_1 + \left(\begin{array}{cccc} 0 & 0 & 0 & i \\ 0 & 0 & -i & 0 \\ 0 & -i & 0 & 0 \\ i & 0 & 0 & 0 \end{array} \right)^\dagger p_2 + \left(\begin{array}{cccc} 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \end{array} \right)^\dagger p_3 - m I_{4 \times 4} \end{array} \right\} \begin{pmatrix} |p1\rangle \\ |p2\rangle \\ |p3\rangle \\ |p4\rangle \end{pmatrix} = 0 \quad (11)$$

$$\Rightarrow \begin{pmatrix} -m & 0 & p_0 + p_3 & p_1 - ip_2 \\ 0 & -m & p_1 + ip_2 & p_0 - p_3 \\ p_0 - p_3 & -p_1 + ip_2 & -m & 0 \\ -p_1 - ip_2 & p_0 + p_3 & 0 & -m \end{pmatrix} \begin{pmatrix} |+\frac{1}{2}, p, 2\rangle \\ |-\frac{1}{2}, p, 2\rangle \\ |+\frac{1}{2}, p, 1\rangle \\ |-\frac{1}{2}, p, 1\rangle \end{pmatrix} = 0 \quad (12)$$

This represents the following system of equations:

$$|+\frac{1}{2}, p, 2\rangle = \frac{1}{m}(p_0 + p_3)|+\frac{1}{2}, p, 1\rangle + \frac{1}{m}(p_1 - ip_2)|-\frac{1}{2}, p, 1\rangle \quad (13)$$

$$|-\frac{1}{2}, p, 2\rangle = \frac{1}{m}(p_1 + ip_2)|+\frac{1}{2}, p, 1\rangle + \frac{1}{m}(p_0 - p_3)|-\frac{1}{2}, p, 1\rangle \quad (14)$$

$$|+\frac{1}{2}, p, 1\rangle = \frac{1}{m}(p_0 - p_3)|+\frac{1}{2}, p, 2\rangle + \frac{1}{m}(-p_1 + ip_2)|-\frac{1}{2}, p, 2\rangle \quad (15)$$

$$|-\frac{1}{2}, p, 1\rangle = \frac{1}{m}(-p_1 - ip_2)|+\frac{1}{2}, p, 2\rangle + \frac{1}{m}(p_0 + p_3)|-\frac{1}{2}, p, 2\rangle \quad (16)$$

That's identical to Eqn.(7), (8), (3), (4)

Last step is only the definition of the wave functions:

$$u_\alpha(p) = \langle p\alpha | u \rangle \quad (17)$$

Inserting this definition into Eq.(10) gives the Dirac equation in 4-dim denotation, page 19 (we have to do a hermitian conjugation, this removes the \dagger in Eq.(10)):

$$(\gamma_\mu p_\mu - m)_{\alpha'\alpha} u_\alpha(p) = 0 \quad (18)$$