LC Forum 2013 – Latest News from WHIZARD

Jürgen R. Reuter

DESY Hamburg





LC Forum 2013, DESY, Oct 11th, 2013

The WHIZARD Event Generator – Release 2.1

- Multi-Channel Monte-Carlo integration
- Efficient phase space and event generation (weighted & unweighted)
- Optimized tree-level matrix elements (O'Mega)
  - $e^+e^- \to t\bar{t}H \to b\bar{b}b\bar{b}jj\ell\nu$ (110,000 diagrams)
  - $e^+e^- \to ZHH \to ZWWWW \to bb + 8j$ (12,000,000 diagrams)
  - $pp \to \ell\ell + nj,\ n = 0, 1, 2, 3, 4, \ldots$ (2,100,000 diagrams with 4 jets + flavors)
  - $pp \to \tilde{\chi}_1^0\tilde{\chi}_1^0 bbbb$ (32,000 diagrams, 22 color flows, $\sim 10,000$ PS channels)
  - $pp \to VVjj \to jj\ell\ell\nu\nu$ incl. anomalous TGC/QGC
  - Test case $gg \to 9g$ (224,000,000 diagrams)

WHIZARD 2.1.1 release: Sep. 18, 2012

Old series: WHIZARD 1.97 (development stopped with 1.94)

The WHIZARD team: F. Bach, [H. Boschmann], [F. Braam], B. Chokoufé, W. Kilian, T. Ohl, JRR, [S. Schmidt], [S. Schwertfeger], M. Sekulla, [C. Speckner], F. Staub, [M. Trudewind], C. Weiss, [D. Wiesler]

Web address: `http://projects.hepforge.org/whizard`

Standard Reference: Kilian/Ohl/JRR, EPJC 71 (2011) 1742, arXive:0708.4233

The WHIZARD Event Generator – Release 2.2

- Multi-Channel Monte-Carlo integration
- Efficient phase space and event generation (weighted & unweighted)
- Optimized tree-level matrix elements (O'Mega)
  - $e^+e^- \to t\bar{t}H \to b\bar{b}b\bar{b}jj\ell\nu$ (110,000 diagrams)
  - $e^+e^- \to ZHH \to ZWWWW \to bb + 8j$ (12,000,000 diagrams)
  - $pp \to \ell\ell + nj,\ n = 0, 1, 2, 3, 4, \ldots$ (2,100,000 diagrams with 4 jets + flavors)
  - $pp \to \tilde{\chi}_1^0\tilde{\chi}_1^0 bbbb$ (32,000 diagrams, 22 color flows, $\sim 10,000$ PS channels)
  - $pp \to VVjj \to jj\ell\ell\nu\nu$ incl. anomalous TGC/QGC
  - Test case $gg \to 9g$ (224,000,000 diagrams)

WHIZARD 2.2.0_α-2 release: Oct. 2nd, 2013

Old series: WHIZARD 1.97 (development stopped with 1.94)

The WHIZARD team: F. Bach, [H. Boschmann], [F. Braam], B. Chokoufé, W. Kilian, T. Ohl, JRR, [S. Schmidt], [S. Schwertfeger], M. Sekulla, [C. Speckner], F. Staub, [M. Trudewind], C. Weiss, [D. Wiesler]

Web address: `http://projects.hepforge.org/whizard`

Standard Reference: Kilian/Ohl/JRR, EPJC 71 (2011) 1742, arXive:0708.4233

The WHIZARD Event Generator – Release 2.2

- Multi-Channel Monte-Carlo integration
- Efficient phase space and event generation (weighted & unweighted)
- Optimized tree-level matrix elements (O'Mega)
  - $e^+e^- \to t\bar{t}H \to b\bar{b}b\bar{b}jj\ell\nu$ (110,000 diagrams)
  - $e^+e^- \to ZHH \to ZWWWW \to bb + 8j$ (12,000,000 diagrams)
  - $pp \to \ell\ell + nj,\ n = 0, 1, 2, 3, 4, \ldots$ (2,100,000 diagrams with 4 jets + flavors)
  - $pp \to \tilde{\chi}_1^0\tilde{\chi}_1^0 bbbb$ (32,000 diagrams, 22 color flows, $\sim 10,000$ PS channels)
  - $pp \to VVjj \to jj\ell\ell\nu\nu$ incl. anomalous TGC/QGC
  - Test case $gg \to 9g$ (224,000,000 diagrams)

WHIZARD 2.2.0 release: Nov. 11, 2013 (LCWS)

Old series: WHIZARD 1.97 (development stopped with 1.94)

The WHIZARD team: F. Bach, [H. Boschmann], [F. Braam], B. Chokoufé, W. Kilian, T. Ohl, JRR, [S. Schmidt], [S. Schwertfeger], M. Sekulla, [C. Speckner], F. Staub, [M. Trudewind], C. Weiss, [D. Wiesler]

Web address: `http://projects.hepforge.org/whizard`

Standard Reference: Kilian/Ohl/JRR, EPJC 71 (2011) 1742, arXive:0708.4233

The WHIZARD Event Generator – Release 2.2

- Multi-Channel Monte-Carlo integration
- Efficient phase space and event generation (weighted & unweighted)
- Optimized tree-level matrix elements (O'Mega)
  - $e^+e^- \to t\bar{t}H \to b\bar{b}b\bar{b}jj\ell\nu$ (110,000 diagrams)
  - $e^+e^- \to ZHH \to ZWWWW \to bb + 8j$ (12,000,000 diagrams)
  - $pp \to \ell\ell + nj,\ n = 0, 1, 2, 3, 4, \ldots$ (2,100,000 diagrams with 4 jets + flavors)
  - $pp \to \tilde{\chi}_1^0\tilde{\chi}_1^0 bbbb$ (32,000 diagrams, 22 color flows, $\sim 10,000$ PS channels)
  - $pp \to VVjj \to jj\ell\ell\nu\nu$ incl. anomalous TGC/QGC
  - Test case $gg \to 9g$ (224,000,000 diagrams)

WHIZARD 2.2.0 release: Nov. 11, 2013 (LCWS)

Old series: WHIZARD 1.97 (development stopped with 1.94)

The WHIZARD team: F. Bach, [H. Boschmann], [F. Braam], B. Chokoufé, W. Kilian, T. Ohl, JRR, [S. Schmidt], [S. Schwertfeger], M. Sekulla, [C. Speckner], F. Staub, [M. Trudewind], C. Weiss, [D. Wiesler] + 3 bachelors

Web address: `http://projects.hepforge.org/whizard`

Standard Reference: Kilian/Ohl/JRR, EPJC 71 (2011) 1742, arXive:0708.4233

WHIZARD 2: Status 2011/12 – Technical Features

- WHIZARD 2: code basically rewritten, only `Fortran 2003` and `O'Caml`
- Clean modularization of code/(First) object-oriented implementation
- OpenMP parallelization
- Operation modes:
  - Dynamic linking (default mode) with on-the-fly generation of process code
  - Static linking (for batch clusters)
  - Library mode, callable from C/C++/Python/...
  - Interactive mode: WHIZARD works as a Shell – WHISH
- Standard conformance: uses `autotools`: `automake`/`autoconf`/`libtool`
- Large self test suite
- Version control (`svn`) at HepForge: use of ticket system and bug tracker
- Continuous integration system (`jenkins`) linked with svn repository

WHIZARD 2 – Installation and Run

- Download WHIZARD from `http://www.hepforge.org/archive/whizard/whizard-2.1.1.tar.gz` and unpack it
- WHIZARD intended to be centrally installed on a system, e.g. in `/usr/local` (or locally on user account)
- Create build directory and `configure`
  External programs (LHAPDF, StdHEP, HepMC) might need flags
- `make`, `make install`
- Create SINDARIN steering file (in any working directory)
- Run `whizard` (in working directory)
- Supported event formats: HepMC, StdHEP, LHEF, LHA, div. ASCII formats

```
O'Mega self tests:
make  check-TESTS
PASS: test_omega95
PASS: test_omega95_bispinors
PASS: test_qed_eemm
PASS: ects
PASS: ward
PASS: compare_split_function
PASS: compare_split_module
==================
All 7 tests passed
==================
WHIZARD self tests:
make  check-am
make  check-TESTS
PASS: empty.run
PASS: vars.run
PASS: md5.run
[..........]
XFAIL: errors.run
PASS: extpar.run
PASS: susyhit.run
PASS: libs.run
PASS: qedtest.run
PASS: helicity.run
PASS: smtest.run
PASS: defaultcuts.run
PASS: restrictions.run
PASS: decays.run
PASS: alphas.run
PASS: colors.run
PASS: cuts.run
PASS: lhapdf.run
PASS: ilc.run
PASS: mssmtest.run
PASS: models.run
PASS: stdhep.run
PASS: stdhep_up.run
====================================
All 53 tests behaved as expected (1 e
====================================
```

WHIZARD Manual



whizard.hepforge.org/manual/

Google

Meistbesucht | Aktuelle Nachr... | WHIZARD | INSPIRE | arXiv.org e-Print... | YouTube - Broa... | Schlagzeilen - S...

WHIZARD is

- Home
- Downloads
- Wiki
- News
- ChangeLog
- Subversion
- Browser
- Tracker
- Internal

WHIZARD 2.1
A generic
Monte-Carlo integration and event generation package
for multi-particle processes
MANUAL [1]

Wolfgang Kilian,[2] Thorsten Ohl,[3] Jürgen Reuter,[4] Christian Speckner [5]

- Contents
- Introduction
  - Disclaimer
  - Overview
  - About examples in this manual
- Installation
  - Package Structure
  - Prerequisites
  - Installation
  - Working With WHIZARD
- Getting Started
  - Hello World
  - A Simple Calculation
- SINDARIN: Overview
  - The command language for WHIZARD
  - SINDARIN scripts
  - Errors
  - Statements
  - Control Structures
  - Expressions
  - Variables

Physics aspects/improvements in WHIZARD 2

- SINDARIN (Scripting INtegration, Data Analysis, Results display and INterfaces) allows for arbitrary expressions for cuts and scales etc. (examples later)

```
cuts = any 5 degree < Theta < 175 degree
         [select if abs (Eta) < eta_cut [lepton]]
cuts = any E > 2 * mW [extract index 2
                         [sort by Pt [lepton]]]
```

- New syntax for decays and chains:

```
process higgsstr = e1, E1 => (Z => e2, E2), (H => b, bbar)
process wtf      = e1, E1 => (Z, h) + (Z, H) + (A, H)
```

- Process libraries: processes of different BSM models can be used in parallel
- Decay cascades including full spin correlations (cf. later)
- `FeynRules` interface Christensen/Duhr/Fuks/JRR/Speckner, EPJC 72 (2012) 1990
- MLM jet matching
- Event-dependent scales in PDFs and running α_s
- Parton Shower: p_T-ordered and analytic Kilian/JRR/Schmidt/Wiesler, JHEP 1204 (2012) 013

Structured Beams

- **Lepton Colliders structured beams**
  - QED ISR (Skrzypek/Jadach, Kuraev/Fadin , incl. p_T distributions)
  - arbitrarily polarized beams (density matrices)
  - Beamstrahlung (CIRCE module) **more later**
  - Photon collider spectra (CIRCE2 module)
  - external beam spectra can be read in (files/generating code)
  - QED FSR (e.g. YFS) not (yet) implemented (charged mesons/hadrons)
- **Hadronic events/hadronic decays**
  - through PYTHIA interface [or HERWIG]
- **Hadron Colliders structured beams**
  - LHAPDF interface
  - Most prominent PDFs directly included
  - QCD ISR and FSR (two different own implementations, interface to PYTHIA)
  - Matching matrix elements/showers (MLM)
  - Underlying event/multiple interactions

O'Mega: Optimal matrix elements

Ohl/JRR, 2001

- [.·.] Replace forest of tree diagrams by
  Directed Acyclical Graph (DAG) of the algebraic expression (including color).



O'Mega: Optimal matrix elements

Ohl/JRR, 2001

- [.·.] Replace forest of tree diagrams by
  Directed Acyclical Graph (DAG) of the algebraic expression (including color).



- simplest examples: $e^+e^- \to \mu^+\mu^-$, and



O'Mega: Optimal matrix elements

Ohl/JRR, 2001

- [.·.] Replace forest of tree diagrams by
  Directed Acyclical Graph (DAG) of the algebraic expression (including color).



- simplest examples: $e^+e^- \to \mu^+\mu^-$, $e^+e^- \to \mu^+\mu^-\gamma$ and





O'Mega: Optimal matrix elements

Ohl/JRR, 2001

- [.·.] Replace forest of tree diagrams by
  Directed Acyclical Graph (DAG) of the algebraic expression (including color).



- simplest examples: $e^+e^- \to \mu^+\mu^-$, $e^+e^- \to \mu^+\mu^-\gamma$ and $e^+e^- \to \mu^+\mu^-\gamma\gamma$







Hard matrix elements: particle types

Possible particle types

- Spin 0 particles
- Spin 1/2 fermions (Majorana and Dirac)
  Fermi statistics for both fermion-number conserving and violating cases
- Spin 1 particles
  - massive and massless
  - Unitarity and Feynman gauge
  - arbitrary R_ξ gauges
- Spin 3/2 particles (Majorana only, gravitinos)
- Spin 2 particles (massless and massive, gravitons)
- Dynamic particles vs. pure insertions
- Unphysical particles for Ward- and Slavnov-Taylor identities

Hard matrix elements: Lorentz structures

Hard-coded set of Lorentz structures

- Purely scalar couplings:
  $\phi^3, \quad \phi^4$
- Scalar couplings to vectors:
  $gV^\mu \phi_1 \mathrm{i}\overleftrightarrow{\partial_\mu} \phi_2, \quad \phi V^2, \quad \phi^2 V^2, \quad \frac{1}{2}\phi F_{1,\mu\nu} F_2^{\mu\nu}, \quad \frac{1}{2}\phi F_{1,\mu\nu} \tilde{F}_2^{\mu\nu}, \quad \phi(\mathrm{i}\partial_\mu V_1^\nu)(\mathrm{i}\partial_\nu V_2^\mu)$
- Pure vector couplings:
  $F_{\mu\nu}F^{\mu\nu}, \quad V_1^\mu((\mathrm{i}\partial_\nu V_2^\rho)\mathrm{i}\overleftrightarrow{\partial_\mu}(\mathrm{i}\partial_\rho V_3^\nu)), \quad gF_1^{\mu\nu}F_{2,\nu\rho}F_{3,}{}^{\rho}{}_{\mu},$
  $g/2 \cdot \epsilon^{\mu\nu\lambda\tau} F_{1,\mu\nu} F_{2,\tau\rho} F_{3,}{}^{\rho}{}_{\lambda}$
- Fermionic couplings to scalars:
  $g_S \bar{\psi}_1 S \psi_2, \quad g_P \bar{\psi}_1 P \gamma_5 \psi_2, \quad \bar{\psi}_1 \phi (g_S + g_P \gamma_5)\psi_2, \quad g_L \bar{\psi}_1 \phi (1-\gamma_5)\psi_2,$
  $g_R \bar{\psi}_1 \phi (1+\gamma_5)\psi_2, \quad g_L \bar{\psi}_1 \phi (1-\gamma_5)\psi_2 + g_R \bar{\psi}_1 \phi (1+\gamma_5)\psi_2$
- Fermionic couplings to vectors:
  $g_V \bar{\psi}_1 \not{V} \psi_2, \quad g_A \bar{\psi}_1 \gamma_5 \not{V} \psi_2, \quad \bar{\psi}_1 \not{V} (g_V - g_A \gamma_5)\psi_2, \quad g_L \bar{\psi}_1 \not{V} (1-\gamma_5)\psi_2,$
  $g_R \bar{\psi}_1 \not{V} (1+\gamma_5)\psi_2, \quad g_L \bar{\psi}_1 \not{V} (1-\gamma_5)\psi_2 + g_R \bar{\psi}_1 \not{V} (1+\gamma_5)\psi_2$

- Fermionic couplings in SUSY Ward identities (not listed here)
- Fermionic couplings to tensors:
  $g_T T_{\mu\nu} \bar{\psi}_1 [\gamma^\mu, \gamma^\nu]_- \psi_2$
- Tensor couplings to vectors:
  $T^{\mu\nu}(V_{1,\mu} V_{2,\nu} + V_{1,\nu} V_{2,\mu})$, $T^{\alpha\beta}(V_1^\mu \mathrm{i}\overleftrightarrow{\partial}_\alpha \mathrm{i}\overleftrightarrow{\partial}_\beta V_{2,\mu}$,
  $T^{\alpha\beta}(V_1^\mu \mathrm{i}\overleftrightarrow{\partial}_\beta (\mathrm{i}\partial_\mu V_{2,\alpha}) + V_1^\mu \mathrm{i}\overleftrightarrow{\partial}_\alpha (\mathrm{i}\partial_\mu V_{2,\beta}))$, $T^{\alpha\beta}((\mathrm{i}\partial^\mu V_1^\nu)\mathrm{i}\overleftrightarrow{\partial}_\alpha \mathrm{i}\overleftrightarrow{\partial}_\beta (\mathrm{i}\partial_\nu V_{2,\mu}))$
- Gravitino couplings:
  $\bar{\psi}\gamma^\mu S \psi_\mu$, $\bar{\psi}\gamma^\mu \not{k}_S S \psi_\mu$, $\bar{\psi}\gamma^\mu \gamma^5 P \not{k}_P \psi_\mu$, $\bar{\psi}\gamma^5 \gamma^\mu [\not{k}_V, \not{V}] \psi_\mu$ etc.

  and many more to fill your advent calendar.....

- Completely general Lorentz structures:
  work in progress, to appear in version 2.2

Hard matrix elements: Color structures

Possible Color structures

- All $SU(N)$ gauge theories supported, but specialize to $N = 3$
- Color flow formalism Stelzer/Willenbrock, 2003; Kilian/Ohl/JRR/Speckner, 2011
- Fundamental representations: $\mathbf{3}, \overline{\mathbf{3}}$
- Adjoint representation: $\mathbf{8}$
- Covers all interactions e.g. in SUSY and extra dimensions
- in preparation: generalized color structures with reps. $\mathbf{6}, \overline{\mathbf{6}}, \mathbf{10}, \overline{\mathbf{10}}$ as well as $\epsilon_{ijk}\phi_i\phi_j\phi_k$ couplings to appear in version 2.2.x

WHIZARD histograms



New completely general syntax in WHIZARD 2.x

```
$title = "Jet Energy in $pp\to \ell\bar\nu j$"
$x_label = "$E$/GeV"
histogram e_jet (0 GeV, 80 GeV, 2 GeV)
analysis = record pt_lepton (eval Pt [extract index 1 [sort by Pt [lepton]]]);
           record pt_jet (eval Pt [extract index 1 [sort by Pt [jet]]]);
           record e_lepton (eval E [extract index 1 [sort by Pt [lepton]]]);
           record e_jet (eval E [extract index 1 [sort by Pt [jet]]])
```

WHIZARD – Overview over BSM Models

MODEL TYPE	with CKM matrix	trivial CKM
QED with e, μ, τ, γ	–	`QED`
QCD with d, u, s, c, b, t, g	–	`QCD`
Standard Model	**`SM_CKM`**	**`SM`**
SM with anomalous gauge coupl.	**`SM_ac_CKM`**	**`SM_ac`**
SM with anomalous top coupl.	**`SMtop_CKM`**	**`SMtop`**
SM with K matrix	—	`SM_KM`
MSSM	`MSSM_CKM`	`MSSM`
MSSM with gravitinos	—	`MSSM_Grav`
NMSSM	`NMSSM_CKM`	`NMSSM`
extended SUSY models	—	`PS/E/SSM`
Littlest Higgs	—	`Littlest`
Littlest Higgs with ungauged $U(1)$	—	`Littlest_Eta`
Littlest Higgs with T parity	—	`Littlest_Tpar`
Simplest Little Higgs (anomaly-free)	—	`Simplest`
Simplest Little Higgs (universal)	—	`Simplest_univ`
3-site model	—	`Threeshl`
UED	—	`UED`
SM with Z'	—	`Zprime`
SM with gravitino and photino	—	`GravTest`
Augmentable SM template	—	`Template`

new models easily: FeynRules interface Christensen/Duhr/Fuks/JRR/Speckner, 1010.3251

Interface to SARAH in the SUSY Toolbox Staub, 0909.2863; Ohl/Porod/Speckner/Staub, 1109.5147

Input files: Basic features

```
model = SM


process halloween = E1, e1 => t, tbar, H


compile


sqrts = 500
beams = E1, e1 => circe1 => isr


integrate (susybg) { iterations = 5:10000, 2:10000 }


n_events = 10000


simulate (full) {
}
```

Example: LHC SUSY cascade decays, Input File

```
model = MSSM

process dec_su_q = su1 => u, neu2
process dec_neu_sl2  = neu2 => SE12, e1

process susybg = u,U => SU1, su1
process full = u, U => SU1, u, e1, SE12

compile

?slha_read_decays = true
read_slha("sps1ap_decays.slha")

integrate (dec_su_q, dec_neu_sl2) { iterations = 1:1000 }

sqrts = 14000
beams = p, p => lhapdf

integrate (susybg) { iterations = 5:10000, 2:10000 }
integrate (full)

n_events = 10000

$title = "Full process"
$description =
  "$p + p \to u+ \bar u \to \bar{\tilde u}_1 + u + \tilde e_{12}^+ + e^-$"
$xlabel = "$M_{\rm inv}(ue^-)$"
histogram inv_mass1_full (0,600,20)

simulate (full) {
  $sample = "casc_dec_full"
  analysis =
    record inv_mass1_full (eval M / 1 GeV [combine[u,e1]])
}

compile_analysis
$analysis_filename = "casc_dec"
write_analysis
```

Example: LHC SUSY cascade decays

$$p + p \to \tilde{u}^* + \tilde{u} \to \tilde{u}^* + u + \tilde{e}^+ + e^-$$

- **Full process:**



Example: LHC SUSY cascade decays

$$p + p \to \tilde{u}^* + \tilde{u} \to \tilde{u}^* + u + \tilde{e}^+ + e^-$$

- **Factorized process w/ full spin correlations:**



Example: LHC SUSY cascade decays

$$p + p \to \tilde{u}^* + \tilde{u} \to \tilde{u}^* + u + \tilde{e}^+ + e^-$$

- **Factorized process w/ classical spin correlations:**



Example: LHC SUSY cascade decays

$$p + p \to \tilde{u}^* + \tilde{u} \to \tilde{u}^* + u + \tilde{e}^+ + e^-$$

- **Factorized process w/ no spin correlations:**



Analytic Parton Shower

JRR/Schmidt/Wiesler, JHEP 2012

- Analytic Parton Shower:
  - no shower veto: shower history is exactly known
  - allows reweighting and maybe more reliable error estimate
- new algorithm for initial state QCD radiation





- matching with hard matrix elements, no ”power-shower”

Analytic Parton Shower

JRR/Schmidt/Wiesler, JHEP 2012

- Analytic Parton Shower:
  - no shower veto: shower history is exactly known
  - allows reweighting and maybe more reliable error estimate
- new algorithm for initial state QCD radiation





- matching with hard matrix elements, no ”power-shower”

Status of NLO development in WHIZARD

- BLHA interface: workflow Speckner, 2012
  1. Process definition in SINDARIN $\Rightarrow$ WHIZARD writes contract file
  2. NLO generator generates code, WHIZARD reads contract
  3. NLO matrix element loaded as shared library

- First implementation: interfacing GoSAM and FeynArts

- Automatic generation of dipole subtraction terms Speckner, 2012; JRR/Weiss, 2013/14
  - proof-of-concept code in WHIZARD 2.1
  - implementation in the context of the revised WHIZARD 2.2 core

First example: $u\bar{u} \to \mu^- \bar{\nu}_\mu e^+ \nu_e$

Input:

```
real mreg = 1 GeV

process test = u, ubar => "mu-", numubar, "e+", nue {
   $method = "dipole_integrated_qed"
   soft_mass_regulator = mreg
   collinear_mass_regulators = mreg, mreg, mreg, 0, mreg, 0
}

me = 0
mmu = 0
alpha_qed = 1.  / alpha_em_i

sqrts = 500 GeV

integrate (test) {iterations = 5:10000, 5:20000}
```

Result:

```
| Integrating process 'test':
|=============================================================================|
| It      Calls  Integral[fb]  Error[fb]   Err[%]    Acc  Eff[%]   Chi2 N[It] |
|=============================================================================|
  10     100000  1.9794090E+00  3.16E-03    0.16    0.50   12.33    0.12   5
|=============================================================================|
```

Simulating Linear Colliders

- High-Energy Linear Lepton Collider (250/350/500/1000/2000/3000 GeV)
- ISR, beamstrahlung, strong fields (CLIC)
- Exhaustive support for these effects in WHIZARD (close collaboration with all LC groups)
- Prime Example $e^+e^- \to b\bar{b}$:

Simulating Linear Colliders

- High-Energy Linear Lepton Collider (250/350/500/1000/2000/3000 GeV)
- ISR, beamstrahlung, strong fields (CLIC)
- Exhaustive support for these effects in WHIZARD (close collaboration with all LC groups)
- Prime Example $e^+e^- \to b\bar{b}$:



Simulating Linear Colliders

- High-Energy Linear Lepton Collider (250/350/500/1000/2000/3000 GeV)
- ISR, beamstrahlung, strong fields (CLIC)
- Exhaustive support for these effects in WHIZARD (close collaboration with all LC groups)
- Prime Example $e^+e^- \to b\bar{b}$:



Luminosity spectrum picks up the Z resonance!

News 2013/early 2014: upcoming official release 2.2.x

- Lost of WHIZARD members 2012: some features delayed in 2013

  (pre-)release version 2.2.0_α-2 available!

News 2013/early 2014: upcoming official release 2.2.x

- Lost of WHIZARD members 2012: some features delayed in 2013

  (pre-)release version 2.2.0_α-2 available!

- WHIZARD core: insert an extra abstraction layer, consistently separate interface from implementation Complete object orientation
  - Replaceable modules with well-defined interface: matrix-elements, beam structure, phase space, integration, decays, shower, ...
  - Much easier to contribute new parts to the code $\Rightarrow$ Industrialization
  - Framework for testing ideas and algorithms
  - Technical changes hidden from the user
  - Modern programming techniques: unit tests, pair programming etc.
  - Much better self checks, regression testing and maintainability

News 2013/early 2014: upcoming official release 2.2.x

- Lost of WHIZARD members 2012: some features delayed in 2013

  (pre-)release version 2.2.0_α-2 available!

- WHIZARD core: insert an extra abstraction layer, consistently separate interface from implementation Complete object orientation
  - Replaceable modules with well-defined interface: matrix-elements, beam structure, phase space, integration, decays, shower, ...
  - Much easier to contribute new parts to the code $\Rightarrow$ Industrialization
  - Framework for testing ideas and algorithms
  - Technical changes hidden from the user
  - Modern programming techniques: unit tests, pair programming etc.
  - Much better self checks, regression testing and maintainability

- Revised model for BSM interactions of electroweak vector bosons (w/ light Higgs)

- Process containers: inclusive production samples (e.g. SUSY)

- Automatic generation of decays, depending on the model

News 2013/early 2014: upcoming official release 2.2.x

- Lost of WHIZARD members 2012: some features delayed in 2013

  (pre-)release version 2.2.0_α-2 available!

- WHIZARD core: insert an extra abstraction layer, consistently separate interface from implementation Complete object orientation
  - Replaceable modules with well-defined interface: matrix-elements, beam structure, phase space, integration, decays, shower, ...
  - Much easier to contribute new parts to the code $\Rightarrow$ Industrialization
  - Framework for testing ideas and algorithms
  - Technical changes hidden from the user
  - Modern programming techniques: unit tests, pair programming etc.
  - Much better self checks, regression testing and maintainability

- Revised model for BSM interactions of electroweak vector bosons (w/ light Higgs)
- Process containers: inclusive production samples (e.g. SUSY)
- Automatic generation of decays, depending on the model
- Easy cascades: `"e+", "e-"=> (t => ("W+"=> "mu+", numu), b), tbar`
- Specification of QCD and electroweak order
- Improvements to the SINDARIN steering language

New (LC-related) features / Plans

- Lumi-linker interface — courtesy of T. Barklow
- LCIO support (C++ interface) — courtesy of F. Gaede
- Support for ILC beam spectra within CIRCE1 — courtesy of G. Wilson
- Module for CLIC beam spectra — provided by S. Poss

New (LC-related) features / Plans

- Lumi-linker interface courtesy of T. Barklow
- LCIO support (C++ interface) courtesy of F. Gaede
- Support for ILC beam spectra within `CIRCE1` courtesy of G. Wilson
- Module for CLIC beam spectra provided by S. Poss
- BSM: general Lorentz structures in matrix-element generator (O'Mega)
- Parsing of UFO files

New (LC-related) features / Plans

- Lumi-linker interface courtesy of T. Barklow
- LCIO support (C++ interface) courtesy of F. Gaede
- Support for ILC beam spectra within `CIRCE1` courtesy of G. Wilson
- Module for CLIC beam spectra provided by S. Poss
- BSM: general Lorentz structures in matrix-element generator (O'Mega)
- Parsing of UFO files
- Working on performance gain: multi-leg, parallelization, smaller expressions etc.

New (LC-related) features / Plans

- Lumi-linker interface courtesy of T. Barklow
- LCIO support (C++ interface) courtesy of F. Gaede
- Support for ILC beam spectra within `CIRCE1` courtesy of G. Wilson
- Module for CLIC beam spectra provided by S. Poss
- BSM: general Lorentz structures in matrix-element generator (O'Mega)
- Parsing of UFO files
- Working on performance gain: multi-leg, parallelization, smaller expressions etc.

- CKKW(-L) matching for jets JRR/Weiss, ca. 2014
- Matched $e^+e^- \to$ jets at LO and NLO, POWHEG box formalism JRR/Weiss, ca. 2015

New (LC-related) features / Plans

- Lumi-linker interface courtesy of T. Barklow
- LCIO support (C++ interface) courtesy of F. Gaede
- Support for ILC beam spectra within `CIRCE1` courtesy of G. Wilson
- Module for CLIC beam spectra provided by S. Poss
- BSM: general Lorentz structures in matrix-element generator (O'Mega)
- Parsing of UFO files
- Working on performance gain: multi-leg, parallelization, smaller expressions etc.

- CKKW(-L) matching for jets JRR/Weiss, ca. 2014
- Matched $e^+e^- \to$ jets at LO and NLO, POWHEG box formalism JRR/Weiss, ca. 2015
- Threshold resummation for $e^+e^- \to W^+W^-$, $t\bar{t}$ etc.

Status of refactoring:

Well, what shall I say ...

Status of refactoring:

Well, what shall I say ...



Status of refactoring:

ok, LC features have pretty high priority



Summary and Outlook



- WHIZARD 2 for ILC (and LHC) physics
- Versatile, user-friendly tool for SM & BSM physics
- Highest-possible support for LC beam structures
- Covers the whole SM, and most possible paths beyond
- Steered via the HepForge page:
  `http://projects.hepforge.org/whizard`
- Expect continuous improvement

Summary and Outlook



- WHIZARD 2 for ILC (and LHC) physics
- Versatile, user-friendly tool for SM & BSM physics
- Highest-possible support for LC beam structures
- Covers the whole SM, and most possible paths beyond
- Steered via the HepForge page:
  `http://projects.hepforge.org/whizard`
- Expect continuous improvement
  - WHIZARD 2.2.0_α-2 available now

Summary and Outlook



- WHIZARD 2 for ILC (and LHC) physics
- Versatile, user-friendly tool for SM & BSM physics
- Highest-possible support for LC beam structures
- Covers the whole SM, and most possible paths beyond
- Steered via the HepForge page:
  `http://projects.hepforge.org/whizard`
- Expect continuous improvement
  - WHIZARD 2.2.0_α-2 available now
  - WHIZARD 2.2.0_β for SiD workshop

Summary and Outlook



- WHIZARD 2 for ILC (and LHC) physics
- Versatile, user-friendly tool for SM & BSM physics
- Highest-possible support for LC beam structures
- Covers the whole SM, and most possible paths beyond
- Steered via the HepForge page:
  `http://projects.hepforge.org/whizard`
- Expect continuous improvement
  - WHIZARD 2.2.0_α-2 available now
  - WHIZARD 2.2.0_β for SiD workshop
  - WHIZARD 2.2.0 for LCWS 2013, Tokio

as usual: **we're open to users wish list!**

`whizard@desy.de`

Interesting times ahead

Interesting times ahead



Kent Nagano, Director General, 2015-