

## Toolkit/checklist

$$I = \sqrt{-1}.$$

Analytic function of complex number  $z$  must be of form  $f(z)$ , i.e. excludes  $f(z, z^*)$ .

Kronecker delta:  $\delta_{ij} = 1$  if  $i = j$ , 0 otherwise.

Completely antisymmetric tensor:

$$\epsilon_{ijk} = \begin{cases} 1 & \text{if } ijk = 123 \text{ or cyclic permutations thereof} \\ -1 & \text{if } ijk = 321 \text{ or cyclic permutations thereof} \\ 0 & \text{otherwise.} \end{cases}$$

Summation convention:  $\sum_i A_i B_i = A_i B_i$ , i.e. if same index ( $i$ ) appears twice in term, sum over it, don't bother writing  $\sum$ .

Note e.g.  $A_{ij} \delta_{jk} = A_{ik}$ .

If  $A_{ij} = A_{ji}$  (symmetric in  $i, j$ ),  $\epsilon_{ijk} A_{jk} = 0$ .

Scalar product:  $A_i B_i$ . (For complex  $A_i, B_i$ , scalar product defined as  $A_i^* B_i$ .)

Contraction of index  $k$  in  $A_{ijk}$  with  $B_i$ :  $A_{ijk} B_k = C_{ij}$ , i.e.  $k$  becomes dummy index, rank of  $A$  reduced from 3 to 2 (a matrix).

A vector  $V$  is an  $n$ -tuple “down the page”

$$V = \begin{pmatrix} 2 + 3I \\ 1 - 4I \\ \vdots \end{pmatrix}.$$

Its *transpose* is  $V^T = (2 + 3I, 1 - 4I, \dots)$ .

Its *conjugate transpose* is  $V^\dagger = V^{T*} = (2 - 3I, 1 + 4I, \dots)$

Scalar product:  $A^{(\dagger)} \cdot B$ .

Matrix  $A$  acting on vector  $v$ :  $A_{ij} v_j = (Av)_i$  (and  $v_j A_{ji} = (v^T A)_i$ ).

Two matrices  $A, B$ :  $(AB)^\dagger = B^\dagger A^\dagger$  (likewise for simple transpose  $T$ ).

Unit matrix  $\mathbf{1}$  (not 1):  $\mathbf{1}_{ij} = \delta_{ij}$ .

Hermitian matrix  $H$ :  $H^\dagger = H$ .

Unitary matrix  $U$ :  $U^\dagger U = \mathbf{1}$ .

Orthogonal matrix  $M$ :  $M^T M = \mathbf{1}$ .

Definition of “exponentiation” of a matrix  $A$ :  $e^A = \mathbf{1} + A + \frac{1}{2!}A^2 + \frac{1}{3!}A^3 + \dots$  (this series is always convergent).

Note  $e^{VAV^{-1}} = Ve^AV^{-1}$  for any  $V$  (because  $VA^nV^{-1} = (VAV^{-1})^n$ ).

$e^{IA}$  unitary if  $A$  Hermitian (because  $(e^{IA})^\dagger e^{IA} = e^{-IA^\dagger} e^{IA} = e^{-IA} e^{IA} = \mathbf{1}$ ).

Trace of matrix  $A$ :  $\text{tr}[A] = A_{ii}$ .

Determinant of matrix  $n \times n$   $A$ :  $\det(A) = \epsilon_{ijk\dots} A_{i1} A_{j2} A_{k3} \dots = \frac{1}{n!} \epsilon_{ijk\dots} \epsilon_{abc\dots} A_{ia} A_{jb} A_{kc} \dots$

Note  $\det(AB) = \det(A)\det(B)$

(because  $\det(AB) = \frac{1}{n!} \epsilon_{ijk\dots} \epsilon_{abc\dots} (A_{im} B_{ma})(A_{jn} B_{nb})(A_{kp} B_{pc}) \dots = \frac{1}{n!} n! \frac{1}{n!} \epsilon_{ijk\dots} \epsilon_{def\dots} A_{id} A_{je} A_{kf} \dots \frac{1}{n!} \epsilon_{qrs\dots} \epsilon_{abc\dots} B_{qa} B_{rb} B_{sc} \dots$ ).

If  $Av = 0$ , then either  $v = 0$ , or  $v \neq 0$  and  $\det(A) = 0$

( $v_m \det(A) = \epsilon_{ijk\dots} (v_m A_{im}) A_{j2} A_{k3} \dots = \epsilon_{ijk\dots} \underbrace{\left( \sum_n v_n A_{in} \right)}_{=0} A_{j2} A_{k3} \dots = 0$ ).

Linearly independent vectors  $v_i$ :  $\sum_i \alpha_i v_i = 0 \implies \alpha_i = 0$ .

$N$  linearly independent vectors  $v_i$  form *complete basis* in some  $N$  dimensional vector space. (i.e. any  $V = \sum_i \alpha_i v_i$ ).

Any other basis of  $N$  linearly independent vectors  $v'_i$  in this space is complete, i.e. any  $V = \sum_i \alpha'_i v'_i$ :

Can write  $v'_i = \sum_j C_{ij} v_j$ , so  $\alpha'_i = \sum_j \alpha_j (C^{-1})_{ji}$ ,

and  $C$  must be invertible for  $v'_i$  to be linearly independent ( $\alpha_i = 0 \implies \alpha'_i = 0$ ).

Eigenvalues  $\lambda_k$  and eigenvectors  $w^{\{k\}}$  of matrix  $A$  from *secular equation*:  $Aw^{\{k\}} = \lambda_k w^{\{k\}}$ .

Requirement of  $(A - \lambda_k \mathbf{1})w^{\{k\}} = 0$  and  $w^{\{k\}} \neq 0$ : *Characteristic equation*  $|A - \lambda_k \mathbf{1}| = 0$ .

If  $A$  is  $N \times N$ , this is polynomial in  $\lambda_k$  with  $N$  solutions  $\rightarrow N$  eigenvectors.

Hermitian  $A$ :

$\lambda_k$  real ( $w^{\{k\}\dagger} \cdot Aw^{\{k\}} = \lambda_k w^{\{k\}\dagger} \cdot w^{\{k\}} = (w^{\{k\}\dagger} \cdot Aw^{\{k\}})^\dagger = \lambda_k^* w^{\{k\}\dagger} \cdot w^{\{k\}}$ ).

$w^{\{k\}}$  orthogonal ( $w^{\{i\}\dagger} \cdot Aw^{\{j\}} = \lambda_i w^{\{i\}\dagger} \cdot w^{\{j\}} = \lambda_j w^{\{i\}\dagger} \cdot w^{\{j\}}$ , so  $i \neq j, \lambda_i \neq \lambda_j \implies w^{\{i\}\dagger} \cdot w^{\{j\}} = 0$  (choose this for  $\lambda_i = \lambda_j$ )).

Normalize  $w^{\{k\}}$  to be orthonormal:  $w^{\{i\}\dagger} \cdot w^{\{j\}} = \delta_{ij}$ .  $w^{\{k\}}$  form complete basis ( $\sum_i \alpha_i w^{\{i\}} = 0 \implies 0 = \sum_i \alpha_i w^{\{j\}\dagger} \cdot w^{\{i\}} = \alpha_j$ ).

Diagonalization of any matrix  $A$ :  $(P^{-1}AP) = \Lambda$ , where  $\Lambda_{ij} = \lambda_i \delta_{ij}$  (no sum)

(equivalent to the secular equation above with  $P_{ij} = w_i^{\{j\}}$ ).

Hermitian  $A$ :  $P$  is unitary ( $(P^\dagger P)_{ij} = w_k^{\{i\}*} w_k^{\{j\}} = w^{\{i\}\dagger} \cdot w^{\{j\}} = \delta_{ij}$ ).

Symmetric  $A$ :  $P$  is orthogonal.

For any  $A$ ,  $\det(e^A) = e^{\text{tr}[A]}$

$(\det(e^A) = \det(P^{-1}e^A P) = \det(e^{P^{-1}AP}) = \det \underbrace{e^\Lambda}_{\text{diagonal}} = (e^\Lambda)_{11}(e^\Lambda)_{22} \dots = e^{\text{tr}[\Lambda]} = e^{\text{tr}[P\Lambda P^{-1]} = e^{\text{tr}[A]})$ .

Positive definite matrix  $A$ :  $v^\dagger Av \geq 0$  for any vector  $v$ , with equality only when  $v = 0$ .

If  $A$  Hermitian, eigenvalues  $\lambda_k > 0$ :  $0 \leq \underbrace{v^\dagger P^\dagger}_{=v'^\dagger} \underbrace{P^\dagger AP}_{\text{diagonal, elements } \lambda_k} \underbrace{Pv}_{=v'} = \sum_k \lambda_k |v'_k|^2$ .

Use natural units where  $c = \hbar = 1$ .

Spacetime indices are Greek letters, e.g.  $\mu, \nu = 0, 1, 2, 3$   
(spatial indices Roman, e.g.  $i, j = 1, 2, 3$ ).

4-vector  $A \equiv A^\mu = (\mathbf{A}, A^0)$ , where  $\mathbf{A} = (A^1, A^2, A^3)$  is spatial 3-vector.

Metric tensor of special relativity:

$$g_{\mu\nu} = g^{\mu\nu} = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & -1 \end{pmatrix}_{\mu\nu}$$

Raising and lowering of indices:  $g_{\mu\nu}A^{\mu\rho} = A_\nu{}^\rho$  etc.

Note  $g_{\mu\nu}g^{\nu\rho} = g_\mu{}^\rho = \delta_\mu{}^\rho$ .

Define  $x^0 = +t$ , so  $x_0 = -t$ . Likewise  $p^0 = E$ ,  $p_0 = -E$  etc.

Scalar product of 4-vectors:  $A \cdot B = A^\mu B_\mu = A_\mu B^\mu = g^{\mu\nu} A_\mu B_\nu = g_{\mu\nu} A^\mu B^\nu = A^1 B^1 + A^2 B^2 + A^3 B^3 - A^0 B^0$ .

Shorthand for spacetime differentiation:  $\partial_\mu = \frac{\partial}{\partial x^\mu}$  and  $\partial^\mu = \frac{\partial}{\partial x_\mu}$ .

d'Alembertian:  $\partial^2 = \partial_\mu \partial^\mu$ .

Commutator of operators  $O_i, O_j$  (e.g.  $O_i$  can be matrices):  $[O_i, O_j] = [O_i, O_j]_- = O_i O_j - O_j O_i$ .

Anticommutator:  $\{O_i, O_j\} = [O_i, O_j]_+ = O_i O_j + O_j O_i$ .

Dirac delta function  $\delta(x - y) = \int \frac{dp}{2\pi} e^{Ip(x-y)}$ :

$\int_a^b dx f(x) \delta(x - y) = f(y)$  if  $a < y, b > y$  (otherwise = 0).

(Continuous version of  $f_i \delta_{ij} = f_j$ .)

$\delta^3(\mathbf{x}) = \delta(x_1) \delta(x_2) \delta(x_3)$ .