

The Multi Messenger Approach and The MAGIC/IceCube Target of Opportunity Program

M. Tluczykont, E. Bernardini & K. Satalecka
for the IceCube & MAGIC Collaborations
Helmholtz Nachwuchsgruppe, DESY, Zeuthen



Venice, TeV III, 2007

- **Multi messenger approach**
 - principles
 - ongoing efforts
- **Neutrino Triggered Target of Opportunity program**
- **Gamma-Ray data archive & lightcurve analysis**

Motivation: Multi-Messenger Approach

- **Multi-Wavelength:**

- different Wavelength bands (simultaneous / non-simultaneous)
- combination of multi-wavelength information: maximize e.m. phenomenology
- better understanding of astrophysical objects

- **Multi-Messenger:**

- other messengers + e.m. information: **Neutrinos**, Gravitational waves
- **complete phenomenology** & better understanding
- **increase detection significance** of new messengers (e.g. coincidence)
- can build **confidence in new messengers**

Neutrinos and Gamma-Rays

Acceleration of cosmic rays: Hadronic interactions lead to γ -ray & ν emission

γ -rays can be produced via hadronic or leptonic processes

ν emission is an unambiguous signature for hadronic acceleration

Neutrino Telescopes	Gamma-Ray Telescopes
$>2\pi$ sky coverage	0.02sr sky coverage
90% time coverage	max. 12 % time coverage
angular res. $O(1^\circ)$	$O(0.1^\circ)$
low statistics / sensitivity	high statistics / sensitivity
bad spectral resolution	excellent spectral resolution

Multi-Messenger studies

Several ongoing Multi-Messenger studies including neutrinos

Can include Radio, Optical, X-rays, gamma-rays + neutrinos

- **AMANDA / IceCube**

- MPI-K: E. Resconi et al.
- UW Madison: T. Monaruli et al.
- DESY: E. Bernardini et al.
- Humboldt Uni: M. Kowalski et al.

- **ANTARES / Km3NeT**


- Predictions based on galactic H.E.S.S. sources: Kappes, Hinton, Stegmann & Aharonian, astro-ph/0607286
- ...

Multi-Messenger Approach


Focus of this work: Very High Energy (VHE) γ -Ray Data

Scope: multi-messenger studies with time variable signals \rightarrow **light curves**

Need: archivation & combination of VHE γ -Ray data from main experiments




Multi-Messenger Approach: Offline analysis



principle: search for $\nu\gamma$ correlations from variable objects (e.g. 1ES 1959+650)
need **simultaneous data**: sparse due to low γ -ray coverage

Neutrino triggered Target of Opportunity program: NToO



ν from predefined list of objects:
trigger γ -ray observations within predefined time window

pure background neutrino sample:
random coincidences with flares

signal neutrino content:
enhanced coincidences

Goals

ensure simultaneous ν/γ data
improve detection chance for ν telescopes
improve phenomenological knowledge of sources

see, e.g. Bernardini et al., Palaiseau 2005, Ackermann et al., ICRC 2007

NToO test run: details

Implementation of NToO idea: cooperation between AMANDA & MAGIC


- 2 X-ray binaries: LSI+61 303, GRS 1915+105
- 3 Blazars: 1ES 2344+514, 1ES 1959+650, Mrk 421
- **predefinition of γ -observation window**: observation of object for 1 hour within a window of 1 day after neutrino trigger was issued
- **predefinition of coincidence**: γ -ray high-state within obs. window
- **predefinition of γ -ray high-state**: observed flux above predefined threshold
- Results: Ackermann et al., ICRC 2007, talk by E. Bernardini

NToO: possible interpretation scheme

$$P = \sum_{i=n_{\text{obs}}}^{+\infty} \frac{(n_{\text{bck}})^i}{i!} e^{-n_{\text{bck}}} \sum_{j=n_{\gamma}}^i \frac{i!}{j!(i-j)!} p_{\gamma}^j (1-p_{\gamma})^{i-j},$$

- n_{obs} observed neutrino events
- n_{bck} expected atmospheric neutrino events
- n_{γ} observed ν / γ coincidences
- p_{γ} Probability of a γ -ray high-state

Ranges of p_{γ} for which an observation of n_{γ} coincidences would yield a Significant detection, $P > 5\sigma$




NToO: possible interpretation scheme

Example neutrino observation: $n_{\text{obs}} = 10$, $n_{\text{bck}} = 10$

In itself this neutrino observation is not really significant...

Adding information from coincident observations:



Need information on γ -ray variability: p_γ

Very little known so far

VHE lightcurve archive:

- central open archive
- definition of common lightcurve format
- lightcurve combination

Gamma-Ray Data Archive

- Collect all available (public) data from VHE experiments
- Current focus: Active Galactic Nuclei
- Extension to other objects...
- Issues:
 - no common lightcurve format → **definition of SLF lightcurve format**
 - normalization difficult: different flux units, partly unknown calibration
 - old data are not available in electronic form
 - ...
- open archive but... use with care!

<http://www-zeuthen.desy.de/multi-messenger/GammaRayData/>

Gamma-Ray Data Archive – Screenshot

<http://www-zeuthen.desy.de/multi-messenger/GammaRayData/>

Objects

[1ES1959+650](#) [Mrk421](#) [Mrk501](#) [PKS2155-304](#)

1ES 1959+650:

publications	available data							
	1999	2000	2001	2002	2003	2004	2005	2006
ITA_99 HEGRA_00/01 Whipple_02 VERITAS_02 VERITAS_06 MAGIC_05 CAT		HEGRA_CTS * HEGRA_CT1	HEGRA_CTS * HEGRA_CT1	HEGRA_CTS * HEGRA_CT1 Whipple	Whipple ?	MAGIC Whipple ?	VERITAS MAGIC ?	VERITAS MAGIC ?
slf format:		HEGRA_CTS HEGRA_CT1	HEGRA_CTS HEGRA_CT1	HEGRA_CTS HEGRA_CT1 Whipple		MAGIC	VERITAS	VERITAS

[1ES1959_combined_v1.slf](#)

Mrk421

publications	available data															
	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	
publ_link	Whipple	Whipple	Whipple	Whipple	HEGRA_CT1 CAT	HEGRA_CTS HEGRA_CT1 CAT	HEGRA_CTS HEGRA_CT1 Whipple CAT	HEGRA_CTS HEGRA_CT1 CAT	HEGRA_CTS HEGRA_CT1 CAT	HEGRA_CTS HEGRA_CT1 MILAGRO	HEGRA_CTS HEGRA_CT1 MILAGRO STACEE	Whipple MILAGRO MILAGRO	Whipple MILAGRO MILAGRO	Whipple MAGIC H.E.S.S. MILAGRO	VERITAS MAGIC	VERITAS
slf format:	Whipple	Whipple	Whipple	Whipple	HEGRA_CT1 CAT	HEGRA_CTS HEGRA_CT1 CAT	HEGRA_CTS HEGRA_CT1 Whipple CAT	HEGRA_CTS HEGRA_CT1 CAT	HEGRA_CTS HEGRA_CT1 CAT	HEGRA_CTS HEGRA_CT1 CAT	HEGRA_CTS HEGRA_CT1	Whipple Whipple	Whipple	HESS MAGIC	MAGIC VERITAS	VERITAS

Mrk421, combined lightcurve: [Mrk421_combined_v2.slf](#)

Current status of conversion/plotting :

The SLF format: Smarties Lightcurve Format

1. MJD of start of observations
2. MJD of end of observations
3. Measured flux
4. Statistical error of measured flux
5. Systematic error of measured flux
6. Spectral index
7. Spectral index statistical error
8. Spectral index systematic error
9. Energy threshold
10. Energy Cutoff
11. Experiment
12. Duration of observation
13. Additional entry 1
14. ...

simple ascii-table


extendable (e.g. spectra with exponential cutoff)

Light-Curve Combination

- **Ingredients:**


- Common lightcurve format (SLF), threshold (1 TeV), Flux unit (Crab)
- Spectral shapes as measured by experiments: power-law including possible cutoff

- $$F(E > 1 \text{ TeV}) = \left(\frac{E_{\text{thr}}}{1 \text{ TeV}} \right)^{-\Gamma+1} \frac{\Phi(>E_{\text{thr}})}{\Phi_{\text{Crab}}(E>1\text{TeV})}$$




Long-Term Lightcurve of Mrk 421

15 years · more than 1500 h of observations



Mrk 501 Long-Term Lightcurve



Lightcurve Analysis:

- flux-state distributions
- relative high-state rates
- comparison to ASM X-ray data



Mrk 421 Flux-State Distributions: Long-Term VHE Data

Flux distribution

- selected observations (no U.L.)
- bias at high flux states: high-state trigger

Zoom into low fluxes

- observation of low state ?
- ... but at low flux states: sensitivity threshold bias




Stochastic flux state distribution: $\frac{dN}{d\Phi} = a \cdot e^{b\Phi}$

Estimation of Relative High-State Rate

Motivation: important ingredient in multi-messenger analyses (\mathcal{V} Correlation)

Hypothesis: stochastic flux-state distribution $\frac{dN}{d\Phi} = a \cdot e^{b\Phi}$

High-State Rate: $R_{HS}(F_{thr}) = \frac{T(F > F_{thr})}{T_{tot}} = \frac{\int_{F_{thr}}^{\infty} e^{b\Phi} dx}{\int_{F_0}^{\infty} e^{b\Phi} dx} = \frac{e^{bF_{thr}}}{e^{bF_0}}$





Back to NToO: highstate rates as input

	LSI+61 303	GRS 1915+105	1ES 2344+514	1ES 1959+650	Mrk 421
n_{bck}	0.86	1.26	0.99	0.92	1.51
n_{obs}	0	1	1	0	3
Follow ups	0	0	1	0	1
n_{γ}	–	–	0	–	0
F_{thr} [C.U.]	0.2	0.2	0.5	1.0	4.0
p_{γ}	–	–	–	< 0.15	< 0.05
P_{ν}	1.0	0.7	0.6	1.0	0.2

VHE to X-Ray Correlation

- observations are **not 100% simultaneous**
- high VHE flux states: observational bias due to external- and self-triggering
- on average: linear correlation up to 4 Crab-level





Comparison to ASM X-Ray Data

Flux-State Distributions from Contemporaneous data ASM X-ray data

- http://xte.mit.edu/ASM_lc.html
- only partly simultaneous !


Component Interpretation

- Noise: Gaussian
- Signal: Gaussian + Exponential



Stochastic flux state distribution: $\frac{dN}{d\Phi} = a \cdot e^{b\Phi}$

Comparison to ASM Data: High-State Rate




expected and found

comparable R_{HS} :


15-year-VHE lightcurve + ASM archive data:

- ASM data: same stochastic behaviour
- only partly simultaneous data
- clear correlation seen




Lightcurve simulation: proof of principle

simulation of exponential flux-state distribution (arb. units)



Lightcurve simulation: proof of principle

Picking randomly observations with a total coverage of 1 %



Plans

- variability simulation
- proper treatment of upper limits

Summary & Outlook

- **Very High Energy Data Archive:**

<http://www-zeuthen.desy.de/multi-messenger/GammaRayData/>

- **archivation of lightcurves: AGN ++**
- **combination of data: long-term lightcurves**
- **analysis: estimation of high state rates**
- **collecting data: archive growing ...**