# The $N^{O}E$ detector performances for large mixing angle long baseline neutrino oscillations


G.C.Barbarino[1], D.Campana[1], F.Guarino[1], A. Lauro[1], G.Osteria[1], U.Rubizzo[1], A.Margiotta[2], M.Spurio[2], P.Bernardini[3], G.Mancarella[3], D.Martello[3], A.Surdo[3], S.Bussino[4], E.Lamanna[4], M.De Vincenzi[4,5], A.Di Credico[6], A.Grillo[6], C.Gustavino[6], S.Mikheyev[6,7], E.Scapparone[6] G.De Cataldo[8], P.Fusco[8], N.Giglietto[8], M.N.Mazziotta[8], A.Rainó[8], P.Spinelli[8]

[1]*Dip. di Scienze Fisiche dell' Universitá di Napoli and INFN Sez. di Napoli - Italy*

[2]*Dip. Fisica dell' Universitá di Bologna and INFN Sez. di Bologna - Italy*

[3]*Dip. Fisica dell' Universitá di Lecce and INFN Sez. di Lecce - Italy*

[4]*Dip. Fisica dell' Universitá di Roma and INFN Sez. di Roma - Italy*

[5]*Dip. Fisica dell' Universitá di Roma III - Italy*

[6]*INFN, Laboratory Nazionali del Gran Sasso, Assergi - Italy*

[7]*Institute of Nuclear Research, Russian Academy of Science, Moscow, Russia*

[8] *Dip. di Fisica dell'Universitá di Bari and INFN Sez. di Bari - Italy*



(July 2, 1997)

# 1 Introduction

One of the open and most debated problems in elementary particle physics is whether neutrinos are massless or not. Very small neutrino masses are unlikely to be measured via the classical $\beta$-decay procedure. If neutrinos have non zero mass and the mass eigenstates are mixed, the most promising way to measure their mass is through oscillation experiments, as foreseen by Bruno Pontecorvo. In the simplified hypothesis of a mixing between two neutrino families, the probability of observing $\nu_i \rightarrow \nu_j$ oscillations at a distance $L$ is given by

$$P(\nu_i \rightarrow \nu_j) = \sin^2 2\theta \sin^2 \left[\frac{1.27\Delta m^2 L}{E_\nu}\right] \quad (1)$$

where $\Delta \mathrm{m}^2 = \mathrm{m_i^2} - \mathrm{m_j^2}$ is in $\mathrm{eV}^2$, L is the distance from the $\nu$ source to the detector in km and $\mathrm{E}_\nu$ is the neutrino energy in GeV.

For maximal mixing, this probability is different from zero if $\Delta \mathrm{m}^2$ is at least $\sim \mathrm{E}_\nu/\mathrm{L}$. Solar neutrino data ($\mathrm{E}_\nu \sim$ few MeV, $L = 150 \cdot 10^6$ km) cover $\Delta \mathrm{m}^2$ regions from $10^{-5}$ to $10^{-10}$ $\mathrm{eV}^2$ and existing accelerator experiments can go up to $\Delta \mathrm{m}^2 \sim 1$ $\mathrm{eV}^2$. The region of $\Delta \mathrm{m}^2 = 10^{-2} \div 10^{-3}$ $\mathrm{eV}^2$ can be covered with long baseline $\nu_\mu$-beams (L $\sim 10^3$ km). The $\Delta \mathrm{m}^2$ region which could be investigated by experiments for oscillation studies using long baseline $\nu$-beams from accelerators coincides with the region of atmospheric neutrinos ($\mathrm{E}_\nu \sim$ GeV, L $= 10 \div 10^4$ km). Long baseline neutrino beams provide a unique way to study $\nu_\mu$ oscillations at small $\Delta \mathrm{m}^2$ values, offering several advantages when compared with atmospheric neutrinos (high flux, known energy spectrum, known initial neutrino direction and flavour and $\nu_e$ contamination). A muon neutrino beam peaked around an energy of several GeV can be used to detect $\tau$ neutrinos appearance via $\nu_\tau$ CC interactions. Long baseline projects are being discussed in various laboratories in the world (Fermilab to Soudan, CERN to Gran Sasso, KEK to SuperKamiokande) and are at different stages of approval.

# 2 Atmospheric neutrinos

## 2.1 Introduction

Atmospheric neutrinos originate mainly from the decays of charged $\pi$, K and muons, produced in cosmic-ray-induced air showers. From these decay chains one expects the ratio of the number of $\nu_\mu(\bar{\nu_\mu})$ to $\nu_e(\bar{\nu_e})$ to be approximatively equal to two at low energy (1-2 GeV). Atmospheric neutrinos can be identified in different energy ranges:

1. At low energy by means of their interaction within a massive detector. These interactions may be studied with water Čerenkov detectors (IMB [1], Kamiokande [3]), or dense tracking calorimeters (NUSEX [2], Soudan 2 [5], Frejus [4]). Neutrino interactions are then classified as fully or partially contained, depending on whether all interaction products are contained within the detector fiducial volume or high energy muons leave the detector.

2. At higher energy by means of the muons produced by $\nu_\mu$ CC interactions in the rock surrounding the detector. In this case the produced muons may either cross or stop in the detector and can be studied with large area tracking detectors. The MACRO experiment measures a deficit of upward-throughgoing $\mu$'s generated by $\nu$ interactions. This result may be another clue of $\nu$ oscillation, even though it is compatible with the expectation with a 8 % confidence level [6], [7].

## 2.2 Contained events

The water Čerenkov experiments Kamiokande and IMB are able to separate $\nu_\mu$ CC from $\nu_e$ CC interactions and find that the ratio

$$R = \frac{\left(\frac{\nu_\mu}{\nu_e}\right)_{\rm data}}{\left(\frac{\nu_\mu}{\nu_e}\right)_{\rm MC}}$$

is smaller than expected. Experimental results on contained atmospheric neutrino events are consistent with a 30-40% $\nu_\mu$ deficit. Such a deficit could be the result of either $\nu_\mu \to \nu_\tau$ or $\nu_\mu \to \nu_e$ oscillations or both. At this stage the simulation of $\nu_\mu$ and $\nu_e$ fluxes favours a $\nu_\mu \to \nu_\tau$ interpretation of the atmospheric neutrino puzzle. In particular, the allowed regions in the oscillation parameter space from the Kamiokande data are: $\Delta \mathrm{m}^2 > 4 \cdot 10^{-3}$ eV$^2$, $\sin^2(2\theta) > 0.3$ for $\nu_\mu \to \nu_e$ and $\Delta \mathrm{m}^2 > 10^{-3}$ eV$^2$, $\sin^2(2\theta) > 0.4$ for $\nu_\mu \to \nu_\tau$. A similar analysis performed by calorimetric experiments (NUSEX and Frejus) are not in agreement with this result; on the contrary Soudan 2 seems to confirm Čerenkov detector results.

### 2.3 Partially contained events

A different analysis has been performed by the Kamiokande group in the so-called multi-GeV energy range. These events are:

1. Fully-contained events with visible energy larger than 1.33 GeV;

2. Partially-contained events which have their interaction vertex in the fiducial volume and a visible track leaving the detector.

The estimated mean energy of neutrinos is 5 GeV for fully contained e-like events, 2.5 GeV for fully contained $\mu$-like events, and 9 GeV for partially-contained events. Also in this data sample an analogous deficit is observed. A feature of this data is the relative excess of electrons and the relatively small deficit of muon neutrinos, compared to what is expected from the $\nu_\mu \rightarrow \nu_\tau$ hypothesis. Assuming oscillation, the best fit gives the following parameters: $\Delta m^2 = 1.8 \cdot 10^{-2}$ eV$^2$ and $\sin^2(2\theta)$=1.0 for $\nu_\mu \rightarrow \nu_e$ and $\Delta m^2 = 1.6 \cdot 10^{-2}$ eV$^2$ and $\sin^2(2\theta)$=1.0 for $\nu_\mu \rightarrow \nu_\tau$. Thus, while the low energy sample suggests $\nu_\mu \rightarrow \nu_\tau$, the high energy sample may suggest $\nu_\mu \rightarrow \nu_e$ oscillations. Finally, combining all their data (sub-GeV and multi-GeV), the Kamiokande group finds a rather limited allowed region, both for $\nu_\mu \rightarrow \nu_\tau$ and for $\nu_\mu \rightarrow \nu_e$ with nearly full mixing and with $\Delta m^2 \sim 10^{-2}$ eV$^2$.

## 3 Three-Flavour Mixing

A more general assumption making use of a three generation neutrino oscillation formalism has been recently carried out by many authors. In particular Harrison, Perkins and Scott[8] have pointed out that atmospheric and solar neutrino data could be explained in a threefold maximal mixing scheme assuming two independent mass-squared differences: $\Delta m^2 = (0.72 \pm 0.18) \cdot 10^{-2}$ eV$^2$ and $\Delta m'^2 < 0.9 \cdot 10^{-11}$ eV$^2$. For L/E $< 1/\Delta m^2$ the flavour survival probability formula $P(\nu_l \rightarrow \nu_l)$ returns a value equal to one, while for L/E $> 1/\Delta m^2$ the value of this quantity, averaged over the neutrino spectrum, is expected to be 5/9. The two plateau regions are separated by a well localized threshold at L/E $\sim 1/\Delta m^2$. The corresponding appearance probability $P(\nu_l \rightarrow \nu_{l'})$ to produce a neutrino of different flavour $l' \neq l$ is equal to 2/9 sharing the oscillation between $\nu_\mu \rightarrow \nu_e$ and $\nu_\mu \rightarrow \nu_\tau$ channel. A similar threshold at L/E $\sim 1/\Delta m'^2$ gives rise to the third plateau characterized by a survival probability equal to 1/3. The data from all the experiments are shown in Fig. 1 as a function of L/E: they suggest the existence of a sharp