DELPHI 97-83 CONF 69
28 July, 1997

A study of the reaction $e^+e^- \to \mu^+\mu^-\gamma_{ISR}$ at LEP and search for new physics at annihilation energies near 80 GeV

Preliminary

DELPHI Collaboration

C. De Clercq
IIHE (VUB-ULB), Brussels

Abstract

A study of the channel $e^+e^- \to \mu^+\mu^-\gamma_{ISR}$, where γ_{ISR} is an initial state radiation photon, is presented using data collected by the DELPHI experiment from 1991 to 1995. The total cross-section at effective annihilation energies ($\sqrt{s'}$) below the Z^0 peak is obtained by using the events with relatively hard initial state radiative photon(s) ($E_\gamma > 1$ GeV). The differential cross-section as a function of the muon polar production angle is also determined in order to extract the forward-backward asymmetries for the reaction $e^+e^- \to \mu^+\mu^-$ at energies $\sqrt{s'}$ between 20 and 87 GeV. The ratio of the helicity cross-sections $\frac{\sigma_{LL}+\sigma_{RR}}{\sigma_{RL}+\sigma_{LR}}$, where the two subscripts stand for the helicities of the incoming e^- and outgoing μ^- respectively, is extracted from the differential cross-sections in order to test the Standard Model and to look for new physics near 80 GeV. No deviations from the Standard Model were found.

Paper submitted to the LP'97 Conference
Hamburg, July 28 - August 1st

1 Introduction

In this paper, experimental results from studies of events collected at LEP1 in the channel $e^+e^- \to \mu^+\mu^-\gamma_{ISR}$, with γ_{ISR} being an initial state radiation photon, are used to probe the cross-sections and forward-backward asymmetries in the energy region between LEP1 and TRISTAN and down to PETRA energies. Similar measurements have been performed previously [1]. In this paper an analysis in terms of helicity cross-sections [2] is also performed, and is shown to be particularly useful in a search for new physics near the effective annihilation energy $\sqrt{s'} = 80$ GeV.

The investigation of the $e^+e^- \to \mu^+\mu^-$ cross-section at energies below the Z^0 peak is attractive because no experiments have ever taken data at energies between 61 and 88 GeV. In addition, an indication of a deviation of the cross-section from the predictions of the Standard Model (SM) at 58 GeV has been reported from TRISTAN [3]. Measurable deviations in the $e^+e^- \to f\bar{f}$ cross-section in this energy range can arise in several models beyond the SM, for instance those which introduce an additional Z' gauge boson.

In a previous paper [4] an analysis was presented of radiative muon events using the data collected by DELPHI up to 1994. In this paper a new analysis is presented based on the data collected between 1991 and 1995.

The paper is organised as follows. Section 2 summarises the theoretical background to the analysis. Section 3 describes briefly the DELPHI detector and presents the data samples used. Section 4 discusses the event selection procedures. Section 5 presents the calculation of the total Improved Born cross-sections. The analysis of the differential cross-sections in terms of the forward-backward asymmetry and the helicity cross-sections is presented in Section 6. Finally, Section 7 summarises.

2 Theoretical background

2.1 Total cross-section

Electromagnetic radiative corrections to the interaction $e^+e^- \to f\bar{f}$ substantially distort the Born-level cross-section at energies around the Z^0 pole [5].

Most of the effects can be understood to arise from initial state radiation (ISR), after which the effective annihilation energy $\sqrt{s'}$ of the e^+e^- system is less than the total centre-of-mass energy $\sqrt{s}$.

The cross-section for the reaction $e^+e^- \to \mu^+\mu^-$ has contributions from direct Z^0 and photon terms and from $\gamma - Z^0$ interference. Radiative corrections can be divided into the following components:

- emission of real photons from the initial and/or final state fermions,
- corrections to the Z^0 and γ propagators, consisting of loop diagrams involving any particles which couple to these bosons,
- vertex corrections, involving virtual photons as well as any other particles which couple to the initial or final fermions,
- box diagrams, involving the exchange of two bosons (γ, Z^0).

At the present level of precision, the box diagrams can be neglected. To a very good approximation, the effect of the vertex and propagator corrections can be absorbed into a redefinition of the Born-level parameters that retains the structure of the Born-level formulae (Improved or Effective Born Approximation; see e.g. [5]).

Pure QED initial state corrections can be described by a radiator function $H_i(s,s')$, giving the probability density of emitting a photon with a given fraction of the beam energy from the initial state, such that the observed cross-section for $e^+e^- \to f\bar{f}(n\gamma)$ can be written:

$$\sigma_{obs}^{f\bar{f}}(s) \;=\; \int_{s_0'}^{s} H_i(s,s')\;\sigma_{IB}^{f\bar{f}}(s')\;ds' \tag{1}$$

where $s_0' = 4m_f^2$, the invariant mass $M_{f\bar{f}}$ of the fermion pair produced is given by the effective annihilation energy $\sqrt{s'}$, and $\sigma_{IB}^{f\bar{f}}(s')$ is the Improved Born cross-section. Thus the observed cross-section involves a convolution of $\sigma_{IB}^{f\bar{f}}(s')$ with the radiator function $H_i(s,s')$. A similar formula can be written for the case where selection criteria are applied to the final state fermions or to the photons produced.

In addition, there is a small contribution to the cross-section from the interference between initial and final state radiation [6]. This interference is very small in the energy region covered by this analysis, $\sqrt{s'} < 87$ GeV and $E_\gamma > 1$ GeV, and can be neglected at the present level of precision.

For the case where a pure ISR sample is selected, the number of events N_{12} in a given interval (s_1', s_2') can be written as

$$N_{12} \;=\; \mathcal{L}\int_{s_1'}^{s_2'} H_i(s,s')\;\sigma_{IB}^{f\bar{f}}(s')\;ds' \tag{2}$$

where $\mathcal{L}$ is the integrated luminosity. Thus, from the known QED radiator function $H_i(s,s')$ (see [7] and references therein) and the measured integrated luminosity $\mathcal{L}$, the Improved Born cross-section $\sigma_{IB}^{f\bar{f}}(s')$ for energies $\sqrt{s'}$ below the nominal LEP energy $\sqrt{s}$ can be determined by measuring the $\sqrt{s'}$ spectrum from the invariant mass distribution of the final state fermion pairs.

2.2 New physics near the zeroes of the $e^+e^- \to \mu^+\mu^-$ cross-section

At tree level, the $e^+e^- \to \mu^+\mu^-$ cross-section can be written, apart from radiative corrections and possible new physics, as

$$\sigma = \sigma_{LL} + \sigma_{RR} + \sigma_{RL} + \sigma_{LR} \tag{3}$$

where $i(j)$ in σ_{ij} stands for the helicity of the incoming e^- (outgoing μ^-) [2]. Within the Standard Model, σ_{LL} and σ_{RR} have nearby zeroes such that $\sigma_{LL} + \sigma_{RR}$ almost vanishes near $\sqrt{s} = 80$ GeV. This is not the case for $\sigma_{RL} + \sigma_{LR}$, which vanishes at $\sqrt{s} > m_Z$. A study of the cross-section sum $\sigma_{LL} + \sigma_{RR}$ around 80 GeV allows a search for new physics in an environment of minimal background.

As shown in [2], due to their angular dependence of the form $(1 \pm \cos\theta)^2$, where θ is the production polar angle of the μ^- relative to the e^- beam direction, the two helicity

components of the cross-section, $\sigma_{LL} + \sigma_{RR}$ and $\sigma_{RL} + \sigma_{LR}$, can be extracted from the data by a polynomial projection of the differential cross-section as a function of θ:

$$\tilde{\sigma}^{\mu}_{+} = \int_{-C_M}^{C_M} \frac{d\sigma}{d\cos\theta} \cdot F_{+} \cdot d\cos\theta \tag{4}$$

$$\tilde{\sigma}^{\mu}_{-} = \int_{-C_M}^{C_M} \frac{d\sigma}{d\cos\theta} \cdot F_{-} \cdot d\cos\theta \ , \tag{5}$$

where the filters $F_{\pm}$ are polynomials of the form $A(1 \pm B\cos\theta)$, and $C_M = \cos\theta_{max}$ represents the acceptance limits in the analysis.

At tree level in the framework of the Standard Model,

$$\tilde{\sigma}^{\mu}_{+} = \sigma_{LL} + \sigma_{RR} \tag{6}$$

$$\tilde{\sigma}^{\mu}_{-} = \sigma_{LR} + \sigma_{RL} \tag{7}$$

and the ratio $\frac{\tilde{\sigma}^{\mu}_{+}}{\tilde{\sigma}^{\mu}_{-}}$ has a minimum just below 80 GeV. Therefore, this ratio is very sensitive to new physics near this energy.

3 The DELPHI detector and data samples

The DELPHI detector and its performance are described in detail in [8]. In this analysis, the tracking was performed by the Vertex Detector (VD), the Inner Detector (ID), the Outer Detector (OD), the Time Projection Chamber (TPC), and the Forward Chambers (FCA,FCB). In the rejection of events with final state radiation (FSR), the barrel and forward electromagnetic calorimeters were used (HPC and FEMC). The identification of muons was based on the muon chambers (MUB and MUF), the hadron calorimeter (HCAL), and the electromagnetic calorimeters. In what follows, the barrel region was defined as $43^\circ \leq \theta \leq 137^\circ$, and the endcaps as $\theta < 43^\circ$ and $\theta > 137^\circ$, where θ is the polar angle with respect to the electron beam axis.

The data used for the calculation of the forward-backward asymmetries and the helicity cross-section ratios were collected during the years 1991 to 1995, and correspond to an integrated luminosity of around 144 pb^{-1}. For the calculation of the total cross-sections, only the data taken during the years 1992-1995 were used, corresponding to an integrated luminosity of around 129 pb^{-1}. About 70% of the events were collected at the Z^0 peak energy, while the others were produced during a scan at different energies below and above the Z^0 peak.

For simulation studies of the signal, dimuon events were generated using the DYMU3 program [9]. The events were generated with up to two initial state photons and at most one final state photon. For the cross-section studies, where a normalisation to the simulated sample is needed, a total of about 833,000 events were generated, spread over the beam energies at which data were taken in the period 1992 to 1995. For the evaluation of the efficiency and purity of the ISR sample (see Section 4.3), about 240,000 radiative muon events with $\mu^+\mu^-$ invariant mass $M_{\mu\mu} < 88$ GeV/c^2 were generated at the same beam energies.

For the study of the background from the channel $e^+e^- \to \tau^+\tau^-$, about 373,000 events were generated with the KORALZ program [10]. For the study of the background coming from $\gamma\gamma$ processes, about 261,000 $e^+e^- \to e^+e^-\mu^+\mu^-$ and 10,000 $e^+e^- \to e^+e^-\tau^+\tau^-$ events were generated using the FERMISV program [11].

All the generated events were passed through the full DELPHI simulation [8] and the same data reconstruction program as the real data.

4 Selection of dimuon events with ISR

The selection of dimuon events with ISR was performed in two steps. First a sample of dimuon events with or without photon production was selected following the procedure detailed in Section 4.1. From this sample, events with ISR were extracted using the procedure described in Section 4.2.

For the calculation of the cross-sections, the same selection procedure was applied to the 833,000 simulated dimuon events (see Section 3).

4.1 Selection of $e^+e^- \to \mu^+\mu^-(n\gamma)$ events

To select a sample of dimuon events allowing for possible photon emission, the procedure explained in [4] was followed.

The events had to contain two charged particles of momentum greater than 10 GeV/c, both of which were identified as muons either by the muon chambers (MUB and MUF), or by the hadron calorimeter (HCAL), or by the electromagnetic calorimeters (HPC and FEMC). Both particles had to come from the interaction region, which was defined as $|z|$ less than 4.5 cm and R less than 1.5 cm, where z is along the e^- beam direction and R is the radius normal to it. The variable $P_{rad} = \sqrt{p_1^2 + p_2^2}/E_{beam}$, where E_{beam} is the beam energy and p_1 and p_2 are the momenta of the two muons, had to exceed 0.3. Events with more than 5 charged particle tracks were rejected.

To reduce the $\tau^+\tau^-$ background, the following three criteria were introduced. Firstly, if the acollinearity angle between the two muons was larger than 1°, the event was rejected if the energy deposited in the HCAL was larger than a cutoff value dependent on the polar angle (see [12]). Secondly, if the event had more than 2 charged particle tracks, either the acollinearity angle between the two muons had to be less than 1° or both muons had to have at least one associated hit in the muon chambers. Thirdly, in the procedure to separate ISR from Final State Radiation (FSR) events, a variable ΔE_γ was introduced, which was defined as

$$\Delta E_\gamma = E'_\gamma - E''_\gamma \tag{8}$$

where

$$E'_\gamma = \sqrt{s} - E_{\mu^+} - E_{\mu^-} \tag{9}$$

and

$$E''_\gamma = \frac{|\sin(\theta_{\mu^+} + \theta_{\mu^-})|}{|\sin(\theta_{\mu^+} + \theta_{\mu^-})| + \sin\theta_{\mu^+} + \sin\theta_{\mu^-}}\sqrt{s}\ . \tag{10}$$

In these formulae, θ_{μ^+} and θ_{μ^-} are the polar angles, and E_{μ^+} and E_{μ^-} the energies of the muons. The variable E'_γ represents the energy taken by the ISR photons if no FSR photons are produced. The variable E''_γ is an approximation to the ISR photon energy if the muon masses are neglected and if one assumes that a single ISR photon is emitted along the beam direction and that no FSR photons are produced. The variable ΔE_γ was also effective in rejecting tau events, as can be seen in ref. [4]. Events with $\Delta E_\gamma > 25$ GeV were rejected as tau candidates.

Because the selection efficiencies could not be estimated reliably at low polar angles, the cross-sections were determined with samples of events with the μ^- polar angle in the region $20^\circ \leq \theta_{\mu^-} \leq 160^\circ$. For the measurement of the forward-backward asymmetries however, the likelihood fit method is not affected by the selection efficiencies if these are forward-backward symmetric. Therefore, for these measurements the μ^- polar angle region was extended to $11^\circ \leq \theta_{\mu^-} \leq 169^\circ$. For the extraction of the helicity cross-sections, efficiency corrections had to be made, and the polar angle acceptance was dependent on the effective energy $\sqrt{s'}$ (see Section 6.3).

After this selection of dimuon events, data runs were rejected if the parts of the DELPHI detector used in this analysis were not fully operational. The total number of dimuons available for the asymmetry analysis (data taking period 1991-1995) amounted to 157,980. These were spread over 7 energy points, but about 88% were produced at the Z^0 peak. The total number of dimuons selected for the cross-section analysis (data taking period 1992-1995) was 130,008 spread over 3 energy points, about 86% of them being produced at the Z^0 peak.

The tau background was estimated with the simulated $\tau^+\tau^-$ events to be 0.25%. The background from two-photon events was also estimated with simulated events, and found to be less than 0.1% for the channel $e^+e^- \to e^+e^-\mu^+\mu^-$. For the channel $e^+e^- \to e^+e^-\tau^+\tau^-$, no simulated events were found to satisfy the dimuon selection criteria.

The cosmic ray background was estimated from the data, by relaxing the definition of the interaction region [12] and counting the number of additional events accepted in the data sample. It was found to amount to 0.55%.

From the 833,000 simulated $\mu^+\mu^-$ events, 691,335 remained after the dimuon selection in the polar angle region $20^\circ \leq \theta_{\mu^-} \leq 160^\circ$. These events were used for the normalisation in the cross-section calculation (see Section 5).

4.2 Selection of $e^+e^- \to \mu^+\mu^-\gamma_{ISR}$ events

To extract the events with ISR from the dimuon sample described in Section 4.1, the following variables were used.

- $p_T^{miss} = |\sum_{i=1}^{N_{ch}} \vec{p_T^i}|$, the missing transverse momentum of the event with respect to the beam axis.
- α_{acol}, the acollinearity angle between the two muons.
- α_{acop}, the acoplanarity angle defined as the angle between the two planes respectively formed by each of the muons and the electron beam axis.
- E_{em}^{neu}, the total neutral electromagnetic energy of the event, defined as

$$E_{em}^{neu} = E_{em}^{tot} - E_{em}^{\mu^+} - E_{em}^{\mu^-} \tag{11}$$

  where E_{em}^{tot} is the total energy deposited in the electromagnetic calorimeters and $E_{em}^{\mu^+}$ ($E_{em}^{\mu^-}$) is the energy deposited by the μ^+ (μ^-).
- ΔE_γ, defined in equations (8) to (10).

- $\alpha_{\mu\gamma}$, the isolation angle. For a given reconstructed photon, this was taken to be the smallest of the following angles: the two angles between the muon tracks and the flightline of the photon, and the two angles between the direction opposite to the muons and the flightline of the photon. The photons were assumed to be produced at the vertex. They were reconstructed following the method described in [8]. Only photons with a reconstructed energy larger than 1 GeV were considered.

To ensure a high purity of the selected sample for all effective annihilation energies, the selection criteria were taken to be different in each $\sqrt{s'}$ interval. The selection criteria are summarised in Table 1.

Two sets of selections were used, called tight or loose, depending on whether or not a photon was detected in the electromagnetic calorimeters close to one of the muons. The tight selections were applied in the barrel region when a photon was found with isolation angle $\alpha_{\mu\gamma} < 36°$, and in the endcaps when either a) a photon was found with isolation angle $\alpha_{\mu\gamma} < 20°$, or b) $\alpha_{\mu\gamma} < \theta_\gamma$, where θ_γ is the polar angle of the photon with respect to the electron beam axis. In all other cases, the loose selections were applied.

For the effective annihilation energy $\sqrt{s'}$, or equivalently the $\mu^+\mu^-$ invariant mass $M_{\mu\mu}$, the following expression was used:

$$\sqrt{s'} = M_{\mu\mu} = \sqrt{s - 2E''_\gamma\sqrt{s}}\ . \tag{12}$$

The justification for this procedure is explained in [4]. The analysis was restricted to the $\sqrt{s'}$ region between 20 and 87 GeV.

In the data sample, 425 ISR events were selected for the cross-section calculation, and 534 for the differential cross-section and asymmetry calculation.

From the 691,335 simulated dimuons selected as described in Section 4.1, 2845 ISR events were selected for the cross-section calculations.

4.3 Efficiency of the selection procedure and purity of the ISR sample

The efficiency of the selection procedure and the contamination by FSR events were studied with the sample of simulated radiative muon events generated by DYMU3. The events were flagged as ISR or FSR on the basis of the generator information. If both initial and final state photons were produced, the events were labelled ISR if the generated final state photon energy was below 1 GeV and FSR otherwise. The effective annihilation energy $\sqrt{s'}$ was calculated using formula (12). The resulting efficiency and purity are displayed as a function of $\sqrt{s'}$ in Figure 1. The purity of the sample is near 90% over the whole energy interval. The selection efficiency is about 80% at low energies and decreases to around 50% at 87 GeV. Figure 1 shows that the purity of the data sample taken between 1991 and 1994 [4] is tha same as the purity of the sample of data taken in 1995. Therefore both samples can simply be merged to determine the forward-backward asymmetry and the helicity component ratios.

The cosmic ray background was checked using the sample of ISR events selected for the cross-section calculation. No additional events were found when the cuts on the interaction region definition were relaxed to allow impact parameters of charged particle tracks to the production vertex of up to 3 cm in the plane transverse to the beam and up to 9 cm parallel to the beam, which should have quadrupled this background.

The background from $e^+e^- \to \tau^+\tau^-$ events was estimated from the sample of 373,000 simulated $\tau^+\tau^-$ events (see Section 3). No events were found to satisfy the ISR selection criteria.

The background from two-photon processes was estimated from simulated events. It was found to be 1.6% for the channel $e^+e^- \to e^+e^-\mu^+\mu^-$, while no events in the channel $e^+e^- \to e^+e^-\tau^+\tau^-$ satisfied the ISR criteria.

5 Estimation of the total Improved Born cross-sections

The data taken in the years 1992–1995 were used to estimate the Improved Born cross-sections $\sigma_{IB}^{obs}(\sqrt{s'})$. The polar angle of the μ^- was required to be in the range $20° \leq \theta_{\mu^-} \leq 160°$. A total of 425 events was selected. The ratio of the averages of the observed to the Standard Model Improved Born cross-sections inside a given $\sqrt{s'}$ interval is given by

$$\frac{< \sigma_{IB}^{obs}(\sqrt{s'}) >}{< \sigma_{IB}^{SM}(\sqrt{s'}) >} = \frac{N_{obs}(\sqrt{s'}) \cdot N_{sim}^{norm}}{N_{sim}(\sqrt{s'}) \cdot N_{obs}^{norm}} \tag{13}$$

where $\sqrt{s'}$ is the mean effective annihilation energy in the interval. The quantities $N_{obs}(\sqrt{s'})$ and $N_{sim}(\sqrt{s'})$ represent the numbers of ISR events in a given $\sqrt{s'}$ interval in the data and in the simulated sample respectively, selected as explained in Section 4.2. The quantities N_{obs}^{norm} and N_{sim}^{norm} represent the number of dimuon events selected as explained in Section 4.1 in the real and simulated data samples. In each $\sqrt{s'}$ interval, the normalisation of the ISR sample to the full dimuon sample was calculated separately for the on-peak and off-peak data, after which the results were averaged. The number $N_{obs}(\sqrt{s'})$ was corrected for the two-photon background, and the number N_{obs}^{norm} was corrected for the background arising from cosmic ray and tau events. The other backgrounds were too small to justify a correction. It was verified that the selection efficiency for ISR events was the same for the observed data and the simulation.

The theoretical Improved Born cross-section, $\sigma_{IB}^{SM}(\sqrt{s'})$, was obtained from the DYMU3 program. The parameters used in this calculation were $M_Z = 91.25$ GeV/c^2, $\Gamma_Z = 2.562$ GeV/c^2, and $\sin^2\theta_W = 0.2296$, which were the default values used by DELPHI for the generation of $\mu^+\mu^-$ events. It was verified that the calculations made by this program and those made by the ZFITTER program [7] agreed to within 1%.

Table 2 shows the number of ISR events selected in the data and in the simulated samples as a function of $\sqrt{s'}$, up to an energy of 87 GeV/c^2. The same table shows the calculated SM Improved Born cross-section in the different $\sqrt{s'}$ intervals as well as the experimental cross-section obtained from formula (13). Only statistical errors were taken into account. The main source of systematic errors was the modelling of the muon momenta in the simulation. To reduce these effects to a negligible size, the muon momenta were smeared in the simulation to match the resolution observed in the data.

Figure 2 shows the ratios between the observed and theoretical Improved Born cross-sections as a function of the effective annihilation energy, and the resulting experimental cross-sections. It also shows the cross-sections for the reaction $e^+e^- \to \mu^+\mu^-$ obtained near the Z^0 peak [12, 14] and those obtained at PEP, PETRA, and TRISTAN [15, 3]. The published values were corrected to obtain Improved Born values. The cross-section

measured with the VENUS detector at 58 GeV has a value of 29.07 ± 0.66 pb [3], which has to be compared with the Standard Model prediction of 30.72 pb. The resolution of Figure 2 is not sufficient to show this deviation.

6 Asymmetries and helicity cross-sections

6.1 Polar angle distributions

The data taken in the years 1991–1995 were merged to obtain the distributions of the muon polar angle in the $\mu^+\mu^-$ rest frame for different $\sqrt{s'}$ intervals. A total of 534 events were selected. The distribution of these events as a function of $\sqrt{s'}$ is shown in Table 3.

For events which are not produced in the e^+e^- c.m. frame, the angle between the μ^- and the e^- beam direction in the $\mu^-\mu^+$ rest frame is given by [13]:

$$\cos\theta^* = \frac{\sin\frac{1}{2}(\theta_{\mu^+} - \theta_{\mu^-})}{\sin\frac{1}{2}(\theta_{\mu^+} + \theta_{\mu^-})} \tag{14}$$

where θ_{μ^+} and θ_{μ^-} are the polar angles of the μ^+ and the μ^- with respect to the e^- beam axis in the laboratory frame.

The raw distributions of $\cos\theta^*$ in the different $\sqrt{s'}$ intervals are shown in Figure 3. They were corrected for selection inefficiencies in the following way. It was assumed that, within the limits of the present statistics, forward-backward asymmetries in detection efficiency could be neglected. Indeed, such asymmetries distort the forward-backward asymmetry by at most 0.001 [14]. Therefore, in each $\sqrt{s'}$ interval of the data, the correction factor was obtained from the folded distribution in $|\cos\theta^*|$, which was fitted to a function of the form

$$\frac{dN_r}{d|\cos\theta^*|} = \frac{3}{8}\, C\, (1 + \cos^2\theta^*) \tag{15}$$

in the region of $|\cos\theta^*|$ where the fit gave the smallest value of χ^2/N_d. The C value obtained from this fit was used to calculate correction factors in the $|\cos\theta^*|$ bins outside the fit region, namely $F(|\cos\theta^*|) = \frac{3}{8}C(1+\cos^2\theta^*)/N_r(|\cos\theta^*|)$, where N_r is the raw number of events. In the unfolded $\cos\theta^*$ distribution, the raw bin contents for positive and negative $\cos\theta^*$ were corrected with the same correction factors $F(|\cos\theta^*|)$. The resulting corrected $\cos\theta^*$ distributions are also shown in Figure 3 for the five $\sqrt{s'}$ intervals considered.

In each $\sqrt{s'}$ interval, the $\cos\theta^*$ distribution was then corrected for the background coming from FSR, which on average amounted to about 10%. In each interval, a distribution of the form

$$\frac{dN_{FSR}}{d\cos\theta^*} = C'\,[\frac{3}{8}(1+\cos^2\theta^*) + A^0_{fb}\cos\theta^*] \tag{16}$$

was subtracted, where A^0_{fb} was taken to be the asymmetry at the Z^0 peak, from which this background dominantly arises. The constants C' were obtained from the integral of equation (16), which is equal to the expected number of FSR events in a given $\sqrt{s'}$ interval as predicted by the simulated radiative muon sample. The contamination by tau events, cosmics or two-photon events was too small to justify a correction.

6.2 Forward-backward asymmetries

In each $\sqrt{s'}$ interval, the asymmetry A_{FB} was obtained by performing a maximum likelihood fit of the uncorrected $\cos\theta^*$ distribution to an expression of the form

$$\frac{dN}{d\cos\theta^*} = C''[(1+\cos^2\theta^*) + \frac{8}{3}A_{FB}\cos\theta^*] \; . \tag{17}$$

Formula (17) does not include radiative corrections. Since the asymmetries determined in this analysis are Improved Born asymmetries, only the electro-weak corrections should be considered. Compared to the experimental precision they are small, and modify the asymmetry by at most 0.02 in the energy region between 40 and 88 GeV. Table 3 shows the asymmetries as a function of $\sqrt{s'}$ up to 87 GeV.

The raw asymmetries obtained with the fit were corrected for the contamination by FSR events in a similar way as the polar angle distributions were corrected (see Section 6.1). These corrected values are also shown in Table 3, and are displayed in Figure 4 together with the SM prediction for the Improved Born asymmetry. Figure 4 also shows the asymmetries measured by DELPHI near the Z^0 peak (see [12, 14]), after correction to Improved Born values. The SM Improved Born asymmetry was calculated with the DYMU3 program with the parameters mentioned in Section 5. The only source of systematic error on the asymmetry, $\delta(A_{FB})^{fit}_{sys}$, which was considered was that resulting from the error on the purity. The values of this error are shown in Table 3.

6.3 Extraction of the helicity cross-sections and test of the Standard Model.

From the $\cos\theta^*$ distributions corrected for selection efficiencies and for the FSR background, the cross-sections $\tilde{\sigma}^{\mu}_{+}$ and $\tilde{\sigma}^{\mu}_{-}$ (see Section 2.2) and their ratio were determined as follows. In each bin of the $\cos\theta^*$ distribution the corrected content was multiplied by a weight factor

$$F_{\pm} \;=\; A\,(1 \pm B\cos\theta^*) \tag{18}$$

where

$$A = \frac{2}{C_M(3+C_M^2)} \qquad B = \frac{3+C_M^2}{2C_M^2} \tag{19}$$

and $C_M = \cos\theta^*_{max}$. These weighted contents were summed for all $\cos\theta^*$ bins between -0.8 and $+0.8$ for $\sqrt{s'}$ below 65 GeV, and between -0.9 and $+0.9$ for the other $\sqrt{s'}$ values. The C_M limits were chosen to match the binning in each $\sqrt{s'}$ interval. The values of $\tilde{\sigma}^{\mu}_{+}$ and $\tilde{\sigma}^{\mu}_{-}$ and their statistical errors were derived from the weighted sums. For the systematic error, the error on the purity of the sample was taken into account. The numbers obtained are given as a function of $\sqrt{s'}$ in Table 3 and are shown in Figure 5, together with the prediction of the Standard Model. The theoretical predictions include all electroweak radiative corrections, apart from the box diagrams which are very small (see [2]). There is no indication of any deviation from the Standard Model.

7 Summary

Using the data collected by DELPHI until 1995, a sample of around 530 dimuon events with initial state radiation was selected. The differential and total cross-sections and forward-backward asymmetries were determined at energies between 20 and 87 GeV. No deviation from the Standard Model was observed.

The polar angle distributions were used to determine the ratio $\frac{\sigma_{LL}+\sigma_{RR}}{\sigma_{RL}+\sigma_{LR}}$, where the two subscripts represent the helicities of the incoming e^- and outgoing μ^- respectively. The aim was to test the Standard Model near the minimum of this ratio, around 80 GeV, where the sensitivity to new physics is greatest. No evidence for new physics was seen.

8 Acknowledgements

We are greatly indebted to our technical collaborators and to the funding agencies for their support in building and operating the DELPHI detector, and to the members of the CERN-SL Division for the excellent performance of the LEP collider.

We would like to thank J.-M. Frêre for suggesting the analysis in terms of helicity cross-sections, and for many fruitful discussions.

References

[1] ALEPH Collab., R. Barate et al., *Study of the muon-pair production at centre-of-mass energies from 20 to 136 GeV with the ALEPH detector*, CERN-PPE/97-10, subm. to Phys. Lett. **B**;
L3 Collab., M. Acciari et al., Phys. Lett. **B374** (1996) 331;
OPAL Collab., P.D. Acton et al., Phys. Lett. **B273** (1991) 338.

[2] J.-M. Frêre, V.A. Novikov, M.I. Vysotsky, Phys. Lett. **B386** (1996) 437.

[3] K. Miyabayashi, *Recent electroweak results from TRISTAN*, Proc. of the XXXth Rencontres de Moriond, Les Arcs, March 1995, ed. J. Trân Thanh Vân.

[4] DELPHI Collab., P. Abreu at al., CERN-PPE/97-45, submitted to Z. Phys. **C**.

[5] F. Berends, *Z lineshape*, in Z Physics at LEP I, CERN 89-08, Volume 1 (1989).

[6] D. Bardin et al., Nucl. Phys. **B351** (1991) 1;
DELPHI Collab., P. Abreu et al., Z. Phys. **C72** (1996) 31.

[7] D. Bardin et al., *ZFITTER, An Analytical Program for Fermion Pair Production in e^+e^- Annihilation*, CERN-TH. 6443/92 and hep-ph/9412201.

[8] DELPHI Collab., P. Aarnio et al., Nucl. Instr. & Meth. **A303** (1991) 233;
DELPHI Collab., P. Abreu et al., Nucl. Instr. & Meth. **A378** (1996) 57.

[9] J.E. Campagne and R. Zitoun, Z. Phys. **C43** (1989) 469;
and Proc. of the Brighton Workshop on Radiative Corrections, Sussex, July 1989.

[10] S. Jadach and Z. Was, Comp. Phys. Com. **36** (1985) 191;
S. Jadach, B.F. Ward and Z. Was, Comp. Phys. Com. **66** (1991) 276.

[11] J. Hilgart, R. Kleis, Comp. Phys. Com. **75** (1993) 191.

[12] DELPHI Collab., P. Abreu et al., Nucl. Phys. **B417** (1994) 3.

[13] Z. Was and S. Jadach, Phys. Rev. **D41** (1990) 1425.

[14] DELPHI Collab., P. Abreu et al., Nucl. Phys. **B418** (1994) 403.

[15] HRS Collab., M. Derrick et al., Phys. Rev. **D31** (1985) 2352;
MAC Collab., W. W. Ash et al., Phys. Rev. Lett. **55** (1985) 1831;
TASSO Collab., W. Braunschweig et al., Z. Phys. **C40** (1988) 163;
CELLO Collab., H.-J. Behrend et al., Phys. Lett. **B191** (1987) 209;
JADE Collab., W. Bartel et al., Z. Phys. **C26** (1985) 507;
MARK J Collab., B. Adeva et al., Phys. Rev. **D38** (1988) 2665;
AMY Collab., A. Bacala et al., Phys. Lett. **B218** (1989) 112;
AMY Collab., C. Velissaris et al., Phys. Lett. **B331** (1994) 227;
TOPAZ Collab., B. Howell et al., Phys. Lett. **B291** (1992) 206.

$\sqrt{s'}$ [GeV]	< 35	35-65	65-80	80-84	> 84
loose selections					
p_T^{miss} [GeV/c]	-	< 5	< 4	< 4	< 3
α_{acol} [°]	> 1.5	> 1.5	> 1.5	> 1	> 1
α_{acop} [°]	< 7.5	< 7.5	< 5	< 4	< 4
E_{em}^{neu} [GeV]	-	< 2	< 2	< 2	< 2
$\lvert\Delta E_\gamma\rvert$ [GeV]	< 25	< 13.2	< 6	< 6	< 5
tight selections					
p_T^{miss} [GeV/c]	-	< 2.5	< 1	< 1	< 1
α_{acol} [°]	> 3	> 3	> 3	> 2	> 2
α_{acop} [°]	< 5	< 5	< 4	< 3	< 3
E_{em}^{neu} [GeV]	-	< 1	< 1	< 1	< 1
$\lvert\Delta E_\gamma\rvert$ [GeV]	< 25	< 4.4	< 3	< 3	< 2.5

Table 1: Requirements for the selection of ISR events. The details are given in the text.

$\sqrt{s'}$ [GeV]	24-38	38-45	45-52	52-59	59-66	66-73	73-80	80-84	84-87
N_{obs}	8	21	14	15	16	28	49	71	203
N_{sim}	39	123	97	133	127	149	331	531	1315
$< \sigma_{IB}^{obs} > / < \sigma_{IB}^{SM} >$	1.11	0.95	0.88	0.79	0.69	1.11	0.90	0.89	1.10
	± 0.46	± 0.23	± 0.27	$\pm .22$	± 0.20	± 0.23	± 0.13	± 0.10	± 0.06
$< \sqrt{s'} >$ [GeV]	34.3	41.7	47.7	55.4	62.2	70.3	77.3	82.4	85.8
σ_{IB}^{SM} [pb]	83.1	56.5	43.6	33.1	27.5	24.9	29.4	49.0	106.1
σ_{IB}^{obs} [pb]	92.2	53.7	38.4	26.1	19.0	27.6	26.5	43.6	116.7
$\delta(\sigma_{IB}^{obs})$ [pb]	± 38.2	± 13.0	± 11.8	± 7.3	± 5.5	± 5.7	± 3.8	± 4.9	± 6.4

Table 2: Numbers of ISR events found in the data and simulated samples (N_{obs} and N_{sim}) for different $\sqrt{s'}$ intervals, and the ratios of the average measured to SM Improved Born cross-sections. Then $< \sqrt{s'} >$ is the mean measured effective annihilation energy in the interval, and σ_{IB}^{SM} is the mean Improved Born cross-section within each energy interval expected in the Standard Model, obtained from the DYMU3 program. Finally σ_{IB}^{obs} with its error $\delta(\sigma_{IB}^{obs})$ is the resulting measured cross-section.

$\sqrt{s'}$ [GeV]	20-50	50-65	65-80	80-84	84-87
N_F	36	16	16	21	72
N_B	29	27	78	70	169
$<\sqrt{s'}>$ [GeV]	42.5	58.9	74.4	82.4	85.9
$A_{FB}^{fit}unc$ [%]	13.2 ± 12.1	-34.5 ± 14.0	-62.3 ± 6.3	-57.1 ± 7.7	-40.8 ± 5.6
$A_{FB}^{fit}corr$ [%]	13.8 ± 12.6	-37.4 ± 15.1	-69.0 ± 7.0	-63.1 ± 8.6	-45.4 ± 6.3
$\delta(A_{FB})_{sys}^{fit}$[%]	± 0.4	± 0.6	± 0.7	± 0.6	± 0.3
$\tilde{\sigma}_+^\mu/\tilde{\sigma}_-^\mu$	1.61 ± 0.54	0.34 ± 0.17	0.08 ± 0.05	0.09 ± 0.06	0.22 ± 0.05
δ_{sys}	0.05	0.01	0.01	0.01	0.01

Table 3:
N_F, N_B: observed number of ISR events in the forward and backward hemispheres ;
$A_{FB}^{fit}unc$: the asymmetry with statistical error calculated with a maximum likelihood fit, without correction for FSR contamination ;
$A_{FB}^{fit}corr$: the asymmetry with statistical error calculated with a maximum likelihood fit, after correction for FSR contamination, with systematic error $\delta(A_{FB})_{sys}^{fit}$;
$\tilde{\sigma}_+^\mu/\tilde{\sigma}_-^\mu$: the helicity component ratio defined in Section 2.2, with its statistical error ;
δ_{sys}: the systematic error on the helicity component ratio.



Figure 1: **(a)** Efficiency for the selection of ISR events and **(b)** purity of the ISR sample with regard to FSR events, based on simulated radiative muon events, for the data sample used in ref. [4] (*black dots*) and for the data taken in 1995 (*stars*)



Figure 2: **(a)** Ratio of measured to SM Improved Born cross-section as a function of the effective annihilation energy. **(b)** Improved Born cross-sections measured in this analysis (*black circles*), at the Z^0 peak (*open circles*), and at PEP, PETRA and TRISTAN. The *solid line* shows the SM prediction.



Figure 3: Distribution of $\cos\theta^*$ in different intervals of the effective annihilation energy $\sqrt{s'}$. The *dashed lines* show the raw distributions and the *solid lines* the distributions after correction for the selection efficiency.



Figure 4: Improved Born asymmetry as a function of the effective annihilation energy. The *black squares* show the measurements made by DELPHI below 87 GeV, using the likelihood fit method described in the text. The *open circles* show the measurements made by DELPHI at the Z^0 peak. The *solid line* shows the SM prediction.



Figure 5: Ratio $\tilde{\sigma}^{\mu}_{+}/\tilde{\sigma}^{\mu}_{-} = (\sigma_{RR}+\sigma_{LL})/(\sigma_{LR}+\sigma_{RL})$ as a function of the effective annihilation energy. The *solid line* shows the SM prediction.