Measurement of the Structure Function $F_2(x, Q^2)$ of the Proton at Low Q^2 with the H1 Detector at HERA Using the New Detector Components Spacal and BDC

DISSERTATION

zur Erlangung des Doktorgrades
des Fachbereichs Physik
der Universität Hamburg

vorgelegt von

Andreas Meyer

aus Bielefeld

Hamburg
1997

Abstract

A measurement of the structure function $F_2(x, Q^2)$ of the proton is presented in the kinematic range of Q^2 between 0.85 and 25 GeV^2 and Bjorken-x between $1.3 \cdot 10^{-5}$ and $5 \cdot 10^{-3}$. The measurement is performed using data recorded with the H1 detector at the ep storage ring HERA in 1995. In the winter shutdown 1994/95 the H1 detector was upgraded with a new calorimeter 'Spacal' and a new drift chamber 'BDC'. These components are used in this analysis for the first time. One of the main topics of this thesis is the energy calibration of the Spacal calorimeter. With the help of the new detector components the precision of the measurement of F_2 is improved compared to previous results by roughly a factor of 2 in the Q^2-range between 2.5 and 6.5 GeV^2.

Kurzfassung

Die Arbeit beschreibt eine Messung der Strukturfunktion F_2 des Protons im kinematischen Bereich von $0.85 \leq Q^2 \leq 25$ GeV^2 und $1.3 \cdot 10^{-5} \leq x \leq 5 \cdot 10^{-3}$. Die Messung beruht auf Daten, die im Betriebsjahr 1995 mit dem H1 Detektor am HERA Speicherring aufgenommen wurden. Im Winter 1994/95 waren ein neues Kalorimeter 'Spacal' und eine Driftkammer 'BDC' im H1 Detektor installiert worden. Diese Detektorkomponenten werden in dieser Arbeit erstmals verwendet. Ein Schwerpunkt dieser Arbeit liegt auf der Energiekalibration des Spacal Kalorimeters. Mithilfe der neuen Detektorkomponenten wird die Präzision der Messung von F_2 im Vergleich zu früheren Ergebnissen im Bereich von $2.5 \leq Q^2 \leq 6.5$ GeV^2 um einen Faktor 2 verbessert.

Contents

Introduction

The scattering of leptons off nuclei, and later nucleons, has been for many decades one of the key measurements in the exploration of the structure of matter. As early as in the late sixties, measurements at SLAC led to the discovery of the spin-$\frac{1}{2}$ constituents of the proton which came to be identified as quarks. In the Quark Parton Model the sum of the momentum distributions of the quarks in the proton is directly related to the structure function F_2.

In the following years Quantum Chromodynamics (QCD) was developed as the field theory of strong interactions. In QCD the quarks interact via exchanged gauge bosons called gluons which, together with sea quarks, carry about half of the proton momentum. At sufficiently large momentum transfer, Q^2, the coupling between quarks and gluons is small, allowing the description of the scattering processes in perturbation theory. Perturbative QCD is able to describe the relative behavior of the parton momentum distributions under variations of Q^2 by virtue of evolution equations. Throughout the seventies and eighties the predictions of QCD have successfully been tested in a series of experiments continuously extending the kinematic range. Since, with increasing beam energy, the electron beams used in the early experiments at SLAC and Fermilab, became more and more difficult to produce, muon beams became a natural choice. Furthermore, neutrino nucleon scattering experiments have been pursued. Until 1992, data with excellent precision, covering a wide kinematic range down to values of Bjorken-$x \sim 10^{-4}$ and up to $Q^2 \sim 200$ GeV^2, have been produced in particular at SLAC (eN) and by the BCDMS, NMC and E665 (μN) as well as CCFR and CDHS (νN) Collaborations at CERN and at Fermilab. These data have been and continue to be the basis for deep insights into the structure of the proton and for improvements in the understanding of QCD.

The electron proton collider HERA at DESY, Hamburg, follows this long tradition of Deep Inelastic Scattering experiments. At HERA both, electrons and protons are accelerated and collide at a center of mass energy of 300 GeV allowing to resolve the structure of the proton down to distances of $\sim 10^{-18}$ m. Compared to former 'fixed target' experiments, where the nucleons probed by the leptons were at rest (in the laboratory frame) the center of mass energy is larger by more than one order of magnitude. HERA therefore provides a unique facility for the study of the structure of the proton in a new kinematic regime allowing measurements of the structure function down to values of Bjorken-$x \gtrsim 10^{-6}$ at values of $Q^2 \gtrsim 0.1$ GeV^2.

Experimentally, the structure function is extracted from the double differential ep cross section measured as a function of x and Q^2. At the two colliding beam experiments ZEUS and H1 not only the scattered lepton but also the hadronic final state is measured giving rise to alternative methods to reconstruct the event kinematics and to control possible systematic effects. Being fully inclusive the measurement of F_2 makes use of the full event statistics and is thus least affected by statistical uncertainties. Furthermore, the results of the inclusive measurement can be interpreted in QCD without theoretical assumptions on the exact behavior of the hadronic final state. As early as in the first year of HERA operation, 1992, a significant rise of the structure function towards small values of x was discovered. The dynamical behavior of the data was found to be well described by the DGLAP evolution equations which are based on perturbative QCD. However, the discrimination of different theoretical models and the precise determination of the parton distributions requires high precision. In the small x regime at HERA in particular the distribution of gluons in the proton can be determined. The continuous increase of the HERA luminosity has reduced the statistical uncertainties. In order to gain further precision it is necessary to optimize the detector performance and to control and minimize the systematic uncertainties of the measurement.

In the winter shutdown 1994/95 the H1 detector was upgraded. The detector components which cover the region of small electron scattering angles ($\lesssim 25°$ relative to the electron beam) were replaced by a new lead / scintillating-fiber calorimeter of spaghetti-type 'Spacal' and a Backward Drift Chamber 'BDC'. The new components improve the detector performance in particular for events with $Q^2 \lesssim 100$ GeV^2 and extend the accessible[1] kinematic range to Q^2-values of ~ 0.85 GeV^2 and $x \sim 10^{-5}$. In addition to a better energy and angular resolution the main improvements lie in the possibility to measure the hadronic final state and in the suppression of beam background on the first trigger level with the help of an excellent time resolution.

For the measurement presented here data are used which were recorded with the H1 detector in 1995, the commissioning year of the Spacal and the BDC. In the context of this analysis the behavior of the new detector components is explored in detail for the first time.

This thesis is organized as follows: Chapter 1 provides an introduction to the theory of Deep Inelastic Scattering and QCD. The physics objective of the analysis is motivated by relating theoretical developments to recent experimental results. The experimental environment of the H1 Detector is covered in chapter 2. A description of the most crucial detector components and the data acquisition is given. Special focus is put on the technical features of the new Spacal Calorimeter. Chapter 3 contains a detailed description of the energy calibration of the Spacal. In chapter 4 the measurement of the structure function is described. Besides a detailed discussion of the event selection based on the identification of the scattered electron in the Spacal a main topic of this chapter is the determination of the photoproduction background. Furthermore, a first investigation of QED final state radiation is performed. Finally, in chapter 5, the result of the measurement is presented. The structure function F_2 and the total virtual photon proton cross section are compared to model predictions currently available to describe the HERA data in the small-Q^2-region.

[1]with data of nominal interaction point position.

Chapter 1

Deep Inelastic Scattering

Inclusive cross section measurements have proven extremely fruitful for the understanding of the structure of matter. The relation of experimental results to theoretical concepts is most direct in the case of inclusive measurements where only the scattered lepton is measured and all possible hadronic final state configurations are summed over. Historically, by the time of the sixties, measurements of elastic electron proton scattering had led to the determination of the nucleon form factors which in the non-relativistic limit describe the charge distributions and the magnetic moments of protons and neutrons.

The notion of 'Deep Inelastic Scattering' (DIS) was coined with the advent of high energy data recorded at the *Stanford Linear Accelerator Center* (SLAC), as of 1967, when events with a very large energy loss of the scattered electron were observed. The famous result of these data was that the deep inelastic structure function F_2 showed very little dependence on the transferred four-momentum squared, Q^2, between the electron and the target, but depended only on the variable x which in the parton model can be interpreted as the fractional momentum of the struck proton constituent. Such a scale invariant behavior, simply called 'scaling', had been predicted by Bjorken already in 1966 [1]. The simplest explanation for scaling was given by the parton model, assuming the proton constituents to be free and point-like partons. Almost immediately, in 1969, the 'Quark Parton Model' (QPM) was established identifying the partons with the spin-$\frac{1}{2}$ quarks [2]. In the Quark Parton Model the structure function F_2 is related to the sum of the parton momentum distributions $x \cdot q_i(x)$ weighted with the square of their electric charge e_i,

$$F_2(x) = \sum_i e_i^2 x q_i(x). \tag{1.1}$$

In the years to follow Quantum Chromodynamics (QCD) was developed as the field theory of strong interactions between quarks and gluons. QCD solves the conceptual puzzle of the partons which, on the one hand are permanently bound in the proton and, on the other hand behave like free particles, introducing a scale dependent ('running') coupling constant which is large at small Q^2 ('confinement') and logarithmically vanishes towards large Q^2 ('asymptotic freedom').

Until the late eighties various fixed target experiments were performed at center of mass energies up to 30 GeV providing a wealth of data by which the structure of the proton could be constrained very precisely for values of $x \gtrsim 10^{-3}$ and $Q^2 \lesssim 200$ GeV^2.

At the ep storage ring HERA the center of mass energy is increased by one order of magnitude to a value of ~ 300 GeV. An entirely new kinematic regime of Deep Inelastic Scattering is thus opened up, extending down to values of $x \gtrsim 10^{-6}$ and Q^2 values of up to 90000 GeV^2. At HERA, besides the scattered electron, the hadronic final state is also measured allowing the use of different independent methods to reconstruct the event kinematics. The HERA research program also covers detailed studies of the properties of the hadronic final state. A concise review of HERA physics can be found in [3].

In this chapter the variables describing the event kinematics of inclusive deep inelastic scattering processes are explained (section 1.1) and the relations between the measured double differential ep cross section and the structure functions are defined (section 1.2). A brief outline of the interpretation of the structure function F_2 in the framework of QCD is given (section 1.3). The present theoretical understanding of ep scattering is confronted with recent experimental results in section 1.4. In this context, the objective of the presented measurement is discussed (section 1.5).

1.1 Event Kinematics

The process of inclusive electron proton scattering, $ep \to lX$ is visualized in fig.1.1(a). The interaction between the electron and the proton, to lowest order in the single boson exchange approximation, takes place via an exchanged gauge boson. The exchanged boson carries the four-momentum q, and p stands for the four-momentum of the incoming proton. The four-momenta of the incoming and the scattered lepton are denoted by k and k', respectively. The system X with the four-momentum p' represents the hadronic final state, i.e. the particles produced in the break-up of the proton.

A distinction is made between two classes of events, namely Neutral Current (NC) events, $e^{\pm}p \to e^{\pm}X$, with a neutral virtual photon or Z^0 exchange; and Charged Current (CC) reactions, $e^{\pm}p \to \overset{(-)}{\nu} X$, with a $W^{\pm}$ exchange and a neutrino as outgoing lepton.

At fixed beam energy the event kinematics is unambiguously defined by two of the three Lorentz invariant variables,

$$Q^2 = -q^2 = -(k-k')^2 \tag{1.2}$$

$$x = \frac{Q^2}{2p \cdot q} \tag{1.3}$$

$$y = \frac{p \cdot q}{p \cdot k} \tag{1.4}$$



Figure 1.1: (a) Schematic representation of inclusive electron proton scattering $ep \to lX$ in the single boson exchange approximation. The four-momenta of the incoming and scattered lepton are k and k', respectively. The incident proton four-momentum is p; p' denotes the four-momentum of the hadronic final state X. The exchanged boson carries the four-momentum q. (b) Representation of ep scattering in the Quark Parton Model. Here the virtual photon interacts with a free pointlike (quark-)parton and the remaining partons ('spectators') do not undergo any interaction. In the infinite momentum frame the incident momentum of the 'Struck Quark' is given by xp.

Q^2, x and y offer the following suggestive interpretations: Q^2 denotes the negative four-momentum squared of the exchanged boson, i.e. its virtuality. In the limit of $Q^2 = 0$ the exchanged photon is called real and its life time becomes large. At a given Q^2 the distance d resolved at fixed x can be estimated following the Heisenberg uncertainty relation $d \cdot |\vec{q}| \sim 1$ yielding $d \sim 2M_p x/Q^2$. Thus, with increasing Q^2 the virtual boson probes smaller and smaller distances d within the proton. For the maximum Q^2 kinematically possible at HERA a value of $\sim 10^{-18}$ m is obtained. In the infinite momentum frame, i.e. in a frame where the incident proton energy is very large compared to the proton mass M_p, x is interpreted as the fraction of proton momentum carried by the struck quark (see fig. 1.1(b)). y gives the fraction of the incident electron energy transferred from the electron to the proton in the proton rest frame, i.e the fractional electron energy carried by the virtual photon. Since the transferred energy is absorbed in the final state of the proton, y quantifies the inelasticity of the event.

It is convenient to define two further quantities:

$$s = (k+p)^2 \tag{1.5}$$

$$W^2 = (q+p)^2. \tag{1.6}$$

s denotes the square of the ep center of mass energy. W gives the center of mass energy of the photon proton system which is equal to the invariant mass of the hadronic final state. At the HERA beam energies, $E_e = 27.5$ GeV for electrons and $E_p = 820$ GeV for protons, the masses of the incident particles can be neglected, yielding the following relations:

$$s = 4 \cdot E_e \cdot E_p \tag{1.7}$$

$$W^2 = \frac{1-x}{x} \cdot Q^2. \tag{1.8}$$

In this approximation the kinematic variables x, y and Q^2 are related via the center of mass energy s:

$$Q^2 = sxy. \tag{1.9}$$

1.2 Inclusive Cross Section and Structure Functions

The cross section of the NC process $ep \to eX$ is written as

$$\sigma \propto L_{\mu\nu} W^{\mu\nu}. \tag{1.10}$$

In this equation the leptonic tensor $L_{\mu\nu}$ describes the interaction between the electron and the exchanged virtual boson. The hadronic tensor $W_{\mu\nu}$ accounts for the scattering of the exchanged boson with the proton. For single photon exchange $L_{\mu\nu}$ is precisely calculable in QED, $W_{\mu\nu}$ is unknown because it contains the internal structure of the proton which is as yet not calculable and must therefore be determined from experiments.

Using Lorentz invariance and current conservation one can, reduce $W_{\mu\nu}$ into three structure functions $F_i(x, Q^2)$, obtaining

$$\frac{\mathrm{d}^2\sigma^{e^\pm p}}{\mathrm{d}x\mathrm{d}Q^2} = \frac{4\pi\alpha^2}{xQ^4}\left[y^2 x F_1(x,Q^2) + (1-y)F_2(x,Q^2) \mp (y - \frac{y^2}{2})xF_3(x,Q^2)\right]. \tag{1.11}$$

A detailed derivation of this formula can be found in [4]. The term xF_3 describes the parity violating contribution from Z^0 exchange and interference terms between γ and Z^0 exchange. It is negligible at low and moderate Q^2 due to the large mass of the Z^0 ($M^2_{Z^0} \sim 10^4$ GeV^2) and becomes measurably large at large values of Q^2. At $Q^2 \lesssim 100$ GeV^2 the contribution is smaller than 0.1%. Consequently, neglecting the term xF_3 for low and moderate Q^2, the equation (1.11) can be reformulated as

$$\frac{\mathrm{d}^2\sigma^{e^\pm p}}{\mathrm{d}x\mathrm{d}Q^2} = \frac{2\pi\alpha^2}{xQ^4}\left[(2(1-y)+y^2)F_2(x,Q^2) - y^2 F_L(x,Q^2)\right] \tag{1.12}$$

$$\text{where} \qquad F_L = F_2 - 2xF_1. \tag{1.13}$$

This expression is called Born cross section.

The ep scattering processes can be viewed as interaction of a flux of virtual photons with the proton[1]. Virtual photons, having an effective mass, may have transverse

[1]The concept of a virtual photon flux is suggestive in particular at small x ($x \ll 10^{-2}$) where the life time, $t = 1/(2M_p x)$, of the exchanged virtual photon in the proton rest frame is large compared to the interaction time with the proton [3].

as well as longitudinal polarization. The double differential ep cross section can be decomposed accordingly,

$$\frac{\mathrm{d}^2\sigma^{e^\pm p}}{\mathrm{d}x\mathrm{d}Q^2} = \Gamma(y)\left(\sigma_T(x,Q^2)+\epsilon(y)\sigma_L(x,Q^2)\right) \tag{1.14}$$

$$= \Gamma(y)\sigma^{\mathrm{eff}}_{\gamma^* p}(x,y,Q^2) \tag{1.15}$$

where $\Gamma(y) = \frac{\alpha(2-2y+y^2)}{2\pi Q^2 x}$ describes the photon flux[2] and $\epsilon(y) = \frac{2(1-y)}{2-2y+y^2}$ is the photon polarization.

σ_T and σ_L respectively correspond to the cross sections of transversely and longitudinally polarized photons with the proton. They add to the total virtual photon cross section

$$\sigma^{\mathrm{tot}}_{\gamma^* p} = \sigma_T + \sigma_L. \tag{1.16}$$

At small x, in the HERA regime, σ_T and σ_L are related to the structure functions as

$$\sigma_T + \sigma_L = \frac{4\pi^2\alpha}{Q^2}F_2 \tag{1.17}$$

$$\sigma_L = \frac{4\pi^2\alpha}{Q^2}F_L. \tag{1.18}$$

The ratio R of σ_L and σ_T is defined as

$$R = \frac{\sigma_L}{\sigma_T} = \frac{F_L}{F_2 - F_L}. \tag{1.19}$$

Due to its relation to longitudinally polarized photons, given in equation (1.18), F_L is called Longitudinal Structure Function. Equation (1.12) shows that the sensitivity of the cross section on F_L is kinematically suppressed with y^2. At low $y \lesssim 0.4$ the effect of F_L on the extracted value of F_2 is therefore negligible ($\lesssim$ 2%).

1.3 Outline of QCD

Quantum Chromodynamics (QCD) is the field theory of the strong interactions between quarks and gluons. A detailed discussion of QCD can be found in [6]. In QCD each quark can appear in three different states called colors (red, green, blue). The strong interaction between quarks is mediated by gluons which carry both a color and an anticolor at the same time. Due to this property gluons are able to couple to gluons (self-coupling). This is in contrast to QED where photon self-coupling does not occur.

[2]It should be emphasized that the definition of a virtual photon flux is somewhat arbitrary since the definition is only constrained in the photoproduction limit ($Q^2 \to 0$) in which the real photon flux must be described, i.e. $\lim_{Q^2\to 0}\sigma^{\mathrm{tot}}_{\gamma^* p}(Q^2,W^2) = \sigma^{\mathrm{tot}}_{\gamma p}(W^2)$ must be satisfied. In this thesis the so-called Hand convention [5] is used.

In QED, due to screening effects from vacuum polarizations, the coupling constant α increases slowly with increasing Q^2. In QCD the self coupling of gluons works in the opposite direction producing antiscreening which leads to a logarithmic decrease of $\alpha_S(Q^2)$ with increasing Q^2 called 'running of $\alpha_S(Q^2)$'. The behavior of α_S is described by the renormalization group equation and the QCD β-function

$$\mu \frac{\delta \alpha_S(\mu)}{\delta \mu} = \alpha_s \cdot \beta(\alpha_S) = -\frac{\beta_0}{2\pi}\alpha_S^2 - \frac{\beta_1}{4\pi^2}\alpha_S^3 - \frac{\beta_2}{64\pi^3}\alpha_S^4 - ... \tag{1.20}$$

$$\text{with} \quad \beta_0 = 11 - \frac{2}{3}N_f; \quad \beta_1 = 51 - \frac{19}{3}N_f; \quad \beta_2 = 2857 - \frac{5033}{9}N_f + \frac{325}{27}N_f^2.$$

N_f is the number of quark flavors with mass less than the energy scale μ. A solution to equation (1.20) is given as an expansion in inverse powers of $\ln(Q^2)$ as

$$\alpha_S(Q^2) = \frac{4\pi}{\beta_0 \ln \frac{Q^2}{\Lambda^2}} \times \left[1 + \text{terms of higher order in } \ln(\mathrm{Q}^2/\Lambda^2)\right]. \tag{1.21}$$

Λ defines the strength of the coupling and must be determined experimentally. It is of the order[3] of 200 MeV [7]. Equation (1.21) is valid where Q^2 is large compared to Λ^2. It accommodates the running properties of α_S with Q^2 between the region of asymptotic freedom $\alpha_S \to 0$ as $Q^2 \to \infty$ and confinement where α_S is large and the partons are tightly bound in the proton.

The QCD factorization theorem states that the interaction can be split into two independent regions, a perturbative ('short distance') part with sufficiently large momentum scale and a non-perturbative ('long distance') part. In the perturbative part the relative smallness of $\alpha_S(Q^2)$ allows the expansion of solutions of QCD in a perturbation series of powers of $\alpha_S(Q^2)$. The non-perturbative part can in general only be determined experimentally. Formally, factorization means that the structure function F_2 can be expressed as a convolution of a coefficient function C_2^i calculable in QCD and the parton distribution functions f_i which are specific to the probed hadron but universal as regards the probing particle,

$$F_2(x, Q^2) \propto \sum_{f_i = q, \bar{q}, g} \int_x^1 dy \; C_2^i \left(\frac{x}{y}, \frac{Q^2}{\mu^2}, \frac{\mu_F^2}{\mu^2}, \alpha_S(\mu^2)\right) f_i(y, \mu_F, \mu). \tag{1.22}$$

The factorization scale parameter μ_F defines the boundary between the perturbative and non-perturbative regions and μ is the renormalization scale parameter used to absorb divergent parts of the perturbation series.
In a physical interpretation factorization means that the structure function F_2 is composed of two contributions, namely a probability density $f_i(y)$ of finding a parton with fractional proton momentum y in the proton and a coefficient term $C_2^i(\frac{x}{y})$ [8]. $C_2^i(\frac{x}{y})$ describes how a parton with fractional momentum x evolves from radiative processes

[3] The exact measured value depends on the renormalization scheme, i.e. the chosen convention.

off the initial parton and subsequently interacts with the exchanged virtual boson. Since the longitudinal momentum of the radiated parton is always smaller or equal to that of the parent parton, the integration in equation (1.22) is restricted to the interval $x \leq y \leq 1$. In this picture F_2 at a given value of x can be viewed as receiving contributions from all partons with a fractional momentum $y \geq x$ in the proton.

Since F_2 is a measurable quantity it should be independent of scale parameters. This means that the parton distribution functions f_i must depend on the choice of μ_F and μ, i.e. the choice of the renormalization scheme. A frequently used scheme is the DIS scheme which is defined by the Quark Parton Model relation between F_2 and the parton momentum distributions $f_i(x)$, given in equation (1.1),

$$F_2(x) = \sum_{i=q,\bar{q}} e_i^2 x f_i(x), \tag{1.23}$$

is satisfied to all orders, i.e. $C_2^{\mathrm{DIS}}(x) = e_i^2 \delta(1-x)$. The DIS scheme can be understood as defining perfect freedom of the partons in the proton as conceptualized in the Quark Parton Model. In this scheme all higher order processes are absorbed in the long range parton distribution functions constituting the structure of the proton. Another scheme, more useful for theoretical calculations, is the $\overline{MS}$ scheme where the above equation holds to leading order (LO) only. In NLO in the $\overline{MS}$ scheme, F_2 is also directly connected with the distribution of gluons in the proton.

The change of the parton distributions as a function of Q^2, i.e. their dynamical behavior or evolution, is described by the DGLAP evolution equations[4][9, 10] given in leading order as:

$$\begin{aligned}
\frac{\mathrm{d}q_i(x,Q^2)}{\mathrm{d}\ln Q^2} &= \frac{\alpha_S(Q^2)}{2\pi}\int_x^1 \frac{\mathrm{d}y}{y}\left[\sum_j q_j(y,Q^2)P_{ij}^{(0)}\left(\frac{x}{y}\right) + g(y,Q^2)P_{ig}^{(0)}\left(\frac{x}{y}\right)\right] \qquad (1.24)\\
\frac{\mathrm{d}g(x,Q^2)}{\mathrm{d}\ln Q^2} &= \frac{\alpha_S(Q^2)}{2\pi}\int_x^1 \frac{\mathrm{d}y}{y}\left[\sum_j q_j(y,Q^2)P_{gj}^{(0)}\left(\frac{x}{y}\right) + g(y,Q^2)P_{gg}^{(0)}\left(\frac{x}{y}\right)\right].
\end{aligned}$$

The splitting functions $P_{ij}(x/y)$ give the probability that a parton j with fractional proton momentum y radiates a parton i with fractional parton momentum x/y. The integration is restricted to $x \leq y \leq 1$ for the same reason as in equation (1.22). The splitting functions are calculable by perturbative expansion,

$$P_{ij}(z,\alpha_S(Q^2)) = \frac{\alpha_S}{2\pi}P_{ij}^{(0)}(z) + \left(\frac{\alpha_S}{2\pi}\right)^2 P_{ij}^{(1)}(z) + ... \tag{1.25}$$

In leading order the splitting functions read:

$$P_{qq}^{(0)}(z) = \frac{4}{3}\left[\frac{1+z^2}{1-z}\right]_+ + 2\delta(1-x) \tag{1.26}$$

[4]DGLAP stands for Dokshitzer, Gribov, Lipatov, Altarelli, Parisi.

$$\begin{aligned}
P_{qg}^{(0)}(z) &= \frac{1}{2}\left[z^2+(1-z)^2\right] \\
P_{gq}^{(0)}(z) &= \frac{4}{3}\frac{1+(1-z)^2}{z} \\
P_{gg}^{(0)}(z) &= 6\left[\left(\frac{z}{1-z}\right)_+ + \frac{1-z}{z} + z(1-z) + \frac{11}{12}\delta(1-x)\right] - \frac{N_f}{3}\delta(1-x).
\end{aligned}$$

Here the definition $\int_0^1 dx \frac{f(x)}{(1-x)_+} = \int_0^1 dx \frac{f(x)-f(1)}{(1-x)}$ is used [7]. To date the splitting functions have been calculated to next to leading order (NLO) [4].

The formal derivation of the DGLAP equations involves the Leading Log Approximation (LLA) where terms of the form $\alpha_S^n \cdot (\ln Q^2)^n$ are summed to all orders. These terms are leading because Q^2 is large by construction of the perturbative ansatz. It has been shown [11] that the LLA summation is equivalent to a sum of gluon ladder diagrams with decreasing fractional proton momentum x and strongly ordered transverse momentum,

$$x < x_n < \dots < x_0 \tag{1.27}$$

$$Q^2 \gg k_{T,n}^2 \gg \dots \gg k_{T,0}^2. \tag{1.28}$$

In the DGLAP approach x must be sufficiently large to ensure that terms of the type $\ln(1/x)$ of the splitting functions remain negligible. At low x it seems appropriate to include leading powers of $\ln(1/x)$ when accompanied by leading $\ln Q^2$ terms. This approach is called double leading logarithmic approximation (DLLA). The summation of $\ln(1/x)$ terms independent of Q^2 and at fixed $\alpha_S(Q^2)$ has been developed by BFKL[5] [12]. In leading order BFKL predict a behavior of F_2 proportional to $x^{-\lambda}$ with $\lambda \approx 0.5$. NLO calculations are still being pursued. In the limit of $s \to \infty$, i.e. $x \to 0$, a power behavior can not persist physically since the cross section must rise slower than $\ln^2 s$ (Froissart bound [13]).

Theory leaves as yet a number of questions to be answered by the experiments. Here, only the aspects most relevant to the present analysis are mentioned: The double differential ep cross section and the structure function $F_2(x, Q^2)$ is presently not calculable but can only be determined experimentally. Once the cross section is known as a function of x and Q^2 in a given kinematic region the evolution equations can be used for predictions of the cross section dependence outside the measured region. Furthermore, the experimental results can be used to extract the parton distribution functions f_i which describe the structure of the proton as a function of x and Q^2. High precision F_2 data at HERA provide a means of determining in particular the gluon density $xg(x, Q^2)$ and the strong coupling constant $\alpha_S(Q^2)$ [14]. As described above, towards low x and low Q^2 the perturbative approach will reach its limit of applicability and non-perturbative effects start to dominate. To date this region of transition, in particular in the low-x regime, has not yet been explored in detail.

[5]BFKL stands for Balitski, Fadin, Kuraev and Lipatov.

1.4 Recent Results

Measurements of F_2 using the HERA data recorded in 1994 [15, 16] showed that the next to leading order (NLO) DGLAP equations describe the data in the wide range from $Q^2 = 5000$ GeV^2 down to the lowest measured values of $Q^2 = 1.5$ GeV^2 within the achieved level of precision. This is illustrated in fig. 1.2 where data from HERA (here the H1 Experiment [15]) are presented together with results from earlier fixed target experiments [17, 18]. F_2 is shown as a function of x in different bins of Q^2. A strong rise of F_2 is seen towards the lowest values of x in all bins of Q^2. The solid line shows the result of the global H1 QCD fit [15, 19] based on the DGLAP evolution equation. QCD fits are performed parametrizing the parton distributions as functions of x at a given starting scale Q_0^2. The parameters are fit to the data such that the DGLAP equations and a set of additional constraints (e.g. QCD momentum sum rules) are obeyed. In the H1 fit [15] the data from H1 and from fixed target experiments are combined in order to achieve maximum possible precision.

The same results are presented in fig. 1.3 showing the total virtual photon proton cross section $\sigma^{\rm tot}_{\gamma^* p}$ as a function of W^2 in bins of Q^2. The strong rise of F_2 towards low x is reflected here in the increase of the cross section with W^2. Note that at low x the relation $W^2 = Q^2/x$ holds (see equation (1.8)). The perturbative model by Glück, Reya and Vogt (GRV) [20], based on the NLO DGLAP evolution equation, succeeds in describing the behavior for all values of $Q^2 \geq 1.5$ GeV^2 (see also section 5.4).

In addition to the previous figure the plot contains the total cross section $\sigma^{\rm tot}_{\gamma p}$ of photoproduction events at $Q^2 = 0$. In this region perturbation theory is not applicable. The line through the data points at $Q^2 = 0$ represents a phenomenological model by Donnachie and Landshoff (DOLA) [21] following a Regge theory type ansatz. An introduction to Regge theory can be found in [22]. DOLA parametrize the energy dependence of the cross section as

$$\sigma^{\rm tot}(W^2) = A \cdot W^{2\epsilon} + B \cdot W^{2\lambda}. \tag{1.29}$$

They predict $\epsilon = -0.4524$ and $\lambda = 0.0808$ from a fit to experimental data of hadron-hadron collisions. It turns out that the γp cross section measured at HERA [23, 24] obeys the same description[6]. At high energies W the 'pomeron' term $B \cdot W^{2\lambda}$ dominates and describes the slow increase of the total photoproduction cross section with the center of mass energy (see also section 5.4).

It is obvious from fig. 1.3 that the energy dependence of the photoproduction data and the data at $Q^2 \geq 1.5$ GeV^2 (DIS) is strikingly different. A measurement of the slope of the ep DIS cross section as $W^{2\lambda}$ at fixed Q^2 yields values of $\lambda \sim 0.2 ... 0.4$ continuously rising towards larger Q^2 [15] in contrast to ~ 0.08 in the case of the photoproduction data with $Q^2 = 0$.

[6] The fact that the energy dependences of hadron-hadron and γp (as well as $\gamma\gamma$) cross sections follow the same exponential behavior is often referred to as 'pomeron universality'.



Figure 1.2: The structure function F_2 as a function of x in bins of Q^2. The results of the H1 Collaboration obtained from data recorded in 1994 [15] are shown by the full points. The fixed target experiments NMC [18] (BCDMS [17]) are represented by open points (squares), respectively. The curves show the results of a NLO QCD fit which includes data for $Q^2 \geq 5$ GeV^2. The dashed lines show the extrapolation of the fit to lower values of Q^2.



Figure 1.3: The total virtual photon proton cross section as a function of W^2 for different bins in Q^2. Open (full) squares denote results from the H1 (ZEUS) Collaboration [23, 24]. At $Q^2 \geq 1.5$ GeV^2 the curves describe the model by GRV [20]. At $Q^2 = 0$ results from total photoproduction cross section measurements are shown. The curve through these points represents the parametrization by DOLA [21].

How does the transition between DIS and photoproduction take place? How can phenomenological cross section models be related to perturbative QCD? These questions are subject of much theoretical debate at present. Various model predictions have been developed to predict the structure function in the region of small Q^2 and small x. A measurement of the structure function F_2 and the energy behavior of the cross section in the transition region at very low values of Q^2 may help to discriminate between the models and to give constraints for further theoretical developments.

1.5 Objective of the Present Analysis

In the winter shutdown 94/95 at the H1 Detector at HERA a substantial increase in the detector acceptance for smaller scattering angles, and hence for smaller values of Q^2, was achieved with the installation of the new detector components 'Spacal' and 'BDC' (see chapter 2).

In the commissioning year 1995 two data sets were recorded. In addition to a sample of data with standard ep beam positions ('nominal vertex data'), a run dedicated to the measurement of the transition region at low Q^2 was performed. Here, the position of the main ep interaction region was shifted by the HERA machine by +70 cm along the proton direction ('shifted vertex data'). This way an acceptance in Q^2 down to $Q^2 \geq 0.35$ GeV^2 was achieved allowing to explore the transition region at very small values of x ($x \geq 5 \cdot 10^{-6}$)[7]. Both data sets have been used to measure the structure function F_2. The results of the shifted vertex data analysis have been published in [25]. Details of the analysis can also be found in [26, 27].

This thesis concentrates on the analysis of the nominal vertex data. Although these data extend the kinematic region compared to results from earlier years, the main objective of the measurement is not centered on the kinematic reach but rather on the precise understanding of the new detector components. In the region of Q^2 between 2 GeV^2 and 6.5 GeV^2 the experimental uncertainty of the results is improved compared to former measurements by roughly a factor of 2.

[7]The ZEUS Collaboration has presented results from 1995 data measured with the new beam pipe calorimeter (BPC) reaching down to Q^2 values of 0.11 GeV^2 covering a slightly different range in x [28].

Chapter 2

The H1 Experiment at HERA

In this chapter the experimental steps of the acquisition and reconstruction of deep inelastic ep scattering events with the H1 Detector at HERA are described. The electron proton storage ring HERA is introduced in section 2.1. The detector components of the H1 Detector and the basic steps of the data acquisition are covered in section 2.2. Special focus is put on the new backward calorimeter Spacal which together with the Backward Drift Chamber (BDC) plays a central role in this analysis (section 2.3). The standard methods used to reconstruct the event kinematics are explained (section 2.4) and the event simulation using Monte Carlo programs is sketched (section 2.5).

2.1 The *ep* Storage Ring HERA

The *Hadron-Elektron-Ring-Anlage* HERA [29] was commissioned in 1992. A schematic view of the HERA ring is shown in fig. 2.1. HERA has a circumference of 6.3 km and collides protons of 820 GeV with 27.5 GeV electrons or positrons. During the first years HERA operated with electrons until in 1994 the operation with positrons was established[1]. The lifetime of the positron beams is longer than that of electrons since the positron beam repels the remaining (positively charged) beam gas ions reducing the beam gas interactions and thus the loss of particles. The HERA particle beams are segmented in 220 bunch positions resulting in bunch crossing intervals of 96 ns corresponding to a rate of 10.4 MHz. In a typical run about 190 bunches are filled with about 10^{10} colliding particles each. The remaining ('pilot' and 'empty') bunches are used for control purposes. The average longitudinal size of the proton bunches is about 60 cm leading to a Gaussian distribution of the interaction points along the beam axis with a spread of 10 cm.

HERA accommodates four high energy physics experiments, namely H1, ZEUS, HERMES and HERA–B. H1 (a in fig. 2.1) and ZEUS (b) are taking data of ep collisions since 1992. HERMES (c) is operated since 1995 and uses the electron beam only.

[1] In the following and throughout this thesis the incident and scattered lepton is referred to as electron.



Figure 2.1: Sketch of the *ep* storage ring HERA (right picture) at DESY, Hamburg. HERA accommodates four experiments, H1 (a), ZEUS (b), HERMES (c) and HERA-B (d). The left sketch shows a zoom with the preaccelerator storage ring PETRA and linear preaccelerators.

Polarized electrons are collided with a polarized gas target (H_2, deuterium, ^{3}He or ^{4}He). The experiment HERA-B (d) is a fixed target experiment which uses the halo of the proton beam for collisions with tungsten wires. HERA-B is presently being commissioned and plans to measure CP violation in the decays of B^0 and $\overline{B^0}$ as of 1999.

2.2 The H1 Detector

The H1 Detector [30] was optimized to measure the energies and momenta of the particles produced in deep inelastic scattering and to allow efficient particle identification. A 3-dimensional sketch of the H1 Detector is given in fig. 2.2. The detector is nearly hermetic, its acceptance for scattered particles is limited only by the forward and backward beam pipe holes (near $\boxed{12}$ and $\boxed{13}$)[2]. Due to the difference between the beam energies of protons and electrons the momenta of most scattered particles have a large component in proton beam (forward) direction. The geometry of the detector, consequently, is asymmetric. The graph $\boxed{16}$ shows the coordinate convention used by the H1 collaboration and in this analysis. The z-axis points to the 'forward' direction along the proton beam. The polar angle is defined with respect to this direction. Looking backwards from the interaction point the x-axis points to the right towards the center of the HERA ring.

The central detector components, described below, are embedded in a superconducting coil $\boxed{6}$, which produces a homogeneous magnetic field of 1.2 T and an iron return yoke $\boxed{10}$, instrumented with streamer tubes for muon detection. In the forward direction

[2]Numbers in $\boxed{}$ refer to fig. 2.2.