Abstract ID: 791

Combination of F2\cc from DIS measurements at HERA

Content:

The charm contribution F2^cc to the proton structure function F2 is determined. The results of D meson production cross section measurements are combined with the measurements using semi-leptonic decays into muons as well as inclusive track measurements. The correlations of the systematic uncertainties between different measurements are taken into account. The data cover the kinematic range of photon virtuality $2 < Q^2 < 1000 \text{ GeV}^2$ and Bjorken scaling variable $10^{-5} < x < 10^{-1}$.

Primary authors: Dr. HAAS, Tobias (DESY); Dr. KRüGER, Katja (Universität Heidelberg)

Co-authors: Dr. REISERT, Burkard (Max-Planck Institut für Physik München); Dr. GEISER, Achim

(DESY); Prof. TASSI, Enrico (Universita della Calabria)

Presenter: Dr. HAAS, Tobias (DESY)

Track classification: 03 - Perturbative QCD, Jets and Diffractive Physics; 04 - Hadronic Structure, Parton

Distributions, soft QCD, Spectroscopy

Contribution type: Parallel Session Talk

Submitted by : Mr. HAAS, Tobias Submitted on Friday 14 May 2010

Last modified on: Friday 14 May 2010

Comments:

These results are presented on behalf of the ${\rm H1}$ and ${\rm ZEUS}$ Collaborations

Thursday 20 May 2010 Page 38