

Matching Threshold & Continuum for exclusive top pairs




Jürgen R. Reuter, DESY



in collaboration with: Fabian Bach, Bijan Chokoufè Nejad, Andre H. Hoang, Wolfgang Kilian, Thomas Teubner, Maximilian Stahlhofen, Christian Weiss
arXiv:1712.02220 [JHEP 1803(2018)184]

HELMHOLTZ
RESEARCH FOR GRAND CHALLENGES






Top Threshold/Continuum at lepton colliders

- LC top threshold scan best-known method to measure top quark mass, $\Delta M \sim 30\text{-}70$ MeV
- LC continuum top production best-known method to measure top couplings

error source	Δm_t^{PS} [MeV]
stat. error (200 fb ⁻¹)	13
theory (NNNLO scale variations, PS scheme)	40
parametric (α_s , current WA)	35
non-resonant contributions (such as single top)	< 40
residual background / selection efficiency	10 – 20
luminosity spectrum uncertainty	< 10
beam energy uncertainty	< 17
combined theory & parametric	30 – 50
combined experimental & backgrounds	25 - 50
total (stat. + syst.)	40 – 75


from 1702.05333



Top Threshold/Continuum at lepton colliders


- LC top threshold scan best-known method to measure top quark mass, $\Delta M \sim 30\text{-}70 \text{ MeV}$
- LC continuum top production best-known method to measure top couplings

Heavy quark production at lepton colliders, qualitatively:



error source	Δm_t^{PS} [MeV]
stat. error (200 fb ⁻¹)	13
theory (NNNLO scale variations, PS scheme)	40
parametric (α_s , current WA)	35
non-resonant contributions (such as single top)	< 40
residual background / selection efficiency	10 – 20
luminosity spectrum uncertainty	< 10
beam energy uncertainty	< 17
combined theory & parametric	30 – 50
combined experimental & backgrounds	25 – 50
total (stat. + syst.)	40 – 75


from 1702.05333



Top Threshold/Continuum at lepton colliders


- LC top threshold scan best-known method to measure top quark mass, $\Delta M \sim 30\text{-}70$ MeV
- LC continuum top production best-known method to measure top couplings

Heavy quark production at lepton colliders, qualitatively:




error source	Δm_t^{PS} [MeV]
stat. error (200 fb ⁻¹)	13
theory (NNNLO scale variations, PS scheme)	40
parametric (α_s , current WA)	35
non-resonant contributions (such as single top)	< 40
residual background / selection efficiency	10 – 20
luminosity spectrum uncertainty	< 10
beam energy uncertainty	< 17
combined theory & parametric	30 – 50
combined experimental & backgrounds	25 – 50
total (stat. + syst.)	40 – 75

Threshold region: top velocity $v \sim \alpha_s \ll 1$ non-relativistic EFT: (v)NRQCD from 1702.05333




Continuum region: “standard” fixed-order QCD



Top Threshold/Continuum at lepton colliders


- LC top threshold scan best-known method to measure top quark mass, $\Delta M \sim 30\text{-}70 \text{ MeV}$
- LC continuum top production best-known method to measure top couplings

Heavy quark production at lepton colliders, qualitatively:




error source	Δm_t^{PS} [MeV]
stat. error (200 fb ⁻¹)	13
theory (NNNLO scale variations, PS scheme)	40
parametric (α_s , current WA)	35
non-resonant contributions (such as single top)	< 40
residual background / selection efficiency	10 – 20
luminosity spectrum uncertainty	< 10
beam energy uncertainty	< 17
combined theory & parametric	30 – 50
combined experimental & backgrounds	25 – 50
total (stat. + syst.)	40 – 75

Threshold region: top velocity $v \sim \alpha_s \ll 1$ non-relativistic EFT: (v)NRQCD from 1702.05333



Continuum region: “standard” fixed-order QCD

MATCH





WHIZARD: our NLO MC framework

WHIZARD v2.6.3 (10.02.2018)

<http://whizard.hepforge.org>

<whizard@desy.de>

WHIZARD Team: *Wolfgang Kilian, Thorsten Ohl, JRR*

*Simon Braß/Vincent Rothe/Christian Schwinn/Marco Sekulla/So Young Shim/Florian Staub/Pascal Stienemeier/
Zhijie Zhao + 2 Master*

PUBLICATIONS

General WHIZARD reference: EPJ C71 (2011) 1742, arXiv:0708.4241

0' Mega (ME generator): LC-TOOL (2001) 040; arXiv:hep-ph/0102195

VAMP (MC integrator): CPC 120 (1999) 13; arXiv:hep-ph/9806432

CIRCE (beamstrahlung): CPC 101 (1997) 269; arXiv:hep-ph/9607454

Parton shower: JHEP 1204 (2012) 013; arXiv:1112.1039

Color flow formalism: JHEP 1210 (2012) 022; arXiv:1206.3700

NLO capabilities: JHEP 1612 (2016) 075; arXiv:1609.03390

Parallelization of MEs: CPC 196 (2015) 58; arXiv:1411.3834

POWHEG matching: EPS-HEP (2015) 317; arXiv:1510.02739

```
alpha_power = 2
alphas_power = 0
```

```
process eett = e1,E1 => t, tbar
{ nlo_calculation = "full" }
```

FKS subtraction [Frixione/Kunszt/Signer, '95]

Resonance-aware treatment
[Ježo/Nason, 1509.09071]


NLO QCD final validation phase

NLO EW work in progress

Working NLO interfaces to:

- ★ GoSam [N. Greiner, G. Heinrich... et al.]
- ★ **OpenLoops** [J. Lindert, P. Maierhöfer, S. Pozzorini et al.]
- ★ Recola [A. Denner, L. Hofer, J.-N. Lang, S. Uccirati]








$e^+e^- \rightarrow tt$ off-shell in the continuum

● Cross checks for $2 \rightarrow 2$ and $2 \rightarrow 4$ with Sherpa & Munich

● Using massive b quarks:
no cuts necessary for $e^+e^- \rightarrow W^+W^-bb$

● Full process $e^+e^- \rightarrow \mu^+\nu_\mu e^-\nu_e bb$ exhibits
Coulomb singularity:







$e^+e^- \rightarrow tt$ off-shell in the continuum

● Cross checks for $2 \rightarrow 2$ and $2 \rightarrow 4$ with Sherpa & Munich

● Using massive b quarks:
no cuts necessary for $e^+e^- \rightarrow W^+W^-bb$

● Full process $e^+e^- \rightarrow \mu^+\nu_\mu e^-\nu_e bb$ exhibits
Coulomb singularity:



INPUT PARAMETERS:

$$m_Z = 91.1876 \text{ GeV},$$
$$m_b = 4.2 \text{ GeV},$$

$$m_W = 80.385 \text{ GeV}$$
$$m_t = 173.2 \text{ GeV}.$$

$$\Gamma_{t \rightarrow Wb}^{\text{LO}} = 1.4986 \text{ GeV},$$
$$\Gamma_{t \rightarrow f\bar{f}b}^{\text{LO}} = 1.4757 \text{ GeV},$$

$$\Gamma_{t \rightarrow Wb}^{\text{NLO}} = 1.3681 \text{ GeV},$$
$$\Gamma_{t \rightarrow f\bar{f}b}^{\text{NLO}} = 1.3475 \text{ GeV}.$$

$$m_H = 125 \text{ GeV}$$

$$\Gamma_H = 0.00431 \text{ GeV}$$

$$\Gamma_Z^{\text{LO}} = 2.4409 \text{ GeV},$$
$$\Gamma_W^{\text{LO}} = 2.0454 \text{ GeV},$$


$$\Gamma_Z^{\text{NLO}} = 2.5060 \text{ GeV},$$
$$\Gamma_W^{\text{NLO}} = 2.0978 \text{ GeV}.$$

Complex Mass Scheme (CMS):

$$\mu_i^2 = M_i^2 - i\Gamma_i M_i \quad \text{for } i = W, Z, t, H$$

$$s_w^2 = 1 - c_w^2 = 1 - \frac{\mu_W^2}{\mu_Z^2}$$







$e^+e^- \rightarrow tt$ off-shell in the continuum

● Cross checks for $2 \rightarrow 2$ and $2 \rightarrow 4$ with Sherpa & Munich

● Using massive b quarks:
no cuts necessary for $e^+e^- \rightarrow W^+W^-bb$

● Full process $e^+e^- \rightarrow \mu^+\nu_\mu e^-\nu_e bb$ exhibits
Coulomb singularity:



INPUT PARAMETERS:

$$m_Z = 91.1876 \text{ GeV},$$
$$m_b = 4.2 \text{ GeV},$$

$$m_W = 80.385 \text{ GeV}$$
$$m_t = 173.2 \text{ GeV}.$$

$$\Gamma_{t \rightarrow Wb}^{\text{LO}} = 1.4986 \text{ GeV},$$
$$\Gamma_{t \rightarrow f\bar{f}b}^{\text{LO}} = 1.4757 \text{ GeV},$$

$$\Gamma_{t \rightarrow Wb}^{\text{NLO}} = 1.3681 \text{ GeV},$$
$$\Gamma_{t \rightarrow f\bar{f}b}^{\text{NLO}} = 1.3475 \text{ GeV}.$$

$$m_H = 125 \text{ GeV}$$

$$\Gamma_H = 0.00431 \text{ GeV}$$

$$\Gamma_Z^{\text{LO}} = 2.4409 \text{ GeV},$$
$$\Gamma_W^{\text{LO}} = 2.0454 \text{ GeV},$$


$$\Gamma_Z^{\text{NLO}} = 2.5060 \text{ GeV},$$
$$\Gamma_W^{\text{NLO}} = 2.0978 \text{ GeV}.$$


Complex Mass Scheme (CMS):

$$\mu_i^2 = M_i^2 - i\Gamma_i M_i \quad \text{for } i = W, Z, t, H$$

$$s_w^2 = 1 - c_w^2 = 1 - \frac{\mu_W^2}{\mu_Z^2}$$


● Typical pentagons:






NLO QCD Results for off-shell $e^+e^- \rightarrow t\bar{t}$

$e^+e^- \rightarrow t\bar{t}$ and $e^+e^- \rightarrow W^+W^-b\bar{b}$




$e^+e^- \rightarrow t\bar{t}$ and $e^+e^- \rightarrow W^+W^-b\bar{b}$ at $\sqrt{s} = 800$ GeV





\sqrt{s} [GeV]	$e^+e^- \rightarrow t\bar{t}$			$e^+e^- \rightarrow W^+W^-b\bar{b}$		
	σ^{LO} [fb]	σ^{NLO} [fb]	K-factor	σ^{LO} [fb]	σ^{NLO} [fb]	K-factor
500	548.4	$627.4^{+1.4\%}_{-0.9\%}$	1.14	600.7	$675.1^{+0.4\%}_{-0.8\%}$	1.12
800	253.1	$270.9^{+0.8\%}_{-0.4\%}$	1.07	310.2	$320.7^{+1.1\%}_{-0.7\%}$	1.03
1000	166.4	$175.9^{+0.7\%}_{-0.3\%}$	1.06	217.2	$221.6^{+1.1\%}_{-1.0\%}$	1.02
1400	86.62	$90.66^{+0.6\%}_{-0.2\%}$	1.05	126.4	$127.9^{+0.7\%}_{-1.5\%}$	1.01
3000	19.14	$19.87^{+0.5\%}_{-0.2\%}$	1.04	37.89	$37.63^{+0.4\%}_{-0.9\%}$	0.993

Chokouf /Kilian/Lindert/Pozzorini/JRR/Weiss, 1609.03390






Differential Results for off-shell $e^+e^- \rightarrow tt$



$$m_t^2 = m_W^2 + \frac{2\langle m_{\ell j_b}^2 \rangle}{1 - \langle \cos \theta_{\ell j_b} \rangle}$$



Chokouf/Kilian/Lindert/Pozzorini/JRR/Weiss, 1609.03390






- NRQCD is EFT for non-relativistic quark-antiquark systems: separate $M \cdot v$ and $M \cdot v^2$
- Integrate out hard quark and gluon d.o.f.
- Resummation of singular terms close to threshold ($v = 0$) Hoang et al. '99-'01; Beneke et al., '13-'14

Phase space of two massive particles

$$R \equiv \frac{\sigma_{t\bar{t}}}{\sigma_{\mu\mu}} = v \sum_k \left(\frac{\alpha_s}{v}\right)^k \sum_i (\alpha_s \ln v)^i \times$$

$$\times \{1 (\mathbf{LL}); \alpha_s, v (\mathbf{NLL}); \alpha_s^2, \alpha_s v, v^2 (\mathbf{NNLL})\}$$

(p/v)NRQCD EFT w/ RG improvement



- NRQCD is EFT for non-relativistic quark-antiquark systems: separate $M \cdot v$ and $M \cdot v^2$
- Integrate out hard quark and gluon d.o.f.
- Resummation of singular terms close to threshold ($v = 0$) Hoang et al. '99-'01; Beneke et al., '13-'14


Phase space of two massive particles

$$R \equiv \frac{\sigma_{t\bar{t}}}{\sigma_{\mu\mu}} = v \sum_k \left(\frac{\alpha_s}{v}\right)^k \sum_i (\alpha_s \ln v)^i \times \{1 (\mathbf{LL}); \alpha_s, v (\mathbf{NLL}); \alpha_s^2, \alpha_s v, v^2 (\mathbf{NNLL})\}$$

(p/v)NRQCD EFT w/ RG improvement

$$R^{\gamma,Z}(s) = \underbrace{F^v(s)R^v(s)}_{\text{s-wave: LL+NLL}} + \underbrace{F^a(s)R^a(s)}_{\text{p-wave} \sim v^2: \text{NNLL}}$$

but contributes
at NLL differentially!



Top Threshold Resummation in (p)NRQCD

- NRQCD is EFT for non-relativistic quark-antiquark systems: separate $M \cdot v$ and $M \cdot v^2$
- Integrate out hard quark and gluon d.o.f.
- Resummation of singular terms close to threshold ($v = 0$) Hoang et al. '99-'01; Beneke et al., '13-'14

Phase space of two massive particles


$$R \equiv \frac{\sigma_{t\bar{t}}}{\sigma_{\mu\mu}} = v \sum_k \left(\frac{\alpha_s}{v}\right)^k \sum_i (\alpha_s \ln v)^i \times \{1 (\mathbf{LL}); \alpha_s, v (\mathbf{NLL}); \alpha_s^2, \alpha_s v, v^2 (\mathbf{NNLL})\}$$


(p/v)NRQCD EFT w/ RG improvement

$$R^{\gamma,Z}(s) = \underbrace{F^v(s)R^v(s)}_{\text{s-wave: LL+NLL}} + \underbrace{F^a(s)R^a(s)}_{\text{p-wave} \sim v^2: \text{NNLL}}$$

but contributes at NLL differentially!

Coulomb potential gluon ladder resummation





Top Threshold Resummation in (p)NRQCD

- NRQCD is EFT for non-relativistic quark-antiquark systems: separate $M \cdot v$ and $M \cdot v^2$
- Integrate out hard quark and gluon d.o.f.
- Resummation of singular terms close to threshold ($v = 0$) Hoang et al. '99-'01; Beneke et al., '13-'14


Phase space of two massive particles

$$R \equiv \frac{\sigma_{t\bar{t}}}{\sigma_{\mu\mu}} = v \sum_k \left(\frac{\alpha_s}{v}\right)^k \sum_i (\alpha_s \ln v)^i \times \underbrace{\{1 (\mathbf{LL}); \alpha_s, v (\mathbf{NLL}); \alpha_s^2, \alpha_s v, v^2 (\mathbf{NNLL})\}}_{\text{(p/v)NRQCD EFT w/ RG improvement}}$$


$$R^{\gamma,Z}(s) = \underbrace{F^v(s)R^v(s)}_{\text{s-wave: LL+NLL}} + \underbrace{F^a(s)R^a(s)}_{\text{p-wave} \sim v^2: \mathbf{NNLL}}$$

but contributes at NLL differentially!

Coulomb potential gluon ladder resummation




can be mapped onto effective $t\bar{t}V$ vertex




$$\mathbb{C} \ni \mathcal{G}_{(N)LL}^{v/a} = \mathcal{G}_{(N)LL}^{v/a}(\alpha_s, M_t^{\text{pole}}, \sqrt{s}, |\vec{p}_t|, \Gamma_t)$$

differential in off-shell $t\bar{t}$ phase space



Top Threshold in WHIZARD

- Implement resummed threshold effects as effective vertex [form factor] in WHIZARD
- $G^{v,a}(0, p_t, E + i\Gamma_t, \nu)$ from TOPPIK code [Jezabek/Teubner], included in WHIZARD



- Default parameters:

$$M^{1S} = 172 \text{ GeV}, \quad \Gamma_t = 1.54 \text{ GeV},$$



$$\alpha_s(M_Z) = 0.118$$


$$M^{1S} = M_t^{pole} (1 - \Delta_{(Coul.)}^{LL/NLL})$$

- ▶ Important effects: beamstrahlung; ISR; LO EW terms
- ▶ Exclusive observables accessible

Theory uncertainties from scale variations: hard and soft scale

$$\mu_h = h \cdot m_t \quad \mu_s = f \cdot m_t v$$






Top threshold: validation and matching

- Transition region between relativistic and resummation effects

$$\begin{aligned}
 \sigma_{\text{NLO+NLL}} = & \sigma_{\text{NLO}} + \left(\tilde{F}_{\text{NLL}} - \tilde{F}_{\text{NLL}}^{\text{exp}} \right) \left(\begin{array}{c} e^+ \\ \\ \\ e^- \end{array} \rightarrow \begin{array}{c} b \\ W^+ \\ W^- \\ \bar{b} \end{array} \right) \left(\begin{array}{c} b \\ W^+ \\ W^- \\ \bar{b} \end{array} \rightarrow \begin{array}{c} e^+ \\ \\ \\ e^- \end{array} \right) \\
 & + \left| \tilde{F}_{\text{NLL}} \left(\begin{array}{c} e^+ \\ \\ \\ e^- \end{array} \rightarrow \begin{array}{c} b \\ W^+ \\ W^- \\ \bar{b} \end{array} \right) \right|^2 \\
 & + \left\{ \tilde{F}_{\text{NLL}} \left(\begin{array}{c} e^+ \\ \\ \\ e^- \end{array} \rightarrow \begin{array}{c} b \\ W^+ \\ W^- \\ \bar{b} \end{array} \right) \left(\begin{array}{c} e^+ \\ \\ \\ e^- \end{array} \rightarrow \begin{array}{c} b \\ W^+ \\ W^- \\ \bar{b} \end{array} \right) + \left(\begin{array}{c} e^+ \\ \\ \\ e^- \end{array} \rightarrow \begin{array}{c} b \\ W^+ \\ W^- \\ \bar{b} \end{array} \right) \tilde{F}_{\text{NLL}} \right\} \\
 & + \left| \tilde{F}_{\text{NLL}} \left(\begin{array}{c} e^+ \\ \\ \\ e^- \end{array} \rightarrow \begin{array}{c} b \\ W^+ \\ W^- \\ \bar{b} \end{array} \right) \right|^2 + \left| \tilde{F}_{\text{NLL}} \left(\begin{array}{c} e^+ \\ \\ \\ e^- \end{array} \rightarrow \begin{array}{c} b \\ W^+ \\ W^- \\ \bar{b} \end{array} \right) \right|^2,
 \end{aligned}$$





Smoothstep matching function:

$$\begin{aligned}
 \sigma_{\text{matched}} = & \sigma_{\text{FO}} [\alpha_{\text{H}}] + \sigma_{\text{NRQCD}}^{\text{full}} [f_s \alpha_{\text{H}}, f_s \alpha_{\text{S}}, f_s \alpha_{\text{US}}] \\
 & - \sigma_{\text{NRQCD}}^{\text{expanded}} [f_s \alpha_{\text{H}}, f_s \alpha_{\text{H}}],
 \end{aligned}$$

$$\begin{aligned}
 \alpha_{\text{H}} &= \alpha_s [\mu_{\text{H}} = hM_t^{1\text{S}}], & \alpha_{\text{F}} &= \alpha_s [\mu_{\text{F}} = hM_t^{1\text{S}} \sqrt{\nu_*}], \\
 \alpha_{\text{S}} &= \alpha_s [\mu_{\text{S}} = hM_t^{1\text{S}} f\nu_*], & \alpha_{\text{U}} &= \alpha_s [\mu_{\text{US}} = hM_t^{1\text{S}} (f\nu_*)^2]
 \end{aligned}$$


$$f_s(v) = \begin{cases} 1 & v < v_1 \\ 1 - 3 \left(\frac{v-v_1}{v_2-v_1} \right)^2 - 2 \left(\frac{v-v_1}{v_2-v_1} \right)^3 & v_1 \leq v \leq v_2 \\ 0 & v > v_2 \end{cases}$$







Top threshold: validation and matching

NLO predictions for on- and off-shell $t\bar{t}$ production




$\Delta_{m_t} = 30$ GeV, expanded, evaluated with α_H , only s-wave contributions







Top threshold: validation and matching

NLO predictions for on- and off-shell $t\bar{t}$ production




$\Delta_{m_t} = 30$ GeV, LL, only s-wave contributions







Top threshold: validation and matching

NLO predictions for on- and off-shell $t\bar{t}$ production





$\Delta_{m_t} = 30$ GeV, NLL, only s-wave contributions





Matching threshold NLL to continuum NLO




Total uncertainty: h - f variation band and matching [switch-off function]

Symmetrization of error bands:



$$\sigma_{\max} = \max \left[\max_{i \in \text{HF}} \sigma_i, \sigma_0 + (\sigma_0 - \min_{i \in \text{HF}} \sigma_i) \right]$$


$$\sigma_{\min} = \min \left[\min_{i \in \text{HF}} \sigma_i, \sigma_0 - (\max_{i \in \text{HF}} \sigma_i - \sigma_0) \right]$$











Threshold matching with QED ISR









Matched threshold differential distributions







Matched threshold differential distributions



DISTRIBUTIONS DEPEND ON M(TOP) NOT YET FULLY EXPLOITED






Summary & Outlook

Next steps: higher QCD order, EW corrections (ISR matching!!), soft gluons

$$e^+e^- \rightarrow W^+bW^-\bar{b}$$






- Top physics is cornerstone of future lepton collider program
- **Leptonic top fully off-shell at NLO QCD**
- Inclusive processes: off-shell background grows with energy

- **Complete v NRQCD threshold / QCD-NLO continuum matching**
- **Offers framework for new differential top mass measurements**
- Can in principle be reweighted to NNNLO QCD accuracy at threshold
- Next steps: EW corrections, semi-leptonic/hadronic top decays, ttH threshold matching, top threshold matched with EW corrections



- WHIZARD 2.6 framework for automated (QCD) NLO
- **NLO QCD** (almost) done → WHIZARD 3.0 [EW in validation]





Outlook to PDG 203x:







BACKUP





- Paradigm processes at lepton colliders: **precision determination of top properties**
- Major background for EW measurements (VVV and VBS); any [most] BSM searches

On-Shell process: $e^+e^- \rightarrow t\bar{t}$


- NLO QCD [Jersak/Laermann/Zerwas, 1982]
- NNLO QCD [Chetyrkin/Kühn/Steinhauser, 1996; Harlander/Steinhauser, 1998]
- NLO EW [Beenakker/von der Marck/Hollik, 1991; Beenakker/Denner/Kraft, 1993]
- Threshold enhancement [Fadin/Khoze, 1987; Strassler/Peskin, 1991; Jezabek/Kühn/Teubner, 1992; Sumino et al., 1992]


Top width: $t \rightarrow W^+b$

- NLO QCD [Jezabek/Kühn, 1989]
- NNLO QCD [Guo/Li/Zhu, 2012]

Off-Shell process: $e^+e^- \rightarrow W^+\bar{b}W^-b$

- NLO QCD [Guo/Ma/Wang/Zhang, 2008] ✗
- NLO QCD diff. [Chokoufe/JRR/Weiss, 2015; Liebler/Moortgat-Pick/Papanastasiou, 2015; Chokoufe/Kilian/Lindert/JRR/Pozzorini/Weiss, 2016]






Working NLO interfaces to:

- ★ GoSam [N. Greiner, G. Heinrich, J. v. Soden-Fraunhofen et al.]
- ★ OpenLoops [F. Cascioli, J. Lindert, P. Maierhöfer, S. Pozzorini]
- ★ RecoLa [A. Denner, L. Hofer, J.-N. Lang, S. Uccirati]


```
alpha_power = 2
alphas_power = 0


process eett = e1,E1 => t, tbar
{ nlo_calculation = "full" }
```



- 🔊 FKS subtraction [Frixione/Kunszt/Signer, 1995]
- 🔊 Resonance-aware treatment [Ježo/Nason, 1509.09071]
- 🔊 Virtual MEs external
- 🔊 Real and virtual subtraction terms internal
- 🔊 NLO decays available for the NLO processes
- 🔊 Fixed order events for plotting (weighted, either LHEF or HepMC)
- 🔊 Automated POWHEG damping and matching
- 🔊 **NLO QCD (massless & massive emitters) fully supported**
- 🔊 **Status of EW corrections: all parts**

technically completed, validation phase started [Rothe et al.]






Working NLO interfaces to:

- ★ GoSam [N. Greiner, G. Heinrich, J. v. Soden-Fraunhofen et al.]
- ★ OpenLoops [F. Cascioli, J. Lindert, P. Maierhöfer, S. Pozzorini]
- ★ RecoLa [A. Denner, L. Hofer, J.-N. Lang, S. Uccirati]


```
alpha_power = 2
alphas_power = 0

process eett = e1,E1 => t, tbar
{ nlo_calculation = "full" }
```



- 🔊 FKS subtraction [Frixione/Kunszt/Signer, 1995]
- 🔊 Resonance-aware treatment [Ježo/Nason, 1509.09071]
- 🔊 Virtual MEs external
- 🔊 Real and virtual subtraction terms internal
- 🔊 NLO decays available for the NLO processes
- 🔊 Fixed order events for plotting (weighted, either LHEF or HepMC)
- 🔊 Automated POWHEG damping and matching
- 🔊 **NLO QCD (massless & massive emitters) fully supported**
- 🔊 **Status of EW corrections: all parts**

technically completed, validation phase started [Rothe et al.]






- Amplitudes (except for pure QCD/QED) contain **resonances** (Z, W, H, t)
- In general: resonance masses *not* respected by modified kinematics of subtraction terms**
- Algorithm to include resonance histories**
[Ježo/Nason, 1509.09071]
- Most important for narrow resonances ($H \rightarrow bb$)
- Additional soft mismatch integration component

$$\begin{aligned} \blacktriangleright D_H^{\text{Born}} &= \left[(\bar{p}_{bb}^2 - m_H^2)^2 + m_H^2 \Gamma_H^2 \right]^{-1}, \\ \blacktriangleright D_H^{\text{Real}} &= \left[(p_{bbg}^2 - m_H^2)^2 + m_H^2 \Gamma_H^2 \right]^{-1} \end{aligned}$$

$$p_{bbg}^2 = \bar{p}_{bb}^2 + \Delta_{bbg}^2$$

$$\frac{D_H^{\text{Born}}}{D_H^{\text{Real}}} \xrightarrow{\bar{p}_{bb}^2 \rightarrow m_H^2} 1 + \frac{\Delta_{bbg}^4}{m_H^2 \Gamma_H^2}$$






- Amplitudes (except for pure QCD/QED) contain **resonances** (Z, W, H, t)
- In general: resonance masses *not* respected by modified kinematics of subtraction terms**
- Algorithm to include resonance histories**
[Ježo/Nason, 1509.09071]
- Most important for narrow resonances ($H \rightarrow bb$)
- Additional soft mismatch integration component

$$\begin{aligned} \blacktriangleright D_H^{\text{Born}} &= \left[(\bar{p}_{bb}^2 - m_H^2)^2 + m_H^2 \Gamma_H^2 \right]^{-1}, \\ \blacktriangleright D_H^{\text{Real}} &= \left[(p_{bbg}^2 - m_H^2)^2 + m_H^2 \Gamma_H^2 \right]^{-1} \end{aligned}$$

$$p_{bbg}^2 = \bar{p}_{bb}^2 + \Delta_{bbg}^2$$

$$\frac{D_H^{\text{Born}}}{D_H^{\text{Real}}} \stackrel{\bar{p}_{bb}^2 \rightarrow m_H^2}{=} 1 + \frac{\Delta_{bbg}^4}{m_H^2 \Gamma_H^2}$$





Resonance mappings for NLO processes


- Amplitudes (except for pure QCD/QED) contain resonances (Z, W, H, t)
- In general: resonance masses *not* respected by modified kinematics of subtraction terms

Algorithm to include resonance histories
 [Ježo/Nason, 1509.09071]

- Most important for narrow resonances ($H \rightarrow bb$)
- Additional soft mismatch integration component

$$\begin{aligned} \blacktriangleright D_H^{\text{Born}} &= \left[(\bar{p}_{bb}^2 - m_H^2)^2 + m_H^2 \Gamma_H^2 \right]^{-1}, \\ \blacktriangleright D_H^{\text{Real}} &= \left[(p_{bbg}^2 - m_H^2)^2 + m_H^2 \Gamma_H^2 \right]^{-1} \end{aligned}$$


$$p_{bbg}^2 = \bar{p}_{bb}^2 + \Delta_{bbg}^2 \quad \frac{D_H^{\text{Born}}}{D_H^{\text{Real}}} \stackrel{\bar{p}_{bb}^2 \rightarrow m_H^2}{=} 1 + \frac{\Delta_{bbg}^4}{m_H^2 \Gamma_H^2}$$



WHIZARD complete automatic implementation: example $e^+ e^- \rightarrow \mu\mu bb$ (ZZ, ZH histories)

It	Calls	Integral[fb]	Error[fb]	Err[%]	Acc	Eff[%]	Chi2	N[It]
1	11988	9.6811847E+00	6.42E+00	66.30	72.60*	0.65		
2	11959	2.8539703E+00	2.35E-01	8.25	9.02*	0.69		
3	11936	2.4907574E+00	6.54E-01	26.25	28.68	0.35		
4	11908	2.7695559E+00	9.67E-01	34.91	38.09	0.30		
5	11874	2.4346151E+00	4.82E-01	19.80	21.57*	0.74		
5	59665	2.7539078E+00	1.97E-01	7.15	17.47	0.74	0.49	5

standard FKS





Resonance mappings for NLO processes

- Amplitudes (except for pure QCD/QED) contain resonances (Z, W, H, t)
- In general: resonance masses *not* respected by modified kinematics of subtraction terms

Algorithm to include resonance histories


[Ježo/Nason, 1509.09071]

- Most important for narrow resonances ($H \rightarrow bb$)

- Additional soft mismatch integration component

$$\begin{aligned} \blacktriangleright D_H^{\text{Born}} &= \left[(\bar{p}_{bb}^2 - m_H^2)^2 + m_H^2 \Gamma_H^2 \right]^{-1}, \\ \blacktriangleright D_H^{\text{Real}} &= \left[(p_{bbg}^2 - m_H^2)^2 + m_H^2 \Gamma_H^2 \right]^{-1} \end{aligned}$$

$$p_{bbg}^2 = \bar{p}_{bb}^2 + \Delta_{bbg}^2 \quad \frac{D_H^{\text{Born}}}{D_H^{\text{Real}}} \stackrel{\bar{p}_{bb}^2 \rightarrow m_H^2}{=} 1 + \frac{\Delta_{bbg}^4}{m_H^2 \Gamma_H^2}$$



- WHIZARD complete automatic implementation: example $e^+ e^- \rightarrow \mu\mu bb$ (ZZ, ZH histories)

It	Calls	Integral[fb]	Error[fb]	Err[%]	Acc	Eff[%]	Chi2	N[It]
1	11988	9.6811847E+00	6.42E+00	66.30	72.60*	0.65		
2	11959	2.8539703E+00	2.35E-01	8.25	9.02*	0.69		
3	11936	2.4907574E+00	6.54E-01	26.25	28.68	0.35		
4	11908	2.7695559E+00	9.67E-01	34.91	38.09	0.30		
5	11874	2.4346151E+00	4.82E-01	19.80	21.57*	0.74		
5	59665	2.7539078E+00	1.97E-01	7.15	17.47	0.74	0.49	5

standard FKS

It	Calls	Integral[fb]	Error[fb]	Err[%]	Acc	Eff[%]	Chi2	N[It]
1	11988	2.9057032E+00	8.35E-02	2.87	3.15*	7.90		
2	11962	2.8591952E+00	5.20E-02	1.82	1.99*	10.91		
3	11936	2.9277880E+00	4.09E-02	1.40	1.52*	14.48		
4	11902	2.8512337E+00	3.98E-02	1.40	1.52*	13.70		
5	11874	2.8855399E+00	3.87E-02	1.34	1.46*	17.15		
5	59662	2.8842006E+00	2.04E-02	0.71	1.72	17.15	0.53	5

FKS with resonance mappings





Top-Forward Backward Asymmetry

$$A_{FB} = \frac{\sigma(\cos \theta_t > 0) - \sigma(\cos \theta_t < 0)}{\sigma(\cos \theta_t > 0) + \sigma(\cos \theta_t < 0)}$$


Gluon emission symmetric in $\theta \Rightarrow$
NLO QCD corrections small

A_{FB} of the top quark



	$e^+e^- \rightarrow$	A_{FB}^{LO}	A_{FB}^{NLO}	$A_{FB}^{NLO} / A_{FB}^{LO}$
A_{FB}	$t\bar{t}$	-0.535	-0.539	1.013
	$W^+W^-b\bar{b}$	-0.428	-0.426	0.995
	$\mu^+e^-\nu_\mu\bar{\nu}_e b\bar{b}$	-0.415	-0.409	0.986
	$\mu^+e^-\nu_\mu\bar{\nu}_e b\bar{b}$, without neutrinos	-0.402	-0.387	0.964
\bar{A}_{FB}	$t\bar{t}$	0.535	0.539	1.013
	$W^+W^-b\bar{b}$	0.428	0.426	0.995
	$\mu^+e^-\nu_\mu\bar{\nu}_e b\bar{b}$	0.415	0.409	0.986
	$\mu^+e^-\nu_\mu\bar{\nu}_e b\bar{b}$, without neutrinos	0.377	0.350	0.928



- Precise predictions of multi-parton final states require properly matched samples
- NLO QCD including POWHEG matching already available [WHIZARD+OpenLoops]
- All descriptions at NLO at the moment for the on-shell process
- Even LO simulations are demanding, e.g.: $e^+e^- \rightarrow b\bar{b}b\bar{b}jjl\nu_e, \quad b\bar{b}jjjjjjl\nu_e$



- Precise predictions of multi-parton final states require properly matched samples
- NLO QCD including POWHEG matching already available [WHIZARD+OpenLoops]
- All descriptions at NLO at the moment for the on-shell process
- Even LO simulations are demanding, e.g.: $e^+e^- \rightarrow b\bar{b}b\bar{b}jjl\nu_e$, $b\bar{b}jjjjjjl\nu_e$



[Chokoufe/JRR/Weiss]