

Looking Beyond the Standard Model with Extremely Energetic Cosmic Particles

Andreas Ringwald

<http://www.desy.de/~ringwald>




XVIII Workshop *Beyond the Standard Model*
March 13–16, 2006, Bad Honnef, D

1. Introduction

1

- There is a high energy universe:
Gamma rays have been identified up to energies $E \lesssim \text{few} \times 10^3$ GeV




■ = Pulsar/Plerion □ = SNR ★ = Starburst galaxy ○ = OB association
◆ = AGN (BL Lac) ▲ = Radio galaxy ✕ = XRB ● = Undetermined

[M. Martinez '05]

1. Introduction

2

- There is a high energy universe:
Gamma rays have been identified up to energies $E \lesssim \text{few} \times 10^3 \text{ GeV}$
Cosmic rays have been observed up to energies $E \lesssim \text{few} \times 10^{11} \text{ GeV}$



[T. K. Gaisser '05]

1. Introduction

3

- There is a high energy universe:

Gamma rays have been identified up to energies $E \lesssim \text{few} \times 10^3 \text{ GeV}$

Cosmic rays have been observed up to energies $E \lesssim \text{few} \times 10^{11} \text{ GeV}$

- It is under active observation:

Gamma ray observatories: e.g.
H.E.S.S., MAGIC

Air shower detectors: e.g. **Pierre Auger Observatory**




[www.auger.org]

1. Introduction

4

- **There is a high energy universe:**
Gamma rays have been identified up to energies $E \lesssim \text{few} \times 10^3 \text{ GeV}$
Cosmic rays have been observed up to energies $E \lesssim \text{few} \times 10^{11} \text{ GeV}$
- **It is under active observation:**
Gamma ray observatories: e.g.
H.E.S.S., MAGIC
Air shower detectors: e.g. **Pierre Auger Observatory**
Neutrino telescopes: e.g. **IceCube**
- **Attack fundamental questions:**
What is it made of? What are the cosmic accelerators? Can we exploit them also for particle physics?




[icecube.wisc.edu]

Outline:

- 2. Observations at ultrahigh energies**
- 3. Non-observations at ultrahigh energies**
- 4. The future ...**
- 5. Conclusions**

2. Observations at ultrahigh energies

- **Spectrum:** Large statistical and systematic uncertainties
 - ↑ low flux
 - ↑ energy from shower simulations



[Ahlers *et al.* '05]

2. Observations at ultrahigh energies


- **Spectrum:** Large statistical and systematic uncertainties
 - ⇐ low flux
 - ⇐ energy from shower simulations
- Crucial improvement by **PAO:**
 - ⇐ huge size ⇒ better statistics
 - ⇐ hybrid observations ⇒ better energy calibration through Fly's Eye technique, direction from ground array



[www.auger.org]


2. Observations at ultrahigh energies

- **Spectrum:** Large statistical and systematic uncertainties
 - ⇒ low flux
 - ⇒ energy from shower simulations
- Crucial improvement by **PAO**:
 - ⇒ huge size ⇒ better statistics
 - ⇒ hybrid observations ⇒ better energy calibration through Fly's Eye technique, direction from ground array
- It works

[Ahlers *et al.* '05]

2. Observations at ultrahigh energies


- **Spectrum:** Large statistical and systematic uncertainties
 - ⇒ low flux
 - ⇒ energy from shower simulations
- Crucial improvement by **PAO**:
 - ⇒ huge size ⇒ better statistics
 - ⇒ hybrid observations ⇒ better energy calibration through Fly's Eye technique, direction from ground array
- It works

[Ahlers *et al.* '05]

2. Observations at ultrahigh energies

10


- Angular distribution: \approx isotrop



2. Observations at ultrahigh energies

11

- **Angular distribution:** \approx isotropic
- **Composition:** Large uncertainty
 \Leftarrow studies rely on simulations
- Cosmic rays above $\gtrsim 10^{8.6}$ GeV dominantly protons




[Heck '05]

2. Observations at ultrahigh energies

12

- **Angular distribution:** \approx isotrop
- **Composition:** Large uncertainty
 \Leftarrow studies rely on simulations
- Cosmic rays above $\gtrsim 10^{8.6}$ eV, the “second knee”, dominantly protons
- Assume that CR's in $10^{[8.6,11]}$ GeV range originate from isotropically distributed extragalactic proton sources, with simple power-law injection spectra $\propto E_i^{-\gamma}(1+z)^n$

[Berezinsky,..'02-'05;...;Ahlers *et al.* '05]



Beyond the Standard Model, Bad Honnef, D

2. Observations at ultrahigh energies


13

- **Angular distribution:** \approx isotrop
 - **Composition:** Large uncertainty
⇐ studies rely on simulations
 - Cosmic rays above $\gtrsim 10^{8.6}$ eV, the “second knee”, dominantly protons
 - Assume that CR's in $10^{[8.6,11]}$ GeV range originate from isotropically distributed extragalactic proton sources, with simple power-law injection spectra $\propto E_i^{-\gamma}(1+z)^n$
- [Berezinsky,..'02-'05;...;Ahlers et al. '05]

\Rightarrow Good fit; inelastic interactions with **CMB** (e^+e^- “dip”; π “bump”) visible; some **post-GZK events?**


A. Ringwald

[Greisen;Zatsepin,Kuzmin '67]




[Ahlers et al. '05]
Beyond the Standard Model, Bad Honnef, D

- Possible sources of these protons:
GRB, AGN, . . .




- Possible sources of these protons:
GRB, **AGN**, . . .
- **Shock acceleration:**
 - p 's, confined by magnetic fields, accelerate through repeated scattering by plasma shock fronts




[Ahlers PhD in prep.]

- Possible sources of these protons:
GRB, **AGN**, . . .
- **Shock acceleration:**
 - p 's, confined by magnetic fields, accelerate through repeated scattering by plasma shock fronts




- Possible sources of these protons:
GRB, **AGN**, . . .
- **Shock acceleration:**
 - p 's, confined by magnetic fields, accelerate through repeated scattering by plasma shock fronts
 - production of π 's and n 's through collisions of the trapped p 's with ambient plasma produces γ 's, ν 's and CR's (n diffusion from source)



[Ahlers PhD in prep.]


- Possible sources of these protons:
GRB, **AGN**, . . .
- **Shock acceleration:**
 - p 's, confined by magnetic fields, accelerate through repeated scattering by plasma shock fronts
 - production of π 's and n 's through collisions of the trapped p 's with ambient plasma produces γ 's, ν 's and CR's (n diffusion from source)
- **Neutrinos as diagnostic tool:**
 - ν 's from sources ($p\gamma \rightarrow n + \pi$'s) close to be measured
 - Cosmogenic neutrino flux (from $p\gamma_{\text{CMB}} \rightarrow N\pi$'s) dominates above 10^9 GeV



[Ahlers *et al.* '05]

3. Non-observations at ultrahigh energies

- $C\nu$'s with $E_\nu \gtrsim 10^8$ GeV probe νN scattering at $\sqrt{s_{\nu N}} \gtrsim 14$ TeV (**LHC**)




[Ahlers *et al.* '05]

3. Non-observations at ultrahigh energies

- $C\nu$'s with $E_\nu \gtrsim 10^8$ GeV probe νN scattering at $\sqrt{s_{\nu N}} \gtrsim 14$ TeV (**LHC**)
- Perturbative Standard Model (**SM**)
 \approx under control (\leftarrow **HERA**)


[Gandhi *et al.* '98; Kwiecinski *et al.* '98; ...]



3. Non-observations at ultrahigh energies

- $C\nu$'s with $E_\nu \gtrsim 10^8$ GeV probe νN scattering at $\sqrt{s_{\nu N}} \gtrsim 14$ TeV (**LHC**)
- Perturbative Standard Model (**SM**)
≈ under control (← **HERA**)

[Gandhi *et al.* '98; Kwiecinski *et al.* '98; ...]
- ⇒ Search for enhancements in $\sigma_{\nu N}$ beyond (perturbative) SM:
 - ◊ Electroweak sphaleron production
($B + L$ violating processes in SM)




[Fodor,Katz,AR,Tu '03; Han,Hooper '03]

3. Non-observations at ultrahigh energies

- $C\nu$'s with $E_\nu \gtrsim 10^8$ GeV probe νN scattering at $\sqrt{s_{\nu N}} \gtrsim 14$ TeV (**LHC**)
- Perturbative Standard Model (**SM**) \approx under control (\leftarrow **HERA**)

[Gandhi *et al.* '98; Kwiecinski *et al.* '98; ...]

- ⇒ Search for enhancements in $\sigma_{\nu N}$ beyond (perturbative) SM:
- ◊ Electroweak sphaleron production ($B + L$ violating processes in SM)
 - ◊ Kaluza-Klein, **black hole**, p -brane or string ball production in TeV scale gravity models
 - ◊ . . .




[AR,Tu '01; Tu '04]

“Model-independent” upper bounds on $\sigma_{\nu N}$

$$\frac{dN}{dt} \propto \int dE_\nu F_\nu(E_\nu) \sigma_{\nu N}(E_\nu)$$

- ⇒ Non-observation of deeply-penetrating particles, together with lower bound on F_ν (e.g. cosmogenic ν 's)
- ⇒ upper bound on $\sigma_{\nu N}$

[Berezinsky,Smirnov '74; Morris,AR '94; Tyler,Olinto,Sigl '01;...]



[Anchordoqui,Fodor,Katz,AR,Tu '04]

“Model-independent” upper bounds on $\sigma_{\nu N}$


$$\frac{dN}{dt} \propto \int dE_\nu F_\nu(E_\nu) \sigma_{\nu N}(E_\nu)$$

- ⇒ Non-observation of deeply-penetrating particles, together with lower bound on F_ν (e.g. cosmogenic ν 's)
⇒ upper bound on $\sigma_{\nu N}$

[Berezinsky,Smirnov '74; Morris,AR '94; Tyler,Olinto,Sigl '01;..]

- Recent quantitative analysis:
[Anchordoqui,Fodor,Katz,AR,Tu '04]
 - ◊ Best current limits from exploitation of **RICE** search results
[Kravchenko *et al.* [RICE] '02,03]

RICE: Radio Ice Cerenkov Experiment



[www2.phys.canterbury.ac.nz/rice]


“Model-independent” upper bounds on $\sigma_{\nu N}$

$$\frac{dN}{dt} \propto \int dE_\nu F_\nu(E_\nu) \sigma_{\nu N}(E_\nu)$$

- ⇒ Non-observation of deeply-penetrating particles, together with lower bound on F_ν (e.g. cosmogenic ν 's)
⇒ upper bound on $\sigma_{\nu N}$

[Berezinsky,Smirnov '74; Morris,AR '94; Tyler,Olinto,Sigl '01;..]

- Recent quantitative analysis:
 - ◊ Best current limits from exploitation of **RICE** search results
[Kravchenko *et al.* [RICE] '02,03]
 - ◊ **Auger** will improve these limits by one order of magnitude



[Anchordoqui,Fodor,Katz,AR,Tu '04]

Strongly interacting neutrino scenarios

- Bounds exploiting searches for deeply-penetrating particles applicable as long as $\sigma_{\nu N} \lesssim (0.5 \div 1) \text{ mb}$
 - For even higher cross sections, e.g. via sphaleron or brane production:
- ⇒ Strongly interacting neutrino scenario for the post-GZK events

[Berezinsky, Zatsepin '69]

A. Ringwald

COSMIC RAYS AT ULTRA HIGH ENERGIES (NEUTRINO?)

V. S. BERESINSKY and G. T. ZATSEPIN
Academy of Sciences of the USSR, Physical Institute, Moscow

Received 8 November 1968

The neutrino spectrum produced by protons on microwave photons is calculated. A spectrum of extensive air shower primaries can have no cut-off at an energy $E > 3 \times 10^{19} \text{ eV}$, if the neutrino-nucleon total cross-section rises up to the geometrical one of a nucleon.

Greisen [1] and then Zatsepin and Kusmin [2] have predicted a rapid cut-off in the energy spectrum of cosmic ray protons near $E = 3 \times 10^{19} \text{ eV}$ because of pion production on 2.7° black body radiation. Detailed calculations of the spectrum were made by Hillas [3]. Recently there were observed [4] three extremely energetic extensive air showers with an energy of primary particles exceeding $5 \times 10^{19} \text{ eV}$. The flux of these particles turned out to be 10 times greater than according to Hillas' calculations.

In the light of this it seems to be of some interest to consider the possibilities of absence of rapid (or any) fall in the energy spectrum of showerproducing particles. A hypothetic possibility we shall discuss* consists of neutrinos being the showerproducing particles at $E > 3 \times 10^{19} \text{ eV}$ due to which the energy spectrum of shower producing particles cannot only have any fall but even some flattening.

The neutrinos under consideration are originated in decays of pions, which are generated in collisions of cosmic ray protons with microwave photons. When calculating the neutrino spectrum the same assumptions were made as by Hillas [3]:

(1) The protons of high and extremely high energies are of extragalactic origin with an output of generation varying with time as t^{-8} after a certain starting time t_0^{**} ,

(2) The integral energy spectrum of generated protons is of the form $E^{-\gamma}$ up to an energy not less than 10^{22} eV .

* Cocconi was the first, who supposed that ultra high energy extensive air showers can be caused by neutrinos [5].

** The Hillas' assumptions about evolution of proton sources are based on Longair's [6] assumptions for evolution of radiogalaxies, the latter chosen to fit experimental data.




Fig. 1.

The calculated neutrino spectrum is represented by curve 3. It has the same spectrum exponent as the spectrum of generated protons. The calculations were made assuming that the pion originating in nucleon-microwave photon collision takes in average near 20% proton energy and the value $\gamma = 1.5$ was used. The calculated ratio of the neutrino intensity to that of the unmodified spectrum of protons (curve 1) at the same energy is $\sim 6 \times 10^{-2}$. We call "unmodified" a proton spectrum at present in the case when a red shift is the only kind of energy losses. The mentioned ratio does not depend on evolution of proton sources and the cosmological model. The proton spectrum at present is shown by curve 2. The curves 1 and 2 were obtained by Hillas using

Beyond the Standard Model, Bad Honnef, D

Strongly interacting neutrino scenarios


- Bounds exploiting searches for deeply-penetrating particles applicable as long as $\sigma_{\nu N} \lesssim (0.5 \div 1) \text{ mb}$
 - For even higher cross sections, e.g. via sphaleron or brane production:
- ⇒ Strongly interacting neutrino scenario for the post-GZK events

[Berezinsky,Zatsepin '69]

- Quantitative analysis:

[Fodor,Katz,AR,Tu '03; Ahlers,AR,Tu '05]

- Very good fit to CR data



Strongly interacting neutrino scenarios

- Bounds exploiting searches for deeply-penetrating particles applicable as long as $\sigma_{\nu N} \lesssim (0.5 \div 1) \text{ mb}$
 - For even higher cross sections, e.g. via sphaleron or brane production:
- ⇒ Strongly interacting neutrino scenario for the post-GZK events


[Berezinsky,Zatsepin '69]

- Quantitative analysis:

[Fodor,Katz,AR,Tu '03; Ahlers,AR,Tu '05]

- Very good fit to CR data
- Need steeply rising cross section, otherwise clash with nonobservation of deeply-penetrating particles

A. Ringwald



[Ahlers,A.R.,Tu '05]

[Han,Hooper '04]

- - - sphalerons

[Anchordoqui,Feng,Goldberg '02] - - - p -branes

[Burgett,Domokos,Kovesi-Domokos '04] ...string
excitations

Strongly interacting neutrino scenarios

- Bounds exploiting searches for deeply-penetrating particles applicable as long as $\sigma_{\nu N} \lesssim (0.5 \div 1) \text{ mb}$
 - For even higher cross sections, e.g. via sphaleron or brane production:
- ⇒ Strongly interacting neutrino scenario for the post-GZK events


[Berezinsky,Zatsepin '69]

- Quantitative analysis:

[Fodor,Katz,AR,Tu '03; Ahlers,AR,Tu '05]

- Very good fit to CR data
- Need steeply rising cross section, otherwise clash with nonobservation of deeply-penetrating particles

A. Ringwald



[Ahlers,A.R.,Tu '05]

[Han,Hooper '04]

- - - sphalerons


[Anchordoqui,Feng,Goldberg '02] - - - p -branes

[Burgett,Domokos,Kovesi-Domokos '04] ...string
excitations

4. The future ...

- Existing observatories for Extreme-
ly High Energy Cosmic ν 's

GLUE: Goldstone Lunar Ultra-high energy
neutrino Experiment




[<http://www.physics.ucla.edu/moonemp/public/>]

4. The future ...

- Existing observatories for Extreme-
ly High Energy Cosmic ν 's

GLUE: Goldstone Lunar Ultra-high energy
neutrino Experiment




[Gorham *et al.* '04]

4. The future ...

- Existing observatories for Extreme-
ly High Energy Cosmic ν 's

FORTE: F_ast O_n-orbit R_ecording of T_{ran}sient E_vents




[nis-www.lanl.gov/nis-projects/forte/]

4. The future ...

- Existing observatories for Extreme-
ly High Energy Cosmic ν 's

FORTE: F_ast O_n-orbit R_ecording of T_{ransient} E_vents




[Lehtinen *et al.* '04]

4. The future ...

- Existing observatories for Extreme-
ly High Energy Cosmic ν 's

ANITA-LITE: Prototype of ANtarctic Impulsive
Transient Antenna




[www.phys.hawaii.edu/anita/web/index.htm]

Beyond the Standard Model, Bad Honnef, D

4. The future ...

- Existing observatories for Extreme-
ly High Energy Cosmic ν 's


ANITA-LITE: Prototype of ANtarctic Impulsive
Transient Antenna



[cosray2.wustl.edu/tiger/index.html]

4. The future ...


- Existing observatories for Extreme-ly High Energy Cosmic ν 's provide sensible upper bounds on flux



4. The future ...

- Existing observatories for **E**xtreme-
ly **H**igh **E**nergy **C**osmic ν 's provide
sensible upper bounds on flux
- Upcoming decade: progressively lar-
ger detectors for **EHEC** ν 's

ANITA:



[www.ps.uci.edu/~anita/]

4. The future ...

- Existing observatories for **E**xtreme-
ly **H**igh **E**nergy **C**osmic ν 's provide
sensible upper bounds on flux
- Upcoming decade: progressively lar-
ger detectors for **EHEC** ν 's

WSRT: WeSterbork **R**adio **T**elescope




[Bacelar, ARENA Workshop '05]

4. The future ...

- Existing observatories for **E**xtreme-
High **E**nergy **C**osmic ν 's provide
sensible upper bounds on flux
- Upcoming decade: progressively la-
ger detectors for **EHEC** ν 's

WSRT: WeSternbork Radio T_Elescope



[Bacelar, ARENA Workshop '05]

4. The future ...

- Existing observatories for **E**xtreme-
ly **H**igh **E**nergy **C**osmic ν 's provide
sensible upper bounds on flux
- Upcoming decade: progressively lar-
ger detectors for **EHEC** ν 's


LOFAR:



[www.lofar.org]


4. The future ...

- Existing observatories for **E**xtreme-
ly **H**igh **E**nergy **C**osmic ν 's provide
sensible upper bounds on flux
- Upcoming decade: progressively lar-
ger detectors for **EHEC** ν 's




4. The future ...

- Existing observatories for **E**xtrremely **H**igh **E**nergy **C**osmic ν 's provide sensible upper bounds on flux
 - Upcoming decade: progressively larger detectors for **EHEC** ν 's
- $\Rightarrow E \geq 10^{16}$ eV:
 → **Astrophysics** of cosmic rays
- $\Rightarrow E \geq 10^{17}$ eV:
 → **Particle physics beyond LHC**




4. The future ...

- Existing observatories for **E**xtrremely **H**igh **E**nergy **C**osmic ν 's provide sensible upper bounds on flux
 - Upcoming decade: progressively larger detectors for **EHEC** ν 's
- $\Rightarrow E \geq 10^{16}$ eV:
 → **Astrophysics** of cosmic rays
- $\Rightarrow E \geq 10^{17}$ eV:
 → **Particle physics beyond LHC**
- $\Rightarrow E \geq 10^{21}$ eV:
 → **Cosmology**: relics of phase transitions; absorption on big bang relic neutrinos



Top-down scenarios for EHEC neutrinos


- Existence of superheavy particles with $10^{12} \text{ GeV} \lesssim m_X \lesssim 10^{16} \text{ GeV}$, produced during and after inflation through e.g.
 - particle creation in time-varying gravitational field



[Kolb, Chung, Riotto '98]

Top-down scenarios for EHEC neutrinos


- Existence of superheavy particles with $10^{12} \text{ GeV} \lesssim m_X \lesssim 10^{16} \text{ GeV}$, produced during and after inflation through e.g.
 - particle creation in time-varying gravitational field
 - decomposition of topological defects, formed during preheating, into their constituents



[Tkachev,Khlebnikov,Kofman,Linde '98]

Top-down scenarios for EHEC neutrinos


- Existence of superheavy particles with $10^{12} \text{ GeV} \lesssim m_X \lesssim 10^{16} \text{ GeV}$, produced during and after inflation through e.g.
 - particle creation in time-varying gravitational field
- ⇒ EHEC ν 's from decay or annihilation of superheavy dark matter (for $\tau_X \gtrsim \tau_U$)
- decomposition of topological defects, formed during preheating, into their constituents



[Berezinsky, Kachelriess, Vilenkin '97]

Top-down scenarios for EHEC neutrinos


- Existence of superheavy particles with $10^{12} \text{ GeV} \lesssim m_X \lesssim 10^{16} \text{ GeV}$, produced during and after inflation through e.g.
 - particle creation in time-varying gravitational field
 - ⇒ EHEC ν 's from decay or annihilation of superheavy dark matter (for $\tau_X \gtrsim \tau_U$)
 - decomposition of topological defects, formed during preheating, into their constituents
 - ⇒ EHEC ν 's from topological defects



[Bhattacharjee,Hill,Schramm '92]

Top-down scenarios for EHEC neutrinos


- **Injection spectra:** fragmentation functions $D_i(x, \mu)$, $i = p, e, \gamma, \nu$, determined via
 - Monte Carlo generators



[Birkel,Sarkar '98]

Top-down scenarios for EHEC neutrinos


- **Injection spectra:** fragmentation functions $D_i(x, \mu)$, $i = p, e, \gamma, \nu$, determined via
 - Monte Carlo generators
 - DGLAP evolution from experimental initial distributions at e.g. $\mu = m_Z$



[Particle Data Group '04]

Top-down scenarios for EHEC neutrinos


- **Injection spectra:** fragmentation functions $D_i(x, \mu)$, $i = p, e, \gamma, \nu$, determined via
 - Monte Carlo generators
 - DGLAP evolution from experimental initial distributions at e.g. $\mu = m_Z$ to $\mu = m_X$



[Fodor,Katz '01]

Top-down scenarios for EHEC neutrinos


- **Injection spectra:** fragmentation functions $D_i(x, \mu)$, $i = p, e, \gamma, \nu$, determined via
 - Monte Carlo generators
 - DGLAP evolution from experimental initial distributions at e.g. $\mu = m_Z$ to $\mu = m_X$
- ⇒ Reliably predicted!



[Aloisio,Berezinsky,Kachelriess '04]

Top-down scenarios for EHEC neutrinos

- **Injection spectra:** fragmentation functions $D_i(x, \mu)$, $i = p, e, \gamma, \nu$, determined via
 - Monte Carlo generators
 - DGLAP evolution from experimental initial distributions at e.g. $\mu = m_Z$ to $\mu = m_X$
- ⇒ Reliably predicted!
- **Spectra at Earth:**
 - for superheavy dark matter, injection nearby: $j_\nu \sim j_\gamma \sim j_p$




[Aloisio, Berezinsky, Kachelriess '04]

Top-down scenarios for EHEC neutrinos

- **Injection spectra:** fragmentation functions $D_i(x, \mu)$, $i = p, e, \gamma, \nu$, determined via
 - Monte Carlo generators
 - DGLAP evolution from experimental initial distributions at e.g. $\mu = m_Z$ to $\mu = m_X$
- ⇒ Reliably predicted!

- **Spectra at Earth:**

- for superheavy dark matter, injection nearby: $j_\nu \sim j_\gamma \sim j_p$
- for topological defects, injection far away: $j_\nu \gg j_\gamma \sim j_p$




[Aloisio,Berezinsky,Kachelriess '04]

Top-down scenarios for EHEC neutrinos

- How natural?
 - **Superheavy dark matter:** need symmetry to prevent fast X decay
 - * gauge $\Rightarrow X$ stable
 - * discrete \Rightarrow stable or quasi-stable

Top-down scenarios for EHEC neutrinos

- **How natural?**
 - **Superheavy dark matter:** need symmetry to prevent fast X decay
 - * gauge $\Rightarrow X$ stable
 - * discrete \Rightarrow stable or quasi-stable
 - **Topological defects:** generic prediction of symmetry breaking (SB) in GUT's, including fundamental string theory, e.g.
 - * $G \rightarrow H \times U(1)$ SB: monopoles
 - * $U(1)$ SB: ordinary or superconducting strings




[Rajantie '03]

Top-down scenarios for EHEC neutrinos

- **How natural?**


- **Superheavy dark matter:** need symmetry to prevent fast X decay
 - * gauge $\Rightarrow X$ stable
 - * discrete \Rightarrow stable or quasi-stable
- **Topological defects:** generic prediction of symmetry breaking (SB) in GUT's, including fundamental string theory, e.g.
 - * $G \rightarrow H \times U(1)$ SB: monopoles
 - * $U(1)$ SB: ordinary or superconducting strings
 - * $G \rightarrow H \times U(1) \rightarrow H \times Z_N$ SB: monopoles connected by strings



[Berezinsky '05]

Top-down scenarios for EHEC neutrinos


- Strong impact of measurement for
 - particle physics
 - cosmology



[Fodor,Katz,AR,Weiler,Wong,in prep.]

Top-down scenarios for EHEC neutrinos


- Strong impact of measurement for
 - **particle physics**
 - * GUT parameters, e.g. m_X
 - **cosmology**



[Fodor,Katz,AR,Weiler,Wong,in prep.]

Top-down scenarios for EHEC neutrinos


- Strong impact of measurement for
 - **particle physics**
 - * GUT parameters, e.g. m_X
 - * particle content of the desert, e.g. SM vs. MSSM
 - **cosmology**



[Barbot,Drees '02]

Top-down scenarios for EHEC neutrinos


- Strong impact of measurement for
 - **particle physics**
 - * GUT parameters, e.g. m_X
 - * particle content of the desert, e.g. SM vs. MSSM
 - * νN scattering at $\sqrt{s} \gg$ LHC
 - **cosmology**



[Tu '04]


Top-down scenarios for EHEC neutrinos

- Strong impact of measurement for
 - **particle physics**
 - * GUT parameters, e.g. m_X
 - * particle content of the desert, e.g. SM vs. MSSM
 - * νN scattering at $\sqrt{s} \gg \text{LHC}$
 - **cosmology**
 - * window on early phase transition
 - * Hubble expansion rate $H(z)$
 - * existence of the big bang relic neutrino background ($C\nu B$)



Top-down scenarios for EHEC neutrinos


- Strong impact of measurement for
 - **particle physics**
 - * GUT parameters, e.g. m_X
 - * particle content of the desert, e.g. SM vs. MSSM
 - * νN scattering at $\sqrt{s} \gg$ LHC
 - **cosmology**
 - * window on early phase transition
 - * Hubble expansion rate $H(z)$
 - * existence of the big bang relic neutrino background ($C\nu B$)



[Fodor,Katz,AR,Weiler,Wong,in prep.]

Top-down scenarios for EHEC neutrinos

- Strong impact of measurement for
 - **particle physics**
 - * GUT parameters, e.g. m_X
 - * particle content of the desert, e.g. SM vs. MSSM
 - * νN scattering at $\sqrt{s} \gg$ LHC
 - **cosmology**
 - * window on early phase transition
 - * Hubble expansion rate $H(z)$
 - * existence of the big bang relic neutrino background ($C\nu B$)



[Fodor,Katz,AR,Weiler,Wong,in prep.]

5. Conclusions

- Exciting times for extremely high energy cosmic rays and neutrinos:
 - many observatories under construction
⇒ appreciable event samples
- Expect strong impact on
 - astrophysics
 - particle physics
 - cosmology

