The Search for Exotic Baryons at the HERMES Experiment

by

Wouter Deconinck

A dissertation submitted in partial fulfillment
of the requirements for the degree of
Doctor of Philosophy
(Physics)
in The University of Michigan
2008

Doctoral Committee:

Professor Wolfgang B. Lorenzon, Chair
Professor Gordon L. Kane
Professor Michael J. Longo
Professor Jianming Qian
Associate Professor Arthur M.F.W. Verhoogt

ABSTRACT

The Search for Exotic Baryons at the HERMES Experiment

by
Wouter Deconinck

Chair: Wolfgang B. Lorenzon

One of the interesting questions of Quantum Chromodynamics, the theory that governs the interactions between quarks and gluons, has been whether it is possible to observe hadrons which can not be explained as a combination of only two or three valence quarks. In numerous searches the existence of these exotic hadrons could not be confirmed. Recently, calculations based on the quark soliton model predicted the narrow exotic baryons Θ^+ and Ξ^{--}. A narrow resonance identified as the Θ^+ was observed by several experiments at the predicted mass of 1540 MeV, but later followed by several dedicated experiments that could not confirm these positive results.

At the HERMES experiment a search for the quasi-real photoproduction of the exotic baryon Θ^+ on a deuterium target and the subsequent decay through $pK^0_S \rightarrow p\pi^+\pi^-$ revealed a narrow resonance in the pK^0_S invariant mass distribution at 1528 MeV. In the search for the corresponding antiparticle Θ^- the result is consistent with zero events.

In this thesis we present the search for the exotic baryon Ξ^{--} on a deuterium target in the data sample used for the observation of the Θ^+. An upper limit on the cross section of the exotic baryon Ξ^{--} is determined. The search for the exotic baryon Θ^+ on hydrogen and deuterium targets at the HERMES experiment is extensively discussed. The event mixing method can be used to estimate the distribution of background events. Several difficulties with this method were addressed, but the background description in the case of the exotic baryon Θ^+ remains unconvincing. Between the years 2002 and 2005 the HERMES experiment operated with a magnetic holding field around the hydrogen target. A method for the reconstruction of displaced vertices in this field was developed. The data collected during the years 2006 and 2007 offer an integrated luminosity that is several times higher than in previous data sets. After investigating all data sets collected with the HERMES experiment on hydrogen and deuterium targets, we are not able to observe a resonance peak at an invariant mass of 1540 MeV consistent with the exotic baryon Θ^+.

ACKNOWLEDGEMENTS

This dissertation would not have been possible without the help of numerous faculty, colleagues, friends and family. Perhaps unknowingly, many individuals contributed to the successful completion of this dissertation, through their expressions of support or concrete assistance in solving problems. Importantly, this also includes a large number of educators, teachers, and mentors who helped shape my approach to this research project before I entered the graduate program at the University of Michigan. I would like to express my gratitude to all of them. Several individuals deserve special mention for their contributions to this dissertation.

Wolfgang Lorenzon has been an exceptional advisor, a constant source of support and inspiration. He put great trust in my independence and always left me the freedom to explore, but was careful to remind me when I was losing sight of my goals. Whenever I asked Wolfgang for his opinion on my work, he offered his insights and comments quickly and accurately. His strategical advice transcended my graduate career; he provided me with opportunities to connect with other academics and was crucial in the formation of my post-doctoral plans. I am very grateful for having had the opportunity to work with Wolfgang. During my future career I can only hope to be as good a mentor as Wolfgang.

My thanks go out to the other members of my dissertation committee for their interest in my work, for the different viewpoints they provided, and for their suggestions for improvement and changes. Gordy Kane provided me with the theoretical background in standard model physics (and beyond) in several courses on particle physics, and continued to be a source of enlightening insight. Mike Longo's perspective on pentaquarks inspired me to carefully review the assumptions underlying this dissertation research, leading to the interesting discovery of ghost tracks in the HERMES data. One of the most interesting courses of my graduate track was taught by Jianming Qian on experimental collider physics. He introduced many of the experimental techniques, from particle detection to statistical analysis, and his lecture notes remain a handy reference. Arthur Verhoogt gracefully accepted the "challenge" of being the ever cheerful cognate member on my dissertation committee after a rather long time away from physics. His questions made me think more about the meaning of the presented results on a very fundamental level.

The original idea of dissertation research at the HERMES experiment was suggested by the spokesperson at the time, Dirk Ryckbosch. His course on medium energy physics at the University of Gent was what sparked my interest in subnuclear physics. Without the inspiration of Dirk I would not have pursued a graduate program at the University of Michigan.

I am very much indebted to my colleague and supervisor Avetik Airapetian. In our office "container" at DESY he was always there (as a walking encyclopedia or even analysis guru) to answer questions and to direct me to the right path when solving problems. Together with Riccardo Fabbri, we spent many evenings in the East Hall, trying to solve odd hardware or software malfunctions in the LPOL system. I want to thank my LPOL friends Avetik and Riccardo not only for being such great colleagues, but also for reminding me of the joys of good Italian food and *caffè*. Or course, I should have put on my soccer shoes more regularly to join Avetik and Riccardo...

I wish to thank all of my colleagues and friends at the HERMES experiment for the interesting discussions, for the critical observations, and for valuable suggestions. In particular the regulars at the "offsite" meetings, arguably more useful than any other meeting, will be missed. Many thanks to Caro and Charlotte for the numerous conversations that had nothing to do with physics, and for the ones that did. Many of the ideas in this dissertations resulted from suggestions by Achim, Alex, Beni, Brian, Caro, Charlotte, Dominik, Elke, Jim, Josh, Larry, Markus, Rebecca, Roberto, Steve, and Yves. Any mistakes result from my misinterpretations of their great ideas. A special thank you goes to Evelyn and Sabine for all their practical advice on wrestling with the German bureaucracy.

I would like to thank my parents, Hilde and Johan, for giving me the opportunities to study and explore, for teaching me to keep an open mind, and for providing me with an example of how to live.

Finally, I most want to thank Susanne who was always there for me, even while we were separated by oceans and time zones. Without her constant support I would not have been able to accomplish this.

TABLE OF CONTENTS

LIST OF FIGURES

Figure

LIST OF TABLES

Table

CHAPTER I

Introduction

In the Standard Model of particle physics the fundamental fermionic constituents of matter are organized in three progressively more massive generations. Each generation of particles consists of two quarks with electromagnetic charge $+\frac{2}{3}$ and $-\frac{1}{3}$ (up u and down d, charm c and strange s, top t and bottom b), a lepton with by definition electromagnetic charge -1 (electron e, muon μ, tau τ), and the corresponding neutral neutrino (ν_e, ν_μ, ν_τ). The particles are subjected to the four fundamental forces: the gravitational force, the electromagnetic interaction responsible for electricity and magnetism, the weak interaction which is important in nuclear processes such as β-decay, and the strong interaction that binds the protons and neutrons in the atomic nucleus. Even though gravity is the most apparent force at macroscopic scales, it is much weaker than the other forces at microscopic scales and will be ignored here.

The fundamental forces act by the exchange of force-mediating particles or gauge bosons. The massless photon γ mediates the electromagnetic force between electrically charged particles. The theory of Quantum Electrodynamics (QED) describes the interactions mathematically as the abelian $U(1)_{em}$ gauge theory. Because of the small value of the electromagnetic coupling constant $\alpha \approx \frac{1}{137}$, the theory of QED lends itself well to perturbative expansion. This allows for the calculation of experimental observables to a very high precision.

The weak force is carried by the three massive vector bosons $W^\pm$ and Z^0 and acts on all fermionic fundamental particles listed above.[1] Because the $W^\pm$ bosons are charged, they are affected by the electromagnetic interaction. The masses of vector bosons are generated by spontaneous symmetry breaking at energies below approximately 10^2 GeV. Above this energy the electromagnetic interaction and the weak interaction are unified in the electroweak interaction with the symmetry group $SU(2) \otimes U(1)_Y$, where Y is the hypercharge. Just as the electromagnetic charge, the hypercharge of a particle indicates how strongly it will couple to the electroweak gauge bosons. At lower energies the electroweak symmetry is spontaneously broken down to $U(1)_{em}$ by the Higgs mechanism, which not only generates the masses of the Goldstone bosons $W^\pm$ and Z^0 but also of all other fermions in the Standard Model, because the interaction of the Higgs field is proportional to the fermion mass.

In addition to the electromagnetic charge and the hypercharge, the quarks (but not the leptons and neutrinos) carry a color charge which can take on one of the values red r, green g, or blue b, the sum of which cancels out. The antiquarks carry an anticolor charge $\overline{r}$, $\overline{g}$, or $\overline{b}$. The eight massless and electromagnetically neutral gluons g mediate the strong force by coupling to the color charge of the quarks. The gluons carry a combined color-anticolor charge. The interaction of the gluons with the quarks, and of the gluons with each other, is described by the theory of Quantum Chromodynamics (QCD), a non-abelian gauge theory with color symmetry group $SU(3)_c$ (for a review see reference [1]).

The self-interaction of gluons, a feature of the $SU(3)$ group structure of QCD that is not present in the simpler $U(1)$ structure of QED (at least at tree level), causes the interaction between two quarks to become weaker when the quarks are closer to each other, and stronger when they are further apart. This is opposite to the behavior of QED, where the effect of electromagnetic charge screening reduces the interaction at larger distances. This particular effect of QCD is called asymptotic freedom, because at short distances the

[1] Naively the strength of the interaction is inversely proportional to the squared mass of the mediating boson. The larger mass of the weak bosons is responsible for the weakness of the weak force.

quarks are quasi-free. Similarly, due to the increase of the interaction at large distances, it is impossible to remove a colored object from a colorless bound state, a property known as color confinement. QCD only allows colorless free objects; all colored objects have to be in a bound state such that their total color charge is canceled.

1.1 Mesons and baryons

The colorless bound states of quarks, antiquarks and gluons in QCD are called hadrons. The simplest colorless combinations of quarks and antiquarks are the *mesons* consisting of a quark and an antiquark ($q\overline{q}$) and the *baryons* composed of three quarks (qqq). These two or three quarks, which carry most of the hadron's momentum, are called the valence quarks. Solely these two or three valence quarks determine the physical observable quantum numbers of the bound state (such as the spin J and the strangeness S). In addition, the valence quarks are embedded in a sea of $q\overline{q}$ pairs, which have no net contribution to the quantum numbers of the bound state, but which can be probed in deep-inelastic scattering experiments. For example, a proton consists of one d and two u valence quarks but $s\overline{s}$ pairs are created and disappear constantly in the quark sea.

Because the masses of the c, b and t quarks are large compared to the masses of the u, d and s quarks ($m_u \approx 3\,\mathrm{MeV}$, $m_d \approx 6\,\mathrm{MeV}$, $m_s \approx 100\,\mathrm{MeV}$, $m_c \approx 1.24\,\mathrm{GeV}$, the masses for the b and t quarks are even larger [1]), they are usually ignored in treatments of the lightest hadrons. When only the three lightest quarks u, d and s are considered, the flavor symmetry group $SU(3)_f$ describes the hadronic bound states. All hadronic states in a perfect symmetry group are expected to have the same energy. $SU(3)_f$ is not an exact symmetry owing to the larger mass of the s quark, but very useful as an approximation. When we also include the c quark, we obtain the badly broken $SU(4)_f$ flavor symmetry. In this work we will limit ourselves to the light quarks only and consider the flavor symmetry group $SU(3)_f$.

Following the group-theoretical relation $\mathbf{3} \otimes \mathbf{3} = \mathbf{8} \oplus \mathbf{1}$ in the flavor symmetry group $SU(3)_f$, the nine possible quark-antiquark combinations $q\overline{q}$ can be grouped in a singlet and an octet of light mesons. Because the s quark has a larger mass than the u and d quarks, the symmetry is not perfect and the mesons with non-zero strangeness are heavier than the mesons without strangeness.[2] Mesons with equal strangeness are approximately degenerate in mass, because of the $SU(2)$ isospin symmetry between u and d quarks. When we include the c quark, we obtain a 15-plet according to $\mathbf{4} \otimes \mathbf{4} = \mathbf{15} \oplus \mathbf{1}$. The mass splittings are now even larger, and the states with charm content are largely independent of the states already obtained in the $SU(3)_f$ singlet and octet.

If we limit ourselves to the ground states without orbital excitations, in other words to bound states with $\ell = 0$, the total spin $J = \ell + s$ of the mesons can be $J = 0$ or $J = 1$ for antiparallel or parallel quarks spins (the sum of the spins is represented by the lower case s, to avoid confusion with the strangeness S). In both spin configurations the meson states can be classified in a singlet and an octet. In figure 1.1 the $J = 0$ states or pseudoscalar mesons are shown in the left panel and the $J = 1$ states or vector mesons are shown in the right panel. The isoscalar singlet state and the two isoscalar states at the center of the octet mix with each other because they have identical quantum numbers (*i.e.* $Q = 0$, $S = 0$).

The multiplets for the baryons consisting of three quarks are determined similarly to the classification scheme of the mesons. The relation $\mathbf{3} \otimes \mathbf{3} \otimes \mathbf{3} = \mathbf{10}_S \oplus \mathbf{8}_M \oplus \mathbf{8}_M \oplus \mathbf{1}_A$, however, does not produce separate ground state multiplets with and without angular momentum ℓ. By treating the quarks with spin up and spin down independently the flavor-spin symmetry group $SU(6)_{fs}$ can be used to determine the ground state multiplets. The baryons are now described by the relation $\mathbf{6} \otimes \mathbf{6} \otimes \mathbf{6} = \mathbf{56}_S \oplus \mathbf{70}_M \oplus \mathbf{70}_M \oplus \mathbf{20}_A$ and the ground state 56-plet decomposes according to $\mathbf{56} = \mathbf{10} \oplus \mathbf{8}$ in an octet with spin $J = \frac{1}{2}$ and a decuplet with spin $J = \frac{3}{2}$. In figure 1.2 the ground state baryon octet is shown on the left and the decuplet on the right. As in the case of mesons, the masses of baryons with the same strangeness S are almost identical because of isospin symmetry. The differences between the masses of baryons with different strangeness is larger.

Because the strong coupling constant α_S is of the order of unity at the energies of the lightest bound

[2] The strangeness S of a hadron is proportional to the number of strange quarks. It follows the convention that the flavor of a quark has the same sign as its electromagnetic charge. Because the strange quark has negative electromagnetic charge $-\frac{1}{3}$, the strangeness of the s quark itself is $S = -1$.



Figure 1.1: The ground state singlets and octets of the light $J = 0$ pseudoscalar mesons (left panel) and $J = 1$ vector mesons (right panel) made of only u, d and s quarks without angular momentum ($\ell = 0$).



Figure 1.2: The ground state $J = \frac{1}{2}$ octet (left panel) and $J = \frac{3}{2}$ decuplet (right panel) of the light baryon states made of only u, d and s quarks without angular momentum ($\ell = 0$).

quark states, a perturbative approach to QCD is not applicable to calculate the masses of the lightest mesons and baryons.[3] At higher energies (and thus shorter distance scales) the strong coupling constant decreases and reaches a value $\alpha_S(M_Z) = 0.12$ at the mass of the Z boson. At higher energies perturbative QCD can be used for high-precision tests of QCD, similar to what is possible in QED. In the non-perturbative region several phenomenological approaches can be used for the determination of the hadronic masses. In the constituent quark model (CQM) the quarks are 'dressed' or given a larger mass ($m_{u/d,CQM} \approx 350\,\mathrm{MeV}$, $m_{s,CQM} \approx 470\,\mathrm{MeV}$) and their interaction in a hadronic potential is determined [2].

1.2 Exotic hadrons

QCD does not prohibit the existence of colorless hadronic states with more than two or three quarks. In fact, studies of the nucleon structure point to a significant contribution from quark-antiquark pairs to the total momentum and spin of the nucleon. The naive quark model with bound states of only two or three valence quark seems incomplete. When one realizes that the quarks in mesons and baryons are bound together by the gluons, also gluons should play an important role in the description of bound states.

The hadrons that fall outside of the expectations of the naive quark model are called manifestly exotic hadrons. Their quantum numbers can not be explained by a bound state of only two or three quarks. Crypto-exotic hadrons are composed of more than two or three quarks, but mix with a regular hadron with the same quantum numbers. The exotic mesons can be classified as glue balls (gg), hybrids ($q\overline{q}g$), and tetraquarks ($q\overline{q}q\overline{q}$). Pentaquarks ($qqqq\overline{q}$) are examples of exotic baryon states.

For mesons the quantum numbers for the total spin J, the parity P, and the charge conjugation eigenvalue C are only allowed to have a limited set of values. The configurations $J^{PC} = 0^{--}$, 0^{-+}, 1^{-+}, 2^{+-},... cannot be obtained in the naive quark model. The observation of several states with $J^{PC} = 1^{-+}$ since the 90s points unambiguously to the existence of exotic mesons, although the exact nature of these exotic mesons is still unknown.[4]

The charge conjugation eigenvalue C is not a good quantum number for baryons, and all combinations of the total spin J and the parity P are allowed. However, an exotic baryon can be identified with a combination of the charge Q and the strangeness S. Since no three-quark baryons with positive strangeness exist, a baryon with charge $Q = +1$ and strangeness $S = +1$ must have the minimal quark configuration $uudd\overline{s}$. This exotic baryon will be referred to as Θ^+. Similarly, the minimal quark content of the exotic baryon Ξ^{--} with charge $Q = -2$ and strangeness $S = -2$ is $ssdd\overline{u}$. In the chiral quark soliton model, discussed in section 2.2, the lightest exotic baryons with spin $J = \frac{1}{2}$ are organized in an antidecuplet, shown in figure 1.3. Due to mixing of the internal member of this antidecuplet with a non-exotic octet, the only manifestly exotic baryons are located on the corners of this antidecuplet.

So far, experimental searches for exotic baryons have been unsuccessful. Early experiments in the 1970s were not able to confidently prove the existence of exotic baryons, although a number of (often contradictory) results in kaon-nucleon scattering reported positive results. In recent years a new wave of positive experimental results and contrasting null results about the exotic baryons Θ^+ and Ξ^{--} has revitalized the discussion, but it is too early to draw conclusions about the existence of these exotic baryons.

At the HERMES experiment in Hamburg, Germany, a search for the exotic baryons Θ^+ and Ξ^{--} was performed [3, 4]. A resonance peak in the decay channel corresponding to the Θ^+ was observed at a mass of $M = 1528\,\mathrm{MeV}$, but no Ξ^{--} resonance could be found. Since these result were published in 2003 and 2004, a large amount of additional data has been collected. This dissertation investigates the search for exotic baryons in these data.

1.3 Overview

In chapter II an overview will be given of the experimental and theoretical efforts to observe, refute or more broadly understand exotic baryons. The evidence in favor of the recently observed exotic baryons Θ^+

[3] The advances in lattice QCD calculations make it possible to determine the masses of the bound states directly from the $SU(3)$ Lagrangian of QCD. This is however not up to par yet with the results from phenomenological quark models.

[4] If each of the reported 1^{-+} state were confirmed, it would result in an overpopulation of the 1^{-+} hybrid multiplet. The experiment GlueX at JLab is proposed to investigate exotic mesons [5].



Figure 1.3: The ground state antidecuplet of the exotic $J = \frac{1}{2}$ baryon states, as predicted by the chiral quark soliton model (see section 2.2.4).

and Ξ^{--} is examined and critically compared with null results. The theoretical efforts in hadronic physics have lead to a number of different approaches in calculating the mass and the width of exotic baryons. The most well-known approach is without doubt the chiral quark soliton model, but the diquark model and the diquark-triquark model are also presented.

The HERMES experiment is introduced in chapter III. It is located in the East Hall of the HERA collider at the DESY laboratory, where energetic electrons or positrons are collided on a fixed gas target. The scattered lepton and the collision fragments are detected in a conventional forward spectrometer with good particle identification capabilities. Although the polarization of the lepton beam is not relevant for the analysis of exotic baryons, the longitudinal polarimeter is discussed in the light of proposed future lepton polarimeter designs.

The analysis of the exotic baryons at the HERMES experiment is presented in chapter IV. Similar to the original analyses of the exotic baryons Θ^+ and Ξ^{--} at the HERMES experiment [3, 4], a search for the exotic baryon Ξ^{--} is presented. The photoproduction cross section ratio of the $\Lambda(1520)$ and $\overline{\Lambda}(1520)$ hyperons is used to determine the expected number of observed Θ^- baryons. For the analysis of the data collected with a transversely polarized hydrogen target, a tracking algorithm was developed for displaced vertices in a homogeneous field. Finally, the technique of event mixing for the determination of distributions of background events is discussed.

In the final chapter V the results of this work are summarized. The analysis method, involving several novel aspects, is critically reviewed. The observation of exotic baryons at the HERMES experiment, in particular of the Θ^+ and the Ξ^{--}, is assessed.

CHAPTER II

The Search for Exotic Baryons

The search for exotic baryons knows a long history, starting immediately after the formulation of the quark model and development of the mathematical structure of QCD. The assumedly lightest five-quark exotic baryon, with a quark configuration $uudd\overline{s}$ and with positive strangeness $S = +1$, has been the focus of the majority of the experimental searches, and it receives therefore also an important position in the theoretical models. In early searches exotic baryons with strangeness $S = +1$ were referred to as Z^* resonances, but in the recent results the symbol Θ^+ has become the accepted notation. The other manifestly exotic baryons in the antidecuplet composed of only u, d and s quarks are denoted by Ξ^{--} for the $ddss\overline{u}$ quark configuration and by Ξ^+ for the $uuss\overline{d}$ quark configuration.[1] Currently only one experiment reported evidence for these states, in contrast with numerous null results.

In this chapter we review the experimental and theoretical results about the light exotic baryons, consisting of only u, d and s quarks.[2] In section 2.1 the older KN partial wave analyses and several recent spectroscopic results are discussed, comparing the positive and null results where possible. In section 2.2 we review the theoretical predictions for exotic baryons and the recent developments that aim to reconcile the seemingly contradictory experimental results.

2.1 Experimental results

Since the early days of QCD, experimental searches for baryons composed of more than three quarks have been performed. In particular, the observation of a baryon with a positive strangeness S would be a clear signature of an exotic baryon, because such a baryon can only be explained by the combination of five or more quarks.

2.1.1 Early controversy about exotic baryons

Using secondary charged kaon beams, a large amount of data has been collected in scattering of kaons on protons and deuterons in the late 60s and 70s. Several indications for baryon resonances with positive strangeness, identified as Z^* resonances at that time,[3] were observed in the partial wave analysis of these data, but none of the reports was ever sufficiently significant to stand beyond the doubt created by contradicting experimental results [6]. The widths of the observed Z^* resonances were usually of the order of 100 MeV, both for the isoscalar resonances Z_0^* around 1800 MeV and for the isovector resonances Z_1^* at 1900 MeV and higher. Due to these large widths, the resonances are now understood as non-exotic pseudo-resonances induced by the opening of $K\pi N$ channels [7].

[1] The Particle Data Group [1] refers to the exotic baryons Ξ^{--} and Ξ^+ with the symbol Φ, but this notation has not found a lot of adoption in the experimental community. In this work we will continue to use the symbol Ξ to avoid inconsistencies with the cited sources.

[2] After the first reports of the observation of the exotic baryon Θ^+ in 2003, similar exotic states with c or b quarks instead of the s quark were predicted [8, 9, 10]. The experimental observation of a resonance decaying to $D^{*-}p$ at the H1 experiment was interpreted as evidence for the charmed exotic baryon Θ_c^0 [11], but in subsequent searches no other experiment was able to confirm this observation. This charmed exotic baryon Θ_c^0 will not be discussed in this work.

[3] The isospin of the resonance is often indicated as a subscript, *i.e.* Z_0^* or Z_1^*.



Figure 2.1: Change of the overall χ^2 when including a narrow P_{01} resonance in a global partial wave analysis of the KN scattering data is only acceptable for resonance widths Γ below a few MeV. The small vertical lines in the right panel indicate the position of experimental observations of the exotic baryon $\Theta^+(1540)$. In the right panel the values for $\Delta\chi^2$ at 1535 MeV are off-scale. Taken from reference [12].

Starting with its 1986 edition of the Review of Particle Physics [13], the Particle Data Group decided to omit the experimental results for the exotic baryons Z^*. The "general prejudice against baryons not made of three quarks" and "lack of experimental activity" were cited as reasons why a decision on the existence of exotic baryons was not to be expected in the foreseeable future. The discussion indeed lay dormant for more than 15 years, until the first of the most recent results was presented in 2003.

The K^+p and K^+d scattering databases were recently re-examined in the light of the evidence for a surprisingly narrow exotic baryon resonance Θ^+ with a mass of approximately 1540 MeV. An upper limit on the width of the exotic baryon Θ^+ of a few MeV was derived, based on fluctuations in the measured cross sections [14, 15, 16]. In a re-examination of the partial wave analysis of the KN scattering data, the new narrow exotic baryons were included and the difference in the goodness-of-fit parameter χ^2 was determined [12]. In figure 2.1 the change of the overall χ^2 of a global fit to the KN scattering data is shown when including a narrow P_{01} resonance. For intrinsic resonance widths Γ above 2 MeV, the increase of the total χ^2 is too large, and only for widths below a few MeV the inclusion of a narrow resonance is possible. Similarly, when including a narrow S_{01} or P_{03} resonance only a width below 1 MeV is consistent with the KN scattering data. This puts a strong constraint on the theoretical models trying to explain the observed exotic baryons.

2.1.2 Experimental results since 2003

Prompted by the theoretical predictions of the chiral quark soliton model [17], a new series of experimental results claiming the observation of the exotic baryon Θ^+ at a mass of approximately 1540 MeV has been published since 2003. This resonance with positive strangeness $S = +1$ can only be explained as a five-quark state with configuration $uudd\bar{s}$, and corresponds to the Z^* resonances discussed in the 70s. Following common practice for baryons, the determined mass is added to the symbol of the resonance to distinguish it from other resonance with the same structure and thus the same symbol, but with a different mass due to orbital excitations. When discussing the observed exotic resonance state around 1540 MeV we will therefore write $\Theta^+(1540)$. The isospin structure of the exotic resonance $\Theta(1540)$ has not been determined conclusively experimentally, with most experiments unable to observe a possible isopartner $\Theta^{++}(1540)$ but some results pointing to the existence of the isospin partner [18]. In the chiral quark soliton model the exotic baryon Θ was predicted to be an isoscalar (as the Z_0^* state).

Initially, the positive results created a lot of enthusiasm in the hadron spectroscopy community. The number of theoretical papers published on the subject, trying to reconcile the different observed masses and attempting to determine the possible production mechanisms rose quickly. However, slowly null results began to trickle in, most of them with better statistical precision and placing strict limits on the allowed cross section or intrinsic width. The apparent disagreement between the positive results and the null results turned around the initial enthusiasm for the experiments that presented evidence in favor. A few experiments could not repeat their previous results with data collected in very similar conditions. Only a small number of experiments has presented additional data in different reaction channels or kinematic regimes in which they confirmed their earlier sightings.

Now, when looking back at the large amount of experimental results, some positive, several null, we can draw some conclusions. Some experimental results were already discarded as invalid, others are in the process of being confirmed or refuted. In this section some experimental results are discussed in more detail. Completeness was not pursued, only experimental results deemed interesting or significant in the eyes of the author were included. Since it is difficult to order the experiments consistently according to any experimental characteristic, they are presented chronologically. But because some experiments published several results, the chronological line has been broken at several points.

2.1.3 Photoproduction on neutrons at the LEPS experiment

In the beginning of 2003, the first recent claim for the observation of the exotic baryon $\Theta(1540)$ was presented by the Laser-Electron Photon experiment at SPring-8 (LEPS) in Japan. At the LEPS experiment high-energetic photons are produced by Compton back-scattering of laser photons from the 8 GeV electrons in the SPring-8 synchrotron [19]. The frequency of the 351 nm Ar laser limits the Compton photon energy to values below 2.4 GeV. The photon energy is determined by measuring the momentum of the scattered electrons using one of the bending magnets of the SPring-8 storage ring. Only for photons with an energy above 1.5 GeV, the scattered electron can be detected.[4]

The photons strike a fixed target cell and the reaction products are detected by the LEPS spectrometer. The design of the spectrometer is optimized for the detection of photoproduced ϕ mesons at forward angles. A silicon strip vertex detector and three drift chambers are used to track charged particles in a magnetic dipole field and determine their momentum. A time-of-flight (TOF) scintillator detector and a Čerenkov counter are used to distinguish electrons from pions.

For the first publication on exotic baryons at the LEPS experiment [20], a 0.5 cm thick plastic scintillator (SC) with an approximately equal number of hydrogen and carbon nuclei was used as the target for the photon beam. This scintillator target was located downstream from the actual liquid hydrogen (LH_2) target for the study of the photoproduction of ϕ mesons. In 2003 the LH_2 target was replaced by a longer target cell which could also hold liquid deuterium (LD_2), to allow for the study of reactions on neutrons without the need for a complicated Fermi motion correction [21, 22].

First results on the scintillator target

For the analysis of the first data sample, collected on the SC target, events with a K^+K^- pair in the detector were considered [20]. The scattered nucleon was reconstructed from the missing mass $MM(\gamma K^+K^-)$,

[4] Because of energy conservation, photons with a lower energy correspond to scattered electrons with an energy closer to the beam energy. The separation between the scattered electron and the electron beam after the first dipole magnet in the synchrotron is then too small and the electron is outside the acceptance of the tagging counter.



Figure 2.2: The distribution of the missing masses $MM(\gamma K^+)$ (left panel) and $MM(\gamma K^-)$ (right panel) for events originating in the SC target and corrected for Fermi motion in the C nucleus (solid line in both panels). When an additional proton track is required in the final state, the $\Lambda(1520)$ resonance becomes visible in the left panel (dashed line). In the right panel, the background contribution is estimated using events with the initial vertex in the LH_2 target (dotted line). Taken from reference [20].

and a veto on a scattered proton track was used to suppress the photo-nuclear reaction $\gamma p \to K^+K^-p$ off protons in the H or C atoms. Assuming the reaction $\gamma n(^{12}\mathrm{C}) \to K^+K^-(n)$ the missing mass $MM(\gamma K^\pm)$ corresponds now to the invariant mass $M(nK^\mp)$. The Fermi motion of the target neutron in the atomic nucleus leads to smearing of the kinematic variables and is corrected for.[5]

In the distribution of the missing mass $MM(\gamma K^+)$, shown in the left panel of figure 2.2, the $\Lambda(1520)$ resonance is not visible in the sample with the proton veto (solid lines), but becomes visible when a proton hit is required (dashed lines) as expected for the reaction $\gamma p(^{12}\mathrm{C}) \to \Lambda(1520)K^+ \to pK^-K^+$. The distribution of the missing mass $MM(\gamma K^-)$, shown in the right panel of figure 2.2, for events originating in the SC target (solid lines) exhibits a peak at a mass of $1540 \pm 10\,\mathrm{GeV}$ with a width smaller than 25 MeV and a naive significance of $\frac{S}{\sqrt{B}} = 4.6\,\sigma$.[6] The structure disappears when the events are not required to originate in the SC target, but instead in the LH_2 target (dotted lines).

Dedicated setup with the liquid deuterium target

In 2003 the target system at the LEPS experiment was upgraded to allow for data collection on liquid hydrogen and liquid deuterium targets with increased thickness [22]. The hydrogen target allowed to estimate the background distributions for reactions on the neutrons, and coherent reactions on deuterons could now be studied without the need for a complicated Fermi motion correction.

The original result for photoproduction on neutrons was confirmed with higher statistical precision, as shown in figure 2.3. This result still depends on the correction for the Fermi motion of the neutron in

[5] Only after the Fermi motion correction a clean separation of the Λ hyperon at 1115 MeV and the Σ^- hyperon at 1190 MeV, both decaying to $K^+\pi^-$, is possible due the improvement in the resolution.

[6] Several expressions are used to calculate the statistical significance of experimental results. The statistical significance of a hypothesis represents the probability that the observation is a random fluctuation, and is usually expressed in units of standard deviations for a normal distribution. The naive estimator $\frac{S}{\sqrt{B}}$ for the significance of a peak with S events on top of B background events is only valid when the background B is known and large. It overestimates the true statistical significance by assuming that the resonance is there, by ignoring the uncertainty in the background description and the change from Gaussian to Poisson distributions for a low number of events. The alternative expression $\frac{S}{\sqrt{B+S}}$ is sometimes used to avoid the assumption that the resonance exists. Only a fully correlated statistical treatment of peak and background allows to calculate the expression $\frac{S}{\delta B}$, where δB is now the uncertainty in the number of background events in the peak region.



Figure 2.3: The missing mass distributions of $MM(\gamma K^+)$ (left panel) and $MM(\gamma K^-)$ (right panel) in the new data, corrected for Fermi motion in the deuterium nuclei of the LD_2 target (solid lines). The values on the horizontal axis are in GeV, with unspecified bin size. The mixed-event background is shown as the dashed line. Taken from references [22, 23].

the deuterium atom. In the left panel the distribution of the missing mass $MM(\gamma K^+)$ with a prominent $\Lambda(1520)$ peak is shown. The distribution of the missing mass $MM(\gamma K^-)$ is shown in the right panel with an excess of events at 1.53 GeV. The dashed line represents the mixed event background, presumably normalized in the invariant mass region below 1.5 GeV.

Inspired by the analysis method used at the CLAS experiment (see section 2.1.6), the LEPS experiment also searched for the $\Theta(1540)$ in the coherent reaction $\gamma d \to pK^- X$ on a deuterium target using the missing mass $MM(\gamma pK^-)$. When only events are selected with an invariant mass $M(pK^-)$ in the region corresponding to the $\Lambda(1520)$ hyperon, the distribution for the missing mass $MM(\gamma K^- p)$ in figure 2.4 is obtained. The total background (indicated by the red line) has a component modeled by the sidebands of the $\Lambda(1520)$ hyperon (blue line) and a component from $\Lambda(1520)$ production estimated from the LH_2 data (green line). A peak structure at 1.53 GeV and a broad bump at 1.6 GeV are observed.

In summary, the LEPS experiment currently claims evidence for the exotic baryons $\Theta(1540)$ in several different reaction channels. The results seem to indicate simultaneous production of the non-exotic $\Lambda(1520)$ hyperon and the exotic baryon $\Theta(1540)$.

2.1.4 Charge-exchange reaction K^+Xe $\to K^0_S p$Xe$'$ at the DIANA experiment

The bubble chamber of the DIANA experiment is filled with liquid Xe and is exposed to a beam of positive kaons K^+ with an energy of 850 MeV separated from the ITEP proton synchrotron. Charged particles are photographically detected by the ionized tracks they leave in the bubble chamber. Using special stereo-projectors the tracks can be fully reconstructed. The momentum of the particles is derived from the range the track covers in the liquid Xe. The strangeness of the final state is fixed to $S = +1$ by the incoming kaon K^+.[7]

First half of the available data sample

The invariant mass $M(K^0_S p)$ was measured using to the charge-exchange reaction $K^+ n(\text{Xe}) \to K^0_S p(\text{Xe}')$ where the neutral kaon decays inside the chamber to two pions, $K^0_S \to \pi^+\pi^-$ [24]. Rescattering in the Xe

[7] The decay of the neutral kaon K^0 does not allow to determine the sign of the strangeness, since both the neutral kaon K^0 and antikaon $\overline{K}^0$ decay to a pion pair $\pi^+\pi^-$.



Figure 2.4: The missing mass distribution of $MM(\gamma K^- p)$ on the deuterium target. A peak structure at $1.53\,$GeV and a broad bump at $1.6\,$GeV are observed above the background (indicated by the red lines). The values on the horizontal axis are in GeV, with unspecified bin size. For a description of the background model refer to the text. Taken from references [22, 23].

nucleus was suppressed by requiring the proton p and neutral kaon K^0 to be in the forward direction and in opposite azimuthal regions with respect to the incoming kaon K^+. A narrow resonance was found at a mass of $1539 \pm 2\,$MeV with a width smaller than 9 MeV, visible in the left panel of figure 2.5. The naive statistical significance was determined to be $4.4\,\sigma$.

Full data sample

In 2006 the DIANA collaboration finished the analysis of the full data sample [25]. The new results confirm the earlier results (see right panel in figure 2.5), and increase the naive statistical significance of the observed $\Theta(1540)$ peak. The intrinsic width of the $\Theta(1540)$ resonance peak could be determined from the ratio of resonant and non-resonant events as $\Gamma = 0.36 \pm 0.11\,$MeV.

The upper limits on the intrinsic width Γ of an exotic baryon, as determined from the re-analysis of the NK scattering data, is consistent with the width determined at the DIANA experiment.

2.1.5 Photoproduction on protons to $nK^+K^0_S$ at the SAPHIR experiment

At the SAPHIR experiment [26], similar to the CLAS experiment below, energetic photons are produced by bremsstrahlung when 2.8 GeV electrons from the continuous mode Electron Stretcher Accelerator (ELSA) strike a copper foil radiator. The energies of the scattered electrons are measured in the tagging system, and allow the determination of the photon energies between 0.9 GeV and 2.7 GeV. The photon beam then passes through a liquid hydrogen target.

The hydrogen target is located in the center of a cylindrical drift chamber. The momentum of charged particles is determined from their motion in the field of a C-shaped magnet. The central drift chamber is surrounded by a scintillator wall to measure the time-of-flight. In the forward direction a planar drift chamber is used for tracking.



Figure 2.5: The distributions of the invariant mass $M(pK^0)$ for the charge-exchange reaction $K^+n(\mathrm{Xe}) \to pK^0\mathrm{Xe}$' at the DIANA experiment. Rescattering in the Xe nucleus is suppressed. The original data sample (left panel) was expanded by a factor two and re-analyzed (right panel). The distribution of the background events is obtained from a fit to the data. Taken from references [24, 25].

In the reaction $\gamma p \to \Theta^+ K^0_S \to nK^+\pi^+\pi^-$ the kinematic parameters of the neutron were determined from missing energy and momentum. In the distribution of the invariant mass $M(nK^+)$ a peak is observed at 1540 ± 4(stat) $\pm$ 2(sys) MeV with a width smaller than 35 MeV. The cross section is estimated to be similar to that of ϕ photoproduction, but a factor 4 smaller than final states with open strangeness [27], or around $0.2\,\mu$b [28].

Recent null results in the same reaction channel by the CLAS experiment [29, 30], with a very similar geometrical acceptance and a statistical precision far exceeding that of the SAPHIR results, have cast a shadow of doubt on the result of the SAPHIR experiment. The upper limit for this reaction channel determined at the CLAS experiment is 0.8 nb, almost three orders of magnitude smaller than the value determined at the SAPHIR experiment.

2.1.6 Photoproduction on protons and deuterons at the CLAS experiment

At the CEBAF Large Acceptance Spectrometer (CLAS) experiment on the 6 GeV CEBAF race-track accelerator at the Thomas Jefferson National Accelerator Facility (JLab) high-energetic electrons or tagged photons are scattered on fixed, light nuclear targets [31]. The tagged photons are produced by colliding the accelerated electron beam with an energy of up to 6 GeV on a thin bremsstrahlung radiator. The tagged photons have energies from 3 to 5.5 GeV.

The detector is, as indicated by its name, a large acceptance device with a toroidal magnetic field and drift chambers for particle tracking, and Čerenkov counters and scintillation time-of-flight detectors for particle identification. Electromagnetic calorimeters are used to detect electrons, photons, and neutrons (although neutron detection is not used in the analyses described here).

The CLAS experiment has published several results on exotic baryons, all in photoproduction reactions on light nuclei. The first two positive results contributed largely to the initial acceptance of the $\Theta(1540)$ baryon, whereas the later null results were considered by some as the proverbial 'nail in the coffin' of the $\Theta(1540)$ baryon.

Photoproduction on deuterons (g2a and g10)

The CLAS experiment obtained its first exotic baryon results with the reaction $\gamma d \to K^+K^-p(n)$ [32]. In this reaction on the neutron, the proton can be regarded as a spectator. It has only a small transverse momentum and is normally not detected by the CLAS spectrometer. Relying on the spectator model for the determination of the neutron momentum is an approximation that increases the number of background events. Therefore, only those events were used where the proton and the K^- have a final state interaction (FSI) and the proton is scattered into the acceptance of the detector by the K^-. This technique allowed for a reduction of the background.



Figure 2.6: The results of the g10 run at the CLAS experiment. The data points collected during the g2a run are compared to the distribution obtained in the g10 run, represented by the solid line (left panel). In the search for associated production of the exotic baryon $\Theta(1540)$ and a Λ hyperon, no resonance could be observed (right panel). Taken from references [33, 34].

The kinematic parameters of the neutron were reconstructed from the missing energy and momentum of the reaction. In the distribution of the invariant mass $M(nK^+)$ a resonance with a naive significance of $5.2\,\sigma$ was found at $1542 \pm 5\,\mathrm{MeV}$, with a width compatible with the experimental resolution. The background was modeled with a Gaussian function.

A repetition of this measurement at the CLAS experiment, with more than six times higher statistical precision and in very similar conditions,[8] could not confirm the existence of a resonance peak in this channel. The background had been underestimated substantially in the first result. In the left panel of figure 2.6 the data points of the original g2a result are compared to the newer results, represented by the solid line. The upper limit of the cross section in the channel $\gamma n \to K^-\Theta^+$ was estimated at 3 nb [33].

In the newer data sample, a search for exotic resonances was also performed in the distribution of the invariant mass $M(K^+n)$ in the reaction $\gamma d \to \Lambda K^+(n)$. The requirement of a Λ hyperon in the final state explicitly tags the strangeness of the reaction. As shown in the right panel of figure 2.6, no resonance was observed and an upper limit on the cross section of 5 nb was determined.

To summarize the results of the CLAS experiment on a deuterium target, we conclude that no exotic resonances could be observed in different reaction channels, and strict upper limits on the production were determined.

Photoproduction on the proton (g6)

A second analysis investigated the reaction channel $\gamma p \to \pi^+K^-K^+(n)$ on a proton target [35]. Again, the neutron was reconstructed from the missing energy and momentum, and in the distribution of the invariant mass $M(nK^+)$ a resonance peak at a mass of $1555 \pm 10\,\mathrm{MeV}$ with a naive significance $7.8\,\sigma$ was found, as shown in figure 2.7. The inset of figure 2.7 was obtained with relaxed event selection requirements. The experimental resolution was also here dominant over the width of the resonance. No resonance structure corresponding to the decay $\Theta^{++} \to pK^+$ was found.

A repetition of this positive experiment is scheduled for the first half of 2008. The g12 run will collect

[8] One substantial difference was the movement of the target 25 cm upstream of the center of the spectrometer. For one half of the data collection, a lower value for the magnetic field of the CLAS spectrometer was chosen. This was done to increase the acceptance for forward angle, negatively charged tracks. For the comparison with the previous results, only the half of the data set taken at the same magnetic field value was used.



Figure 2.7: The distributions of the invariant mass $M(nK^+)$ in the reaction $\gamma p \to \pi^+ K^- K^+ (n)$ for data collected at the CLAS experiment. The neutron was reconstructed from the missing energy and momentum. The inset shows the distribution with relaxed event selection requirements. Taken from reference [35].

a higher number of photoproduction events on protons, and should either confirm this result with high precision or refute it with a strict upper limit on the allowed cross section.

Photoproduction on the proton with increased statistics (g11)

In 2005, new results on a proton target were presented by the CLAS experiment, once again with very high statistical precision. In the reaction channels $\gamma p \to \overline{K}^0 K^0 p$ [29] and $\gamma p \to \overline{K}^0 K^+(n)$ [30] a search was performed for a resonance with decay channels $\Theta^+ \to K^0_S p$ and $\Theta^+ \to K^+ n$, but no resonance could be observed. The results are reproduced in figure 2.8. A combined analysis of both channels allows to estimate an upper limit on the production cross section $\gamma p \to \overline{K}^0 \Theta^+$ of 0.7 nb. Using several theoretical models, the width of a possible resonance is limited to values below 3.2 MeV.

Due to the implemented hardware triggers and the lower photon energy, this data sample can not be used to confirm or refute the earlier, positive result on a proton target in the reaction channel $\gamma p \to \pi^+ K^- K^+ (n)$.

The previous results seem to be in disagreement with the results of the LEPS experiment. A detailed analysis, taking into account the small differences in acceptance and kinematic coverage, indicates that the results are still consistent for several possible theoretical scenarios for the exotic baryons [36].

It is possible that seemingly unrelated changes to the analysis method or selection criteria reduce the number of observed exotic events, due to the interference with another process. A recent development in the search for exotic baryons at the CLAS experiment exploits the interference between different reactions with identical final states [37, 38].

2.1.7 Quasi-real photoproduction at the HERMES experiment

The HERMES experiment will be described in detail in chapter III. Here we summarize the published evidence for a narrow resonance compatible with the exotic baryon Θ^+ in quasi-real photoproduction on a deuterium target through the decay channel $pK^0_S \to p\pi^+\pi^-$ [3].

The distribution of background events is modeled using a simulation with the PYTHIA Monte Carlo generator, which does not include several excited Λ^* hyperon resonances that thus have to be added in by hand. The shape of the mixed event background distribution agrees with the non-resonant contribution of the Monte Carlo simulation, providing further justification of this method.[9] In the left panel of figure 2.9 the distribution of the invariant mass $M(pK^0_S)$ is shown with a fit of a Gaussian resonance shape on a



Figure 2.8: The distributions of the invariant mass $M(K_S^0 p)$ (left, top) and $M(K^+n)$ (right) in the reaction channels $\gamma p \to \overline{K}^0 K^+(n)$ and $\gamma p \to \overline{K}^0 K^0 p$ at the CLAS experiment. The missing mass of the second reaction, when only a K_S^0 is required in the final state, is shown in the left, bottom panel. No resonances are visible, and upper limits on the photoproduction cross section are estimated. Taken from references [29, 30].



Figure 2.9: Observation of the exotic baryon Θ^+ at the HERMES experiment. In the left panel the background events in the distribution of the invariant mass $M(pK^0_S)$ are modeled using the mixed event background distribution (solid black line) normalized to the non-resonant contribution determined in a PYTHIA Monte Carlo simulation (shaded histogram). In addition the excited Λ^* hyperon resonances (dotted lines) and a Gaussian resonance shape corresponding to the exotic baryon Θ^+ (dashed line) are added. In the right panel the sum of third order polynomial function and a Gaussian resonance shape is used. Taken from reference [3].

background model composed of the Monte Carlo simulation, mixed event distribution, and additional Λ^* hyperon resonances. In the right panel of figure 2.9 the data is fitted with the sum of a third order polynomial function and a Gaussian resonance shape, effectively reducing the assumptions on the understanding of the distribution of background events.

A search for the exotic antibaryon Θ^- was also performed at the HERMES experiment, but did not result in an observation. In figure 2.10 the distribution of the invariant mass $M(\overline{p}K^0_S)$ is shown. The number of events was determined using the sum of a third order polynomial function and a Gaussian resonance shape. The result of 3 ± 6 events is consistent with zero.

2.1.8 Fixed-target proton-proton collisions with the COSY-TOF experiment

The COSY-TOF experiment is a time-of-flight detector at the cooler synchrotron proton storage ring COSY at FZ-Jülich. Protons of 2.95 GeV are focused on a liquid hydrogen target and the reaction products are detected in several time-of-flight hodoscopes. The momentum is reconstructed geometrically, without the use of a spectrometer magnet. The investigation of the predictions of the chiral quark soliton model [17] was already in 1998 one of the objectives of the COSY-TOF experiment.

In the exclusive reaction channel $pp \rightarrow \Sigma^+ K^0_S p$, a narrow resonance was observed in the distribution of the invariant mass $M(pK^0_S)$ at 1530 ± 5 MeV, with a width compatible with the experimental resolution [39]. The cross section was estimated as $0.4 \pm 0.1(\mathrm{stat}) \pm 0.1(\mathrm{sys})\,\mu$b.

More data was collected by the COSY-TOF experiment at marginally higher beam energy. Using three different analysis methods, no $\Theta(1540)$ resonance could be observed in the new data with high statistical precision. An upper limit on the cross section of $\sigma(pp \rightarrow \Sigma^+\Theta^+) < 0.15\,\mu$b was determined (at 95% confidence level) [40].

[9]The event mixing method that was used for this analysis is known to be incorrect. For a detailed discussion, refer to section 4.7.



Figure 2.10: Search for the exotic baryon Θ^- at the HERMES experiment. The distribution of the invariant mass $M(\overline{p}K^0_S)$ is fitted with the sum of a third order polynomial function and a Gaussian resonance shape.

2.1.9 Proton-proton collisions at the NA49 experiment

At the NA49 experiment at the CERN SPS accelerator complex protons with an energy of 158 GeV were collided on a liquid hydrogen target. The reaction fragments were detected with four large time projection chambers. Particle identification is achieved using the energy loss dE/dx.

By reconstructing the particles in the decay channel to $\Xi^-\pi^-$ and further to $\Lambda\pi^-\pi^-$, the invariant mass distribution in figure 2.11 is obtained. After subtracting the background determined by event mixing, a narrow peak is visible at a mass of 1862 MeV. This resonance was interpreted as the exotic baryon Ξ^{--} [41].

2.1.10 Inclusive deep-inelastic scattering at the ZEUS experiment

The ZEUS experiment was the first collider experiment to provide evidence for the existence of the $\Theta(1540)$ state. At the ZEUS experiment inclusive deep-inelastic scattering of electrons and protons at a center-of-mass energy of about 310 MeV is studied. In the analysis at the ZEUS experiment resonance decays to $K^0_S\overline{p}$ are considered as well. They can only be explained as an exotic baryon $\overline{\Theta}$ with minimal quark content $(\overline{uudds})$. A resonance at 1521.5 ± 1.5(stat) $\pm$ 2.8(sys) MeV is observed. The width is consistent with the experimental resolution of 2 MeV [42].

2.1.11 Hadronic interactions at the SVD-2 experiment

At the SVD-2 experiment, the 70 GeV proton beam of the IHEP accelerator in Protvino, Russia, collides on a fixed C, Si, or Pb target. The spectrometer was designed to study charm hadroproduction near threshold. It consists of a high-precision microstrip vertex detector, a large-aperture spectrometer magnet with multi-wire proportional chambers, and a threshold Čerenkov counter for separation of pion and proton tracks between 4 and 21 GeV. A narrow resolution for several established resonances could be demonstrated.

The search for exotic baryons at the SVD-2 experiment concentrated on the inclusive scattering of protons from nuclei $pA \to pK^0_S X$. Two results have been presented by the SVD collaboration, with partly overlapping data samples. The first result presents a resonance in the distribution of the invariant mass $M(pK^0_S)$ at 1526 ± 3(sys) $\pm$ 3(stat) MeV [43]. One could argue that the resonance seems to exist of only a single data point.



Figure 2.11: Evidence for the exotic baryon Ξ^{--} at the NA49 experiment is found in the decay channel to $\Xi^{--} \rightarrow \Xi^-\pi^- \rightarrow \Lambda\pi^-\pi^-$. The normalized mixed event background (shaded histogram) is subtracted from the invariant mass distribution in the top panel. A narrow peak is visible at a mass of 1862 MeV in the subtracted distribution in the bottom panel. Taken from reference [41].



Figure 2.12: The distribution of the invariant mass $M(pK^0_S)$ for inclusive DIS events at the ZEUS experiment. A clear peak is visible at 1521.5 MeV.

A more recent analysis, with improved tracking algorithms, confirmed the earlier indications. An increase of a factor 3 to 4 in total number of K_S^0 events was obtained. Moreover, the full data sample was separated in two distinct kinematic regimes, depending on the location of the K_S^0 decay vertex. Even without event selection criteria, both subsamples exhibit a peak at 1522.2 ± 3(stat) MeV or 1523.6 ± 3.1 MeV (see figure 2.13). The total proton-nuclear cross section is estimated as 6 μb.

2.1.12 Secondary kaons in the BELLE detector

Using the interaction of secondary kaons with the detector material of the BELLE experiment at the KEKB e^+e^- collider, production of the exotic baryon $\Theta(1540)$ was studied [44]. Interactions in the inner Si vertex detectors were used, and both inclusive and exclusive reaction channels were investigated. The result of the inclusive search for pK^0 decays are shown in the left panel of figure 2.14. No structure is observed in the distribution of the invariant mass $M(pK_S^0)$, but the $\Lambda(1520)$ hyperon is clearly visible in the distribution of the invariant mass $M(pK^-)$. The estimated upper limit on the ratios of the inclusive $\Lambda(1520)$ and $\Theta(1540)$ cross sections is 2.5%.

Exclusive production of $\Theta(1540)$ baryons is studied in the reaction $K^+n \to pK_S^0$. Since the (secondary kaon) projectile is not reconstructed, assumptions have to be made about the contribution of other reaction channels to the same final state. Destructive interference between different reaction channels is expected to be negligible, though. The resulting distribution of $M(pK_S^0)$ is shown in the right panel of figure 2.14, but does not indicate any structure at the position of the reported $\Theta(1540)$ baryon. The expected yield is indicated by the open points, and the signal expected from the results of the DIANA experiment are indicated by the solid line. These results do not seem to confirm the results of the DIANA experiment. The upper limit on the width of the $\Theta(1540)$ resonance is estimated as $\Gamma < 0.64$ MeV.

2.1.13 Exotic baryons in e^+e^- annihilation at the BABAR experiment

At the BABAR experiment electrons and positrons are collided at a center of mass energy of $\sqrt{s} = 10.58$ GeV. An inclusive search for the $\Theta(1540)$ and for most of the other members of the exotic antidecuplet was performed on the very expansive data set with an integrated luminosity of 123 pb^{-1}. Clear signals are observed for the known baryon resonances, but no evidence is found for the production of exotic baryons.

In the left panel of figure 2.15 the distribution of the invariant mass $M(pK_S^0)$ is shown for events satisfying the decay channel $\Theta(1540) \to pK_S^0$. No $\Theta(1540)$ resonance is observed at a mass of 1540 MeV or anywhere else in the distribution. Similarly, the existence of the exotic $\Xi^{--}(1860)$, $\Xi^-(1860)$ and $\Xi^0(1860)$ resonances could not be confirmed.

For the manifestly exotic baryons $\Theta^+(1540)$ and $\Xi^{--}(1860)$ upper limits on the production rate per $e^+e^- \to q\overline{q}$ interaction are determined to allow comparison with the regular baryons [45]. The comparison between regular baryons (at two different center-of-mass energies) and the determined upper limits on the production rate for the manifestly exotic baryons (for two assumptions on the width) is shown in the right panel of figure 2.15.

Although baryon production in e^+e^- collisions is known for the democratic production of hadrons with nonzero strangeness (and other heavier quark constituent), depending only on the mass and the spin of the produced hadron but not directly on the specific quark content, the production of an exotic baryon such as the $\Theta^+(1540)$ requires the creation of five $q\overline{q}$ pairs out of the vacuum compared to only three for regular baryons. Comparing the production rates of exotic hadrons to the expectations for regular hadrons could therefore appear unmotivated.

More recently an inclusive search for the $\Theta^+(1540)$ was performed using the interactions of secondary hadrons and of electrons or positrons in the beam halo with the beam pipe and the material of the inner tracking detectors [46, 47]. Again no evidence for the exotic baryon $\Theta^+(1540)$ was found, but the results allow for a comparison with other photo-, electro-, and hadroproduction (see figure 2.16).



Figure 2.13: Recent results of the SVD-2 experiment. Both figures were obtained with statistically independent subsamples.



Figure 2.14: Secondary kaon scattering with the Si vertex detectors at the BELLE experiment is used to study the reaction $K^+n \to pK^0$. In the left panel, the distributions of the invariant mass $M(pK^-)$, represented by the data points with error bars with a prominent $\Lambda(1520)$ resonance peak, and the invariant mass $M(pK^0)$, represented by the solid histogram, are shown for inclusive events. In the right panel, the distribution of the invariant mass $M(pK^0)$ for exclusive events is shown, with the expected yield indicated by the open symbols. Taken from reference [44].



Figure 2.15: Production of exotic baryons in e^+e^- annihilation at the BABAR experiment. In the left panel the distribution of the invariant mass $M(pK_S^0)$ is shown. The expected position of the $\Theta^+(1540)$ peak is indicated but no structure is visible. In the right panel the production rates per $e^+e^- \rightarrow q\overline{q}$ interaction are compared for the regular baryons and for the determined upper limits on the production of the exotic baryons $\Theta^+(1540)$ and $\Xi^{--}(1860)$. Taken from reference [45].



Figure 2.16: Electro- and hadroproduction of exotic baryons at the BABAR experiment. The results of the BABAR experiment are compared with the results of the HERMES experiment (left panel) and the ZEUS experiment (right panel). Taken from reference [47].



Figure 2.17: The search for $\Theta^+(1540)$ decays at the HyperCP experiment. The expected resonance shape is shown in the upper panel, and used in a fit to the distribution of the invariant mass $M(pK^0_S)$ in the lower panel. At most 0.3% of the pK^0_S candidates could come from an exotic decay $\Theta^+(1540) \to pK^0_S$. Taken from reference [48].

2.1.14 Fixed target proton-tungsten collisions at the HyperCP experiment

The HyperCP experiment at Fermilab was designed to study CP symmetry violation in the decays of the cascade hyperons Ξ^- and Ξ^+ [49]. A secondary beam is produced by 800 GeV protons striking a copper target, and the collimator channel immediately downstream of the target and embedded in a dipole magnet is used to select a positively charged beam, consisting of mainly protons and pions with approximately 5% kaons, with a broad momentum distribution between 120 and 220 GeV. Events produced in the tungsten exit region of the collimator were used in a search for the exotic baryons $\Theta^+(1540)$ [48].

The HyperCP experiment does not have dedicated particle identification detectors. For the identification of the K^0_S mesons the signature of their decay to two oppositely charged pions with a reconstructed invariant mass $M(\pi^+\pi^-)$ consistent with the K^0 mass allows for a clean selection. Charged particles with more than 50% of the total momentum were assumed to be protons, motivated by the high probability of being protons for these events. Thus, the search was restricted to the exotic baryon decay $\Theta^+(1540) \to pK^0_S$.

In the distribution of the invariant mass $M(pK^0_S)$ no peak was observed, as shown in figure 2.17. The distribution is fitted with the expected resonance shape obtained in a Monte Carlo simulation. At most 0.3% of the pK^0_S candidate events could come from the decays of exotic Θ^+ baryons. The same data sample contained over 140,000 combined Ξ^- and Ξ^+ decays.

The high value and broad spread of beam momentum avoids the possibility of peaks generated by kinematic reflections. Unfortunately, together with the mixed beam composition it precludes the estimation of limits on the production cross section of the $\Theta(1540)$ baryons.

2.1.15 High-energetic proton-nucleus interactions at the HERA-B experiment

At the HERA-B experiment collisions of high-energetic protons in the halo of the 920 GeV HERA proton beam with the nuclei of atoms in target wires were studied. The HERA-B detectors has a large acceptance, precision silicon vertex detectors and particle identification detectors. The resonance decays $K^0_S \to \pi^+\pi^-$, $\Lambda \to p\pi^-$, $\Lambda(1520) \to pK^-$, and $\Xi^- \to \Lambda\pi^-$ are cleanly identified and reconstructed. In the selected momentum region between 22 and 55 GeV the misidentification of protons is less than 1%.

Using the decay channel to pK^0_S an inclusive search for the exotic baryon $\Theta^+(1540)$ was performed in the data samples collected on carbon, titanium and tungsten wires. The three obtained invariant mass distributions are shown in top panels of figure 2.18. Any structures in the distributions are consistent with statistical fluctuations. The upper limits on the cross section for the production of $\Theta(1540)$ baryons are



Figure 2.18: The search for Θ^+ and Ξ^{--} in pA reactions at the HERA-B experiment. For all three investigated target materials, carbon (left panel), titanium (middle panel), and tungsten (right panel), no statistically significant peak for the exotic baryon Θ^+ is invisible in the invariant mass distribution $M(pK^0_S)$. In the bottom panels the featureless distribution of the invariant masses $M(\Xi^-\pi^+)$ (left panel) and $M(\Xi^-\pi^-)$ (right panel) and charge conjugate modes are shown. Taken from references [50, 51].

between 4 and 20 μb per nucleon. The ratio of the production rates of the exotic Θ^+ to other hyperons is five times smaller than the value observed at the ZEUS experiment, and two orders of magnitude smaller than the value observed at the HERMES experiment.

A search for the exotic baryon $\Xi(1860)$ was performed in the decay channels to $\Xi^-\pi^-$ and $\Xi^-\pi^+$ and the corresponding antiparticle channels, but also there upper limits stronger than 4 μb per nucleon could be determined. The distributions of the invariant mass $M(\Xi\pi)$ for the $\Xi^0(1860)$ and $\Xi^{--}(1860)$ combinations (and their charge conjugates) are shown in the bottom panels of figure 2.18. A clear excess is visible at 1530 MeV corresponding to the well-established $\Xi^{0*}(1530)$ hyperon.

2.1.16 Discussion

In this experimental overview several observations of the exotic baryons Θ^+ and Ξ^{--} have been discussed. Among the original observations of the Θ^+ there are several that are not valid anymore (SAPHIR, CLAS on deuterons, COSY-TOF), and only the LEPS and SVD-2 experiments have been able to confirm the original analysis with additional data, although these results have only been shown at conferences or published in proceedings. The observation of the exotic baryon Ξ^{--} has not been confirmed by any experiment. For the exotic baryon Θ^+ a long list of experiments with null results, often associated with upper limits on the branching ratios or cross sections, is shown in table 2.1.

A small number of experiments is or will be collecting data in very similar conditions as for their earlier observation. With this new data they are expected to confirm or refute the earlier results. At the ZEUS experiment the original observation of the exotic baryon Θ^+ only involved data collected during the HERA I running period until the year 2000. Since the luminosity upgrade and the installation of an improved vertex detector in 2001, the ZEUS experiment has collected additional data in the HERA II running period until the year 2007. This data is still being analyzed. The CLAS experiment will collect

Table 2.1: A large number of experiments is unable to confirm the existence of the exotic baryon Θ^+. These null results are listed here alongside the limits they impose on the branching ratio (B.R.) or relative yield of other resonances.

Group	Reaction	Limit
BES [52]	$e^+e^- \to J/\Psi \to \overline{\Theta}\Theta$	$< 1.1 \times 10^{-5}$ B.R.
BaBar [45, 53]	$e^+e^- \to \Upsilon(4S) \to pK^0X$	$< 1.0 \times 10^{-4}$ B.R.
Belle [44]	$e^+e^- \to B^0\overline{B}^0 \to p\overline{p}K^0X$	$< 2.3 \times 10^{-7}$ B.R.
LEP [54]	$e^+e^- \to Z \to pK^0X$	$< 6.2 \times 10^{-4}$ B.R.
HERA-B [55]	$pA \to K^0pX$	$< 0.02 \times \Lambda^*$
SPHINX [56]	$pC \to K^0\Theta^+X$	$< 0.1 \times \Lambda^*$
HyperCP [48]	$pCu \to K^0pX$	$< 0.3\%\ K^0p$
CDF [57]	$p\overline{p} \to K^0pX$	$< 0.03 \times \Lambda^*$
FOCUS [58]	$\gamma BeO \to K^0pX$	$< 0.02 \times \Sigma^*$
Belle [44]	$\pi + Si \to K^0pX$	$< 0.02 \times \Lambda^*$
PHENIX [59]	$Au + Au \to K^-\overline{n}X$	(not given)

new data on a proton target during the first half of 2008. This will allow for the refutation or confirmation of the observation of the exotic baryon Θ^+ in the original data set collected during the g6 run [35].

In table 2.2 several observations are listed alongside with a repetition of the measurement with higher statistical precision. This list is not intended to be complete, and it is certainly debatable whether some can be considered a repetition, but it indicates that several of the original experiments with only a small and unconvincing number of events have not stood up to the scruntiny of experiments with a very high statistical precision. In the few cases where the original observation was confirmed, the precision of the repeated result is still not convincing.

2.2 Theoretical models

In this section the theoretical models for hadrons and more specifically exotic baryons are discussed. Since a perturbative approach to QCD is impossible at the low energies of hadronic bound states, several phenomenological models have been proposed in the past. The concept of constituent quarks is important in all of these models. A quite different approach was taken by the chiral quark soliton model (χQSM), which provided the predictions of a narrow exotic baryons that lead to the recent experimental search for exotic baryons [17].

Several theoretical reviews on exotic baryons have been published before [60, 61, 62], but the most important theoretical approaches are summarized in this section.

2.2.1 The MIT bag model

A conceptually very simple model for hadrons is the MIT bag model [63, 64]. The massless (or very light) quarks and gluons are confined in a finite region with constant energy density, the ‘bag’. Inside the bag the quarks do not interact with each other, leading to the property of asymptotic freedom. Outside the bag the masses of the quarks become infinitely large, and thus also the color confinement property of QCD is trivially satisfied. The content of the bag can always be chosen as a color-singlet.

This simple model with a very limited number of parameters is surprisingly successful in the description of hadrons, reproducing not only the masses of the light baryons but also the proton radius and even quark distributions in the nucleon. The bag model can be readily extended to include five or more quarks. For the mass of the lightest exotic baryons values from 1700 to 2400 MeV were predicted [65].

Table 2.2: Comparison of observations of the exotic baryon Θ^+ with repeat experiment. The original positive results, claiming the observation of the exotic baryon $\Theta^+(1540)$, are compared to repeat experiments in very similar kinematic conditions but with higher statistical precision. Taken from reference [66].

Observation	Reaction	Repetition	Reaction	Increase	Result
LEPS [20]	$\gamma C \to K^+K^-(n)$	LEPS [22, 23]	$\gamma d \to K^+K^-(n)$	$\times 5$	Confirmed $5\,\sigma$
SVD-2 [43]	$pA \to pK^0_S X$	SVD-2 [67]	$pA \to pK^0_S X$	$\times$3–4	Confirmed 6–9 σ
DIANA [24]	$K^+\text{Xe} \to K^0 pX$	Belle [44]	$K^+\text{Si} \to K^0 pX$	$\times 10$	$\Gamma_\Theta < 1\,\text{MeV}$
SAPHIR [27]	$\gamma p \to K^0 K^+ n$	CLAS (g11) [29]	$\gamma p \to K^0 K^+ n$	$\times 10$	$\sigma_{tot} < 1\,\text{nb}$
CLAS (g2a) [32]	$\gamma d \to K^+K^-p(n)$	CLAS (g10) [33]	$\gamma d \to K^+K^-p(n)$	$\times 6$	$\sigma_{tot} < 3\,\text{nb}$
HERMES [3]	$e^+d \to K^0 pX$	BABAR [47]	$e^+\text{Be} \to K^0 pX$	$\times 200$	No Θ^+ observed
COSY-TOF [39]	$pp \to \Sigma^- K^0 p$	COSY-TOF [40]	$pp \to \Sigma^- K^0 p$	$\times$4–8	$\sigma < 0.15\,\mu\text{b}$

2.2.2 Group theory of exotic baryons

To determine the configuration of exotic baryons states, we start from similar group theoretical arguments that were used to explain the regular meson and baryon multiplets in chapter I. In the expansion of $\mathbf{6} \otimes \mathbf{6} \otimes \mathbf{6} \otimes \mathbf{6} \otimes \overline{\mathbf{6}}$ the exotic baryons end up in antidecuplets, 27-plets, and 35-plets [68, 69, 70].

When we limit ourselves to states without angular momentum ($\ell = 0$), the exotic baryons with the lowest energy are organized in the antidecuplet ($\overline{\mathbf{10}}$) with spin and parity $J^\pi = \frac{1}{2}^+$. It is nearly degenerate with an octet ($\overline{\mathbf{8}}$) of states with negative parity, but in the octet no manifestly exotic states are present. The antidecuplet $\overline{\mathbf{10}}$ is shown in figure 2.19, where the masses are predictions of the chiral quark soliton model. To determine numerical values for the masses and widths of the exotic baryons, specific models have to be used.

2.2.3 The chiral structure of QCD

At low energies the chiral structure of QCD defines the dynamic behaviour of quarks, mesons and baryons [71, 72]. Because the u and d quarks are almost massless compared to the nucleon, the theory of QCD with only two flavors is invariant under transformation in the chiral symmetry group $SU(2)_R \times SU(2)_L$.[10] Since we do not observe pairs of hadrons with opposite parity, this chiral symmetry must be spontaneously broken down to the symmetry group $SU(2)_V$, where V refers to the invariant vector charge of the QCD ground state or vacuum. The three Goldstone bosons associated with the symmetry breaking are the pions π^-, π^0 and π^+. They are light but not massless due to the small but non-zero masses of the u and d quarks. When we consider the s quark in addition to the u and d quarks, the same mechanism of spontaneous chiral symmetry breaking occurs, but now eight massive Goldstone bosons are formed (π^-, π^0, π^+, K^-, K^0, $\overline{K}^0$, K^+, η). The η' is substantially heavier than the η because it is associated with the broken $U(1)$ axial symmetry.

The spontaneous symmetry breaking is accompanied by the formation of the gluon condensate $G_{\mu\nu}G^{\mu\nu}$ and the chiral quark condensate $\langle q\overline{q}\rangle$, which fills the vacuum with quark-antiquark pairs. The light 'bare' quarks and antiquarks that compose the mesons and baryons are 'dressed' in these fields and receive a momentum-dependent dynamical mass ($m_{u,d} \approx 350\,\text{MeV}$, $m_s \approx 470\,\text{MeV}$). These so-called constituent quarks have now masses approximately one half of the mass of the ρ meson or one third of the mass of the nucleon.

Baryons can now be described as three constituent quarks surrounded by a cloud of massless mesons. The interactions between the constituent quarks and the meson cloud determine the properties of the baryon.

2.2.4 The chiral quark soliton model

In the 1960s a different approach was proposed by Skyrme to study the nucleon [73, 74] and, by extension to $SU(3)_f$, baryons with strangeness [75, 76]. At low energies the baryons are regarded as spherically symmetric soliton solutions of the pion field or chiral field that forms due to the spontaneous breaking of the chiral symmetry of QCD.

This suggests a classification of the light baryons as rotational states. The minimal generalization of spherical symmetry in a system with three isospin components (*i.e.* the three pions) is

$$\pi^\alpha(x) = \frac{x^\alpha}{r} P(r),$$

where $P(r)$ is the spherically symmetric profile of the soliton. This implies that a space rotation of the field is equivalent to a rotation in isospace. From the analogy of this situation with the classical spherical top, one sees that the rotational states are $(2J+1)^2$-fold degenerate in spin and isospin. Applying this for $J = \frac{1}{2}$ we find the four nucleon states. For $J = \frac{3}{2}$ we find the sixteen Δ resonance states. The relations between the characteristics of the states are generally within about 30% of the experimental values [77].

[10] For massless particles the helicity h, defined as the projection of the spin σ on the direction of the momentum $\sigma \cdot \frac{\vec{p}}{|\vec{p}|}$, is an invariant. It is then also referred to as the chirality or handedness. It can take the value $+1$ for right-handed or -1 for left-handed particles.



Figure 2.19: The ground state antidecuplet of the exotic $J=\frac{1}{2}$ baryon states, with the masses predicted by the chiral quark soliton model.

If extended to three quark flavors, the rotations are performed in ordinary space and in $SU(3)$ space. Quantization shows that the lowest baryon state is the octet of regular baryons with spin $J=\frac{1}{2}$ and the second excitation is the decuplet of regular baryons with spin $J=\frac{3}{2}$. The relations between the mass, width and branching ratio of the members of the octet and decuplet are satisfied to an accuracy of up to one percent [17].

Already in 1987 the existence of the lightest exotic baryon with strangeness $S=+1$ at a mass of approximately 1530 MeV was predicted [78, 79].

The third rotational excitation in the $SU(3)$ case is an antidecuplet with spin $J=\frac{1}{2}$ (see figure 2.19). This exotic antidecuplet can be alternatively considered in a primitive way as states made of three quarks plus a quark-antiquark pair. The lightest member Θ^+ can then be considered a K^+n or K^0p bound state. The advantage of the chiral soliton model is that all concrete numbers follow from symmetry considerations and do not rely on a specific dynamical realization. If one number is extracted from experiment, one can completely fix all the other members of the antidecuplet, together with their widths and branching ratios.

By using the mass of the established neutron resonance N at 1710 MeV as the 'anchor' for the antidecuplet, the mass of the Θ^+ was predicted to be 1530 MeV and the width less than 15 MeV.[11] These predictions partially motivated the experimental searches for Θ^+ and are in agreement with the results of the different experiments.

However, in this model the predicted width of $N_{\overline{10}}$ and $\Sigma_{\overline{10}}$ are both below the experimental values listed by the Particle Data Group. 'Anchoring' the antidecuplet to these states increases the width of the Θ^+ to exceed the bounds set by experimental data [80].

2.2.5 The Jaffe-Wilczek diquark model

In an alternative positive-parity interpretation the exotic baryon Θ^+ is a bound state of two highly correlated ud diquarks and an $\overline{s}$ antiquark [81]. The quarks in the ud diquarks are necessarily in an antisymmetric color configuration $\overline{\mathbf{3}}_c$, otherwise no color singlet would be formed with the remaining antiquark. When the quarks have opposite spins, then they are simultaneously in an antisymmetric color, flavor, isospin, and spin configuration. A strong correlation between the quarks in the diquarks is then expected.

While this models predicts the same exotic antidecuplet of states as the chiral quark soliton model in addition to a crypto-exotic octet, there are distinct differences in the masses of the states. The exotic antidecuplet and crypto-exotic octet of baryons is shown in the left panel, and the relative mass spectrum in the right panel of figure 2.20. Due to the mixing of the antidecuplet and the octet, one crypto-exotic state with a mass below that of the Θ^+ is formed. This state was identified as the Roper resonance $N(1440)$ in this model.

[11] After the first observations of the exotic baryon Θ^+ a numerical mistake in the calculation of the width was discovered. The correct value of the width following from the chiral quark soliton model is therefore closer to 30 MeV. Refer to references [82, 83, 84] in that order for a discussion of this issue.



Figure 2.20: In the diquark model exotic baryons consist of two highly correlated diquarks and an anti-quark. In the left panel the predicted $J=\frac{1}{2}$ antidecuplet and $J=\frac{3}{2}$ are shown. When the mass of the Θ^+ baryon is fixed to the value in the χQSM, the relative mass spectrum with the label "quark" in the right panel is obtained. Taken from reference [81].

The diquark model predicts that the exotic baryon Ξ^{--} has a mass of 1750 MeV, when the mass of the Θ^+ is fixed at the observed value of 1540 MeV. Although no strong explanations for the narrow width are given, it may be possible to explain the narrowness by the weak coupling of the Θ^+ to the KN mode. However, the large width of the Roper resonance in the same multiplet makes this highly non-trivial.

2.2.6 The Karliner-Lipkin diquark triquark model

A challenge for any exotic baryon model is to explain the positive parity predicted by the Skyrme model for the Θ^+ state, while the 'standard' pentaquark involves five quarks in an S-wave and therefore has negative parity. If Θ^+ has indeed positive parity, then there is clearly one unit of angular momentum, which makes the calculations difficult.

In the diquark-triquark model two color nonsinglet clusters (a ud diquark and a $us\bar{s}$ triquark) are in a relative P-wave and kept together by the color-electric force [85, 9]. The P-wave configuration ensures that the parity is positive. The clusters are separated by a distance larger than the range of the color-magnetic force. The repulsive hyperfine interaction between two quarks of the same flavor is then not felt between the clusters.

CHAPTER III

The HERMES Experiment

The HERMES experiment[1] was designed to study the quark-gluon spin structure of the nucleon. This is accomplished by studying asymmetries in the cross section for deep-inelastic scattering (DIS) of longitudinally polarized leptons on light nuclear targets. Several other experiments have provided accurate data for the *inclusive* cross section asymmetry, *i.e.* when only the scattered lepton is detected. At the HERMES experiment, *semi-inclusive* deep-inelastic scattering (SIDIS) processes are studied, *i.e.* some of the hadrons created in the fragmentation of the target nucleon or the hadronization of the struck quark are detected as well. These data offer a way to determine the flavor of the struck quark, and thus access the flavor-dependent spin structure functions. Along with the accurate determination of the particle track and momentum, good particle identification (PID) is crucial for the HERMES experiment. It can identify pions, kaons, protons, and leptons with high efficiency and low contamination.

The HERMES experiment is located in one of the four experimental halls of the lepton-proton collider HERA[2] at the research institute DESY[3] in Hamburg, Germany. It only uses the lepton beam of the HERA collider. The lepton storage ring can be filled with either electrons or positrons,[4] which are accelerated to an energy of 27.6 GeV.

One of the characteristic properties of the HERA lepton storage ring is the large transverse self-polarization of the lepton beam built up in the arcs of the accelerator. Before the leptons reach the HERMES experiment, a spin rotator aligns the spin longitudinally, parallel or antiparallel to the beam direction. Behind the HERMES experiment the polarization of the leptons is measured with the longitudinal polarimeter (LPOL). Finally, before the leptons continue their way through the arcs of the HERA collider, their spin is rotated back transverse to the beam direction.

In the following sections the HERA lepton-proton collider, the HERMES target, and the HERMES spectrometer are described in more detail. A section on Monte Carlo simulations of reactions at the HERMES experiment follows. Finally, the operation of the longitudinal polarimeter (LPOL) of the lepton beam is discussed in light of high-precision lepton beam polarimetry measurements at future collider experiments (in particular the electron-ion collider EIC).

3.1 The HERA lepton-proton collider

Until the end of operations in 2007, the HERA lepton-proton collider at the research institute DESY was one of the world's highest-resolution probes into the quark structure of the proton. At the HERA collider electrons or positrons were accelerated to an energy of 27.6 GeV, and protons to an energy of 920 GeV. A schematic overview of the different accelerator systems, including the pre-accelerator PETRA, at the research institute DESY is shown in figure 3.1.

---

[1] HERA Measurement of Spin

[2] Hadron-Elektron Ringanlage, hadron-electron ring facility

[3] Named after the Deutsches Elektron Synchrotron, the first synchrotron built at the research institute, which accelerated its first electrons to 7.4 GeV on January 1, 1960.

[4] Since the analysis presented in this work is independent of the charge of the colliding leptons, positrons are included when the text refers only to electrons, unless explicitly mentioned otherwise.



Figure 3.1: The research institute DESY consists of several accelerator systems. After generation of a low-energetic proton and lepton beam in the LINAC and DESY accelerators, the synchrotron PETRA was used as pre-accelerator for the approximately 6.3 km long HERA storage rings. The electrons or positrons (red) were accelerated to 27.6 GeV, the protons (blue) were accelerated to 920 GeV. At two interaction points (North and South) the leptons and protons were brought in collision for the H1 and ZEUS experiments. The HERA-B experiment in the West hall used only the proton beam. The HERMES experiment was located in the East hall and only used the electron or positron beam. Taken from reference [86].

At two interactions points the lepton and proton beams collided at a center-of-mass energy of approximately 300 GeV. At these two interaction points, located in the North and South Hall, the experiments H1 and ZEUS studied deep-inelastic scattering reactions in the low-x region.[5] The other two experimental halls of the HERA collider were occupied by fixed target experiments. The HERA-B experiment searched for CP-symmetry breaking in the decay of B-mesons generated in the interactions of protons in the proton beam halo with movable metal wires. The HERA-B experiment was terminated in 2003 and the HERA-B spectrometer was removed from the West Hall. The second fixed target experiment at the HERA collider was, of course, the HERMES experiment. The HERMES experiment used only the lepton beam of the HERA collider.

In the HERA lepton storage ring, electrons or positrons circulate in bunches with a length of 37 ps and separated by 96 ns. At most 220 bunches can be filled simultaneously, although in practice rarely more than 185 bunches are filled.[6] The lepton bunches with an energy of 12 GeV are injected in several steps from the PETRA pre-accelerator into the HERA lepton storage ring, until a current of approximately 45 mA is reached. They are then accelerated to the full energy of 27.6 GeV. During data collection with polarized target gases, the lepton beam current decreases exponentially with a life time of over 10 hours until it reaches approximately 15 mA. By injecting unpolarized gases with higher density in the HERMES target the life time is then further reduced and one hour later the remaining leptons are usually dumped. A new lepton beam is injected and data taking resumes. In a very similar fashion the HERA proton storage ring is operated. Because the proton beam current decreases more slowly, up to three regular lepton fills can be obtained for every proton fill.

Only very few leptons in the lepton beam interact in the HERMES target. Essentially, there is no influence of the HERMES target on the lepton beam at the usual low densities. To remove the synchrotron radiation travelling with the beam, two collimators are positioned upstream of the HERMES spectrometer.

A small asymmetry in the spin-flip amplitudes for leptons when emitting synchrotron radiation in a magnetic dipole field is amplified by the repeated revolutions in the HERA lepton storage ring and enhances the population of the spin state parallel (antiparallel) to the magnetic field for positrons (electrons), resulting in the transverse polarization of the lepton beam, in particular vertically upwards for both electrons and positrons. This effect, known as the Sokolov-Ternov effect [87], allows for a theoretical asymptotic self-polarization of the lepton beam up to 92.4% with a rise time of 37 minutes [88]. Due to the interaction with the proton beam and various other depolarizing effects, the lepton polarization obtained in the HERA storage ring was almost two times lower.

To perform physics experiments with longitudinally polarized lepton beams, the spin direction is rotated by 90°, before the leptons reach the HERMES experiment, to the direction parallel to the lepton beam. This is achieved in a spin rotator, which consists of three consecutive pairs of dipole magnets [89]. The now longitudinally polarized leptons interact with the target gas in the HERMES experiment, or with the proton beam in the collider experiments. The spin direction is rotated back to the transverse direction in a second spin rotator, before the leptons leave the straight section and enter the bending section again. During ther HERA I period until 2001, only one pair of spin rotators was installed, positioned around the HERMES experiment. During this period the asymptotic lepton polarization was typically around 55% with a risetime of 23 minutes [90]. In 2001 spin rotator pairs were installed around the two collider experiments, as part of the HERA II upgrade. After this upgrade, which resulted mainly in an increase of the collider luminosity, the lepton polarization was typically only around 50%, due to the increased beam-beam interactions.

At two locations on the HERA lepton storage ring, the beam polarization is measured continuously. Both polarimeter systems are based on asymmetries in the Compton back-scattering cross section of polarized laser photons from the lepton beam. Downstream from the HERMES experiment, but before the spin rotator, the longitudinal polarization of the leptons is measured. The energy-asymmetry in the Compton cross section is used in the longitudinal polarimeter (LPOL), which will be described in section 3.5. In a straight section without spin rotators, the transverse polarimeter (TPOL) measures the spatial asymmetry of the back-scattered Compton photons. Both polarimeters provide polarization measurements with a sta-

[5] In DIS reactions the variable x can be interpreted as the fraction of the total momentum of the nucleon that is carried by the struck quark or gluon.

[6] Empty lepton bunches, not necessarily combined with a corresponding empty proton bunch, provide a convenient way of studying background events. However, the real reason for the empty bunches is purely technical: A long bunch-free region is needed for the ramp-up kicker magnets of the proton beam abort system.



Figure 3.2: Schematic view of the HERMES target system. Taken from reference [91].

tistical precision of approximately 1% and a systematic uncertainty of 1.6% (LPOL) and 3.0% (TPOL).

3.2 The HERMES target region

The HERMES target consists of a polarized or unpolarized gas in a storage cell internal to the HERA lepton storage ring [92]. A schematic view of the target region is shown in figure 3.2. The target cell is constructed as a 40 cm long, open-ended T-shaped tube that confines the gas atoms in a region around the lepton beam. The gas atoms leak out the ends and are pumped away by a high speed pumping system. The requirement of data collection without significant reduction of the lepton beam lifetime[7] limits the target density. At the end of lepton fills unpolarized high density data are collected with H_2, D_2, ^{3}He, N_2, Xe, or Kr gases, providing a substantial data sample for the study of unpolarized reactions and nuclear effects.

Since the initial commissioning of the HERMES experiment in 1995, several target gases and target polarization states have been used. In 1995, the gas in the storage cell was supplied by an optically pumped, longitudinally polarized $^3\overrightarrow{\mathrm{He}}$ cell. In the period 1996–1997, an atomic beam source (ABS) was used to produce a longitudinally polarized $\overrightarrow{\mathrm{H}}$ target, and in 1998–2000 the ABS was modified for operation with longitudinally polarized $\overrightarrow{\mathrm{D}}$.

In 2001, the target region was modified substantially for the operation with the transversely polarized target gases $\uparrow$H and $\uparrow$D [93]. A magnet providing the transversely oriented holding field for the target gases was installed.[8] The synchrotron radiation generated by the deflection of the lepton beam limited the strength of the magnetic field to approximately 297 mT. The variations in the volume of the storage cell were smaller than $\Delta B_x = 0.60\,\mathrm{mT}$, $\Delta B_y = 0.15\,\mathrm{mT}$, and $\Delta B_z = 0.05\,\mathrm{mT}$. The variations outside of the storage cell volume were larger. The shift of the beam position in the transverse holding field amounts to approximately 2 mm, an effect that can be observed easily in the collected data.

At the end of 2005, the transversely polarized target system was removed. For the recoil data taking period 2006–2007 only unpolarized target gases were needed. To accomodate the recoil detector, the length of the storage cell was reduced to 15 cm. The elliptical cross section of $21 \times 9\,\mathrm{mm}^2$ was unchanged.

[7]This is quantified by the requirement of a lepton beam lifetime larger than 45 hours if all other contributions to the lifetime are zero.

[8]A superconducting magnet, with longitudinal field direction, was installed during the years 1998–2000, but the effects of the transverse magnetic field on the tracking of scattered particles are more important.



Figure 3.3: Schematic view of the HERMES spectrometer during the years 2002–2005. The target is shown in yellow, the spectrometer magnet in blue, the tracking detectors in red, and the particle identification detectors in green. Taken from reference [91].

3.3 The HERMES spectrometer

The HERMES spectrometer is a forward spectrometer, a design typically used for fixed-target experiments. It consists of several detector components to track and identify scattered and produced particles. The spectrometer is described in great detail in reference [91]. In this section, only the detector components most relevant for the analysis of exotic baryons are described in detail.

The central component of the HERMES spectrometer is the H-shaped dipole magnet with an integrated field strength of $1.3\,\mathrm{T\cdot m}$. In the symmetry plane of the spectrometer, a massive steel plate shields the lepton and proton beams as they pass through the spectrometer magnet, separated by a distance of $72\,\mathrm{cm}$. Inside the shielding of the positron beam pipe, a correction coil corrects for fringe fields and the imperfections of the magnet shielding. The coil also serves to compensate the magnetic holding field of the target system when operating with transversely polarized target gases.

The horizontal symmetry plane defined by the shielding plate inside the spectrometer magnet separates the spectrometer in two identical *top* and *bottom* halves. The steel plate limits the acceptance at small angles in the vertical direction. A schematic side-view of the HERMES spectrometer during the years 2002–2005 is shown in figure 3.3.

The initial trajectory of the scattered particles is determined by the front tracking system, which consists of the front drift chambers (DVC and FC). Behind the spectrometer magnet, the momentum measurement is completed by the back drift chambers (BC). Inside the magnet, proportional chambers (MC) help match the front and back tracks, and allow track reconstruction for low-momentum particles that do not reach the back region.

In the front region, the acceptance for the decay of $\Lambda(1115)$ hyperons was increased in 2001 by the installation of two silicon detectors just downstream of the target region [94, 95]. In the beginning of 2006 a recoil detector was installed, adding acceptance for the recoiling target proton in DVCS events [96]. In the present analysis, these detectors have not been used.

Particle identification (PID) is provided by scintillator hodoscopes (H1 and H2), a transition radiation detector (TRD), a lead-glass calorimeter (CALO), and a ring imaging Čerenkov detector (RICH). The RICH detector identifies pions, kaons, and protons. Combination of the signals in the particle identification detectors allow for a hadron rejection factor of higher than 10^4. Until 1997, before the installation of the RICH detector, a single-gas radiator threshold Čerenkov detector (CER) was installed. The threshold Čerenkov detector allowed only for limited separation of pions from heavier hadrons.

3.3.1 The particle tracking detectors

The particle tracking system is used to determine the event vertex in the target cell and to measure the momentum and the angles of the scattered particles. The momentum resolution is better than 1.3%, and is limited by bremsstrahlung in the walls of the target cell, the vacuum window, and the front tracking detectors. The resolution in the scattering angle is better than 0.6 mrad and limited by multiple scattering in these materials.

The drift vertex chambers (DVC) and the front drift chambers (FC) provide the vertex reconstruction in the target cell and the definition of the scattering angles. The back drift chambers (BC) measure the magnetic deflection and hence the momentum of the detected particles. Proportional chambers inside the magnet (MC) are used for the momentum analysis of low energy decay products that are deflected too strongly to reach the back chambers.

The horizontal length of the drift chambers precluded the use of long horizontal wires. All planes use one of three wire orientations, either vertically (X) or tilted 30° right or left from the vertical axis (U and V).

The drift chambers

The drift chambers DVC, FC, MC, and BC are of the conventional horizontal-drift type. Each layer of drift cells consists of a plane of alternating anode and cathode wires between a pair of cathode foils. The charged particles passing through the detector generate an ionization avalanche, which is detected by the wires spun across the chambers. The chambers are assembled as modules consisting of six layers in three coordinate doublets (UU', XX' and VV'). The X', U' and V' planes are staggered by half the cell size in order to help resolve tracking ambiguities.

To ensure redundancy in the front tracking system and to provide a larger acceptance for muon detection, the drift vertex chamber DVC was constructed and installed after one year of HERMES operation. They are followed by the front chambers FC1 and FC2, providing good spatial resolution immediately in front of the spectrometer magnet.

The three magnet chambers MC are located in the gap of the spectrometer magnet and were originally intended to help resolve multiple tracks in case of high-multiplicity events. Since low backgrounds have made this unnecessary, their primary function is now the momentum analysis of particles with energies below 2 GeV, for example from the decay of the Λ hyperon. With the MCs particle tracks with a momentum as low as 0.5 GeV can be reconstructed.

Finally, the four back chambers BC are located behind the spectrometer magnet.

Track reconstruction

The HERMES track reconstruction program uses a tree-search algorithm for fast track finding and a look-up table for fast momentum determination of the tracks. Except for small curvatures caused by the magnetic fringe fields, the track projections are approximately straight lines outside of the magnetic field region. The basic idea of pattern recognition using the tree-search algorithm is to look at the whole hit pattern of the detectors with increasing resolution. In each step of the iteration, the algorithm checks if the pattern (at a given resolution) contains a sub-pattern that is consistent with an allowed track.

All allowed patterns are generated at the initialisation of the program. They are stored in a database. Because a given pattern at one resolution constrains the number of allowed patterns at a lower resolution, it is possible to reduce the CPU time involved. Symmetry is used to reduce the number of patterns that have to be stored.

To avoid the CPU intensive task of tracking a particle in a magnetic field, a large look-up table is generated during the initialization of the tracking program using the measured magnetic field map. It contains the momentum of a given track as a function of the track parameters in front and behind the magnet.

Transverse magnet correction After the installation of the transversely polarized target in 2001, the existing tracking algorithm was updated to take into account the slight deflection of the scattered particles in the transverse target magnetic field. Two algorithms were developed to perform this correction [97]. The first method (TMC1) is based on a transfer-matrix approach, the second method (TMC2) uses reference tracks. Both algorithms return the corrected, reconstructed interaction point of the track with the lepton beam. For secondary vertices, which are displaced from the lepton beam, these methods are not applicable.

3.3.2 The particle identification detectors

The HERMES particle identification (PID) system discriminates between positrons, pions, kaons, and protons. The PID system consists of four subsystems: two plastic scintillator hodoscopes, a transition radiation detector, a lead-glass calorimeter, and a dual radiator ring imaging Čerenkov detector. The scintillator hodoscopes and the electromagnetic calorimeter are also used in the first level trigger to select which events to consider for further processing.

Forward trigger scintillator (H0)

While technically not a PID detector, the forward trigger scintillator (H0) is located directly upstream of the front drift chamber to eliminate triggers from particle from the proton beam. A backward-going particle produces a pulse that is displaced by 36 ns from the normal trigger condition.

Scintillator hodoscopes (H1 and H2)

A scintillator hodoscope (H1) and a preshower scintillator counter (H2) provide fast trigger signals and information for particle identification. Both hodoscopes are composed of vertical, fast scintillator paddles of 1 cm thick and 9 cm wide with 2–3 mm overlap between each paddle. The scintillation light is detected by photomultiplier tubes coupled with light guides to the outer end of each scintillator paddle. In the passive radiator of the preshower scintillator counter H2, consisting of two radiation lengths of lead, electromagnetic showers are generated. The energy deposit in the scintillator paddles facilitates the discrimination between leptons and hadrons. Positrons typically deposit at least ten times more energy than hadrons, allowing for a pion rejection of 10 for lepton detection.

The scintillator hodoscopes can also be used as a time-of-flight (TOF) particle identification detector for low momentum hadrons [98]. This complements the hadron identification capabilities of the RICH detector (see below). The timing of the output signals in the PMTs of H1 and H2 are measured with respect to the bunch crossing time in the center of the target cell. The path length of each track is calculated from the straight partial tracks in the front region and the back region.[9] To account for differences in cable length, the time-of-flight timings are calibrated with cleanly identified electrons which move essentially at the speed of light. The (squared) mass of hadrons is then extracted from the relativistic momentum

$$p = m\frac{\beta}{\sqrt{1-\beta^2}}, \tag{3.1}$$

where β is the relativistic speed v/c. The squared particle mass obtained from the timing in H1 is shown in figure 3.4 for hadrons with a momentum in the range 0.6–1.1 GeV (left) and 1.5–2.0 GeV (right). Kaon identification is limited to momenta below 1.5 GeV, but pions and protons can be distinguished up to 2.9 GeV with efficiencies (contamination) above 98% and 85% (below 4% and 6%).

Electromagnetic calorimeter (CALO)

The calorimeter provides a first level trigger and measures the energy of electrons, positrons, and photons, but suppresses pions by a factor of more than 1000. It is composed of two times 420 lead-glass blocks of transverse dimensions $9 \times 9\,\mathrm{cm}^2$ and 18 radiation lengths long, viewed from the rear by a photomultiplier tube. Monitoring of aging effect in the glass-blocks is achieved using a dye laser, which sends light through glass fibers to every photomultiplier and a reference photodiode. Combining the preshower hodoscope (H2) with the electromagnetic calorimeter, the pion rejection factor integrated over all energies is approximately $2.5 \cdot 10^3$.

[9] This approximation is adequate because the deflection angle in the spectrometer magnet is only $\pi/20$.



Figure 3.4: Time-of-flight hadron identification with the scintillater hodoscopes H1. Pions, kaons, and protons can be separated in the distribution of the squared mass m^2 for the momentum range 0.6–1.1 GeV (left). For the momentum range 1.5–2.0 GeV pions and protons can be distinguished. Taken from reference [98].

Transition radiation detector (TRD)

The purpose of the transition radiation detector (TRD) is to provide a pion rejection factor of at least 100 at positron energies higher than 5 GeV. At these energies, only electrons and positrons produce transition radiation in the form of X-rays in the 6 cm thick two-dimensional matrix of dielectric fibers. The generated X-rays are detected in proportional wire chambers. Pions deposit energy directly in the chambers due to ionisation of the chamber gas, but leptons deposit on average twice as much energy in addition to the energy in the X-rays. The average deposited energy of hadrons and leptons is shown in figure 3.5. The pion rejection factor satisfies the design goals, and reaches a value of 80 even below an energy of 5 GeV.

Single gas-radiator threshold Čerenkov counters (CER)

In the original setup of the HERMES spectrometer, used until 1997, pion identification was provided by a pair (top and bottom) of single gas-radiator threshold Čerenkov counters (CER). The Čerenkov counters were located between the back drift chambers, where the RICH was later installed. They were filled with freon, C_4H_{10}, and the produced Čerenkov radiation were detected by an array of 20 spherical mirrors that focus the light on 12.7 cm wide photomultiplier tubes. The momentum threshold was 3.8 GeV for pions, 13.6 GeV for kaons, and 25.8 GeV for protons. The number of detected photoelectrons for a track allows to distinguish between pions and heavier hadrons, but not between kaons and protons.

Lepton-hadron separation

The lepton-hadron separation of the separate detectors can be improved by combining the responses of the transition radiation detector (TRD), the Čerenkov counters (CER), the preshower hodoscope (PRE), and the electromagnetic calorimeter (CALO) using a Bayesian probabilistic approach. For each detector the so-called parent distributions specify the probability $\mathcal{L}^i$ to obtain a specific detector response for a given particle type i. They are determined from the measured responses in each detector using stringent selection requirements on the other detectors. The probability P^i that a given detector response originates in a particle of type i is then defined as

$$P^i = \frac{\phi^i \mathcal{L}^i}{\sum_j \phi^j \mathcal{L}^j}, \tag{3.2}$$



Figure 3.5: The average energy deposited in the transition radiation detector (TRD) by hadrons and leptons, averaged over all momenta. Taken from reference [91].

where the sum runs over all particle types. The flux factor ϕ^i is the *a priori* probability that a track with given track parameters originates from a particle of type i. These flux factors are determined from the collected data with an iterative procedure starting from equal flux factors for all particle types.

The logarithm of the ratio of the probabilities P^h and P^e is calculated for each detector,

$$PID = \log_{10} \frac{P^e}{P^h}. \tag{3.3}$$

The following sums of the particle identification detectors are defined:

$$PID_3 = PID_{CALO} + PID_{PRE} + PID_{CER}, \tag{3.4}$$
$$PID_5 = PID_{TRD}. \tag{3.5}$$

In figure 3.6 the variables PID_3 and PID_5 are shown for events collected by the spectrometer. The separation between leptons (left) and hadrons (right) is clearly visible.

Usually, the sum $PID_3 + PID_5$ is required to be smaller than zero for hadrons and larger than zero for leptons. In hadron samples obtained by requiring $PID_3 + PID_5 < -1$, the probability that a track originated from a hadron is at least 10 times larger than the probability that it originated from a lepton.

Dual radiator ring imaging Čerenkov detector (RICH)

The dual radiator ring imaging Čerenkov detector (RICH) was designed to provide hadron identification for pions, kaons and protons in the momentum range from 2 to 15 GeV. The RICH detector is described in detail in reference [99]. A schematic view of the detector setup is shown in the left panel of figure 3.7. It uses a wall of silica aerogel radiator tiles and a gas radiator volume with freon, C_4H_{10}, and the whole detector is located in the same location of the previous Čerenkov counter. The charged particles transversing the radiator materials emit Čerenkov radiation in a cone with an opening angle depending on the momentum and the particle type.



Figure 3.6: The combination of the signals in the Čerenkov counters, preshower-hodoscopes, and electromagnetic calorimeter (left axis) and the deposited energy in the TRD detector (right axis) allows for separation of leptons (left) and hadrons (right) with high efficiency and low contamination. Taken from reference [91].



Figure 3.7: The dual radiator ring imaging Čerenkov detector (RICH). A schematic view of the detector setup and radiator configuration is shown in the left panel. The observed hit pattern for a typical three-track event is shown in the right panel. Taken from reference [99].