Status Report of HERMES

Pasquale Di Nezza

(on behalf of HERMES Collaboration)

- First measurement of transversity
- Exotic baryons: the pentaquark
- The spectrometer and the data taking

Physics Research Committee, DESY Oct 30/31, 2003

Even they were puzzled ...





DIS + SIDIS cross section



$$d\sigma = d\sigma_{UU} + \cos 2\phi d\sigma_{UU} + \frac{1}{Q}\cos\phi d\sigma_{UU} + \lambda\frac{1}{Q}\sin\phi d\sigma_{LU}$$

$$+ S_L\left[\sin 2\phi d\sigma_{UL} + \frac{1}{Q}\sin\phi d\sigma_{UL}\right] + \lambda S_L\left[d\sigma_{LL} + \frac{1}{Q}\cos\phi d\sigma_{LL}\right]$$

$$+ S_T\left[\sin(\phi+\phi_S)d\sigma_{UT} + \sin(\phi-\phi_S)d\sigma_{UT} + \sin(3\phi-\phi_S)d\sigma_{UT} + \frac{1}{Q}\sin(2\phi-\phi_S)d\sigma_{UT}\right]$$

$$+ \lambda S_T\left[\cos(\phi-\phi_S)d\sigma_{LT} + \frac{1}{Q}\cos(2\phi-\phi_S)d\sigma_{LT} + \sin(3\phi-\phi_S)d\sigma_{LT}\right] + ...$$

Operator decomposition of the Correlation Function at Twist-2

$$\Phi^{Tw\,2}_{Corr}(x) = \frac{1}{2}\{f_1(x) + S_L\, g_1(x)\gamma_5 + h_1(x)\gamma_5\gamma S_T\}\gamma^-$$

$f_1 =$ 

Unpolarized quark in unpolarized nucleon

$g_{1L} =$ 

Helicity difference

$\frac{1}{2}$ 

Extremely well known function

$\frac{1}{2}$ 

Function measured by HERMES (EMC,SMC, E142-143-154-155)

Operator decomposition of the Correlation Function at Twist-2

$$\Phi^{Tw\,2}_{Corr}(x) = \frac{1}{2}\{f_1(x) + S_L\, g_1(x)\gamma_5 + h_1(x)\gamma_5\not{S}_T\}\not{n}^-$$

$h_1 =$  $-$

Single helicity flip



NOT ALLOWED IN E.M. INTERACTIONS

Operator decomposition of the Correlation Function at Twist-2

$$\Phi^{Tw\,2}_{Corr}(x) = \frac{1}{2}\{f_1($$

$h_1 =$  $-$

Single helicity flip



flip

Peculiarity of transversity

Relativistic nature of quark.
In absence of relativistic effects $h_1(x)=g_1(x)$

Q^2 -evolution. Unlike for $g_1^p(x)$,
the gluon doesn't mix with quark in $h_1^p(x)$

High sensitivity to the valence quark polarization
q and $\bar{q}$ have opposite sign.

Tensor charge: first moment of h_1.
Calculable by lattice QCD.

DIS + SIDIS cross section



$$d\sigma = d\sigma_{UU} + \cos 2\phi d\sigma_{UU} + \frac{1}{Q}\cos\phi d\sigma_{UU} + \lambda\frac{1}{Q}\sin\phi d\sigma_{LU}$$

$$+ S_L\left[\sin 2\phi d\sigma_{UL} + \frac{1}{Q}\sin\phi d\sigma_{UL}\right] + \lambda S_L\left[d\sigma_{LL} + \frac{1}{Q}\cos\phi d\sigma_{LL}\right]$$

$$+ S_T\left[\underbrace{\sin(\phi+\phi_S)d\sigma_{UT}}_{h_1\otimes H_1^\perp} + \underbrace{\sin(\phi-\phi_S)d\sigma_{UT}}_{f_{1T}^\perp\otimes D_1} + \sin\right]$$

(Collins)

(Sivers)

Peculiarity of $f_{1t}^\perp$

- Chiral-even naïve T-odd DF
- Related to parton orbital momentum
- Violates naïve universality of PDF
- Different sign of $f_{1t}^\perp$ in DY

Definition of Angles and Asymmetries



$(\phi - \phi_S)$

Angle of hadron relative to initial quark spin (Sivers)

$(\phi + \phi_S)$

Angle of hadron relative to final quark spin (Collins)

- Sivers-Collins effects can be distinguished only with transverse polarised target.
- Large asymmetry has been measured in inclusive π production ($p^{\uparrow}p \rightarrow \pi X$) → jet axis not known. Both mechanisms involved.

Definition of Angles and Asymmetries



$$A_{UT}^{h}(\phi,\phi_S) = \frac{1}{|S_T|}\frac{N_h^{\uparrow}(\phi,\phi_S) - N_h^{\downarrow}(\phi,\phi_S)}{N_h^{\uparrow}(\phi,\phi_S) + N_h^{\downarrow}(\phi,\phi_S)} =$$

$$= A_{UT}^{Collins}\sin(\phi+\phi_S) + A_{UT}^{Sivers}\sin(\phi-\phi_S)$$

$$A_{UT}^{Collins} = \frac{\langle |p_{h\perp}|/M_h \cdot \sin(\phi+\phi_S)\rangle_{UT}}{\langle 1\rangle_{UU}} = |S_T| \frac{\sum_q e_q^2 h_1^q H_1^{\perp(1),q\to h}}{\sum_q e_q^2 f_1^q D_1^{q\to h}} \cdot \frac{1-y}{1-y+y^2/2}$$

$$A_{UT}^{Sivers} = \frac{\langle |p_{h\perp}|/M_p \cdot \sin(\phi-\phi_S)\rangle_{UT}}{\langle 1\rangle_{UU}} = |S_T| \frac{\sum_q e_q^2 f_{1T}^{\perp,q} D_1^{q\to h}}{\sum_q e_q^2 f_1^q D_1^{q\to h}}$$

Transverse asymmetry for π^+, π^-, π^0

Sivers Moments



Collins Moments



- First measurement of naïve T-odd DF in DIS (orbital mom)
- Opposite sign from RHIC DY ?

- Much of plausible value ranges of transversity and disfavoured Collins function are excluded

Hadron spectroscopy

- Standard Quark Model
  - allows hadrons as
    - mesons ($q\bar{q}$)
    - baryons (qqq)
  - also allows "non-standard" or exotic hadron states
    - multiquark mesons ($qq\bar{q}\bar{q}$)
    - multiquark baryons ($qqqq\bar{q}$) -> appear as baryon resonances
    - hybrid states ($q\bar{q}g$ or $qqqg$)
    - dibaryons ($qqqqq$)
    - glueballs







Search for Exotic Baryon States

Predictions:

Bag models (Jaffe '77; De Swart '80): 1.8-1.9 GeV

Skyrme model (Praszalowicz '87, Walisser '92): 1.3 – 1.8 GeV

Chiral-Soliton Model (Diakonov, Polyakov '97)

NA49



New anti-decuplet



The 3 corners are exotics

Spring-8



CLAS



hermes

- Complete Particle Identification with high efficiency: 98% (π), 88% (K), 85% (p)
- Direct reconstruction: detection of each decayptc, Inv mass reconstuction (other exp. missing mass)
- Excellent invariant mass systematics: ± 2 MeV

Event Reconstruction

$$e^+ d \rightarrow K^0_s p X \rightarrow \pi^+ \pi^- p X$$



- RICH ID: 1-15 GeV for π, 4-9 GeV for p
- K_s decay length>7 cm
- Distance cut: $\pi^+\pi^-$, K_s p, θ^+ beam
- Collinearity K_s $\pi\pi$ < 45 mrad
- 485< K_s-mass<509 MeV
- No Λ

Particle Identification Proof





Good identification of Λ means clear identification of p

Good K_s identification within the kinematical cuts of this analysis

Detector Mass calibration



$\rho^0(770)$



$\phi(1020)$

Masses in agreement with PDG values ± 2 MeV



$\bar{\Lambda}(1116)$



$\Lambda^*(1520)$

Measured Invariant Mass

Excess at 1526 M

Background approximated by a polynomial

Widt



Monte Carlo Simulation

- Simulated resonance at 1540 MeV
- Simulated Γ=2 MeV
- Decay in $K_s p$
- Full detector simulation

- Recons. Mass at 1540 ± 0.3 MeV
- Recons. σ=7 ± 0.2 MeV



Status of the Spectrometer



Spectrometer fully debugged after the shutdown
Each single detector is operative!
80K DIS collected in 2003-runII with transverse polarized hydrogen target

Conclusions

- First observation of non-zero Sivers effect
- Sizeable Collins asymmetries measured for π^0 and π^-
- A kind of brain storm is underway for model interpretation

2003: milestone from HERMES transverse asymmetries
>2004: results from HERMES, COMPASS, BELLE(RHIC-rIII, CLAS)

Conclusions

- First observation of non-zero Sivers effect
- Sizeable Collins asymmetries measured for π^0 and π^-
- A kind of brain storm is underway for model interpretation
- A narrow exotic baryon resonance has been directly reconstructed
- Most precise determination of the mass { $M = 1526 \pm 2 \pm 2$ MeV; $\sigma = 7.5 \pm 2.4$ MeV
- Background description and Monte Carlo simulation

Taking data for physics