


Nuclear medium dependence of transverse Λ polarisation in quasi-real photoproduction

Klaus Rith

University of Erlangen-Nürnberg & DESY

On behalf of the  collaboration


27.5 GeV e^+ / e^- beam of HERA



Internal gas targets

polarized : $^1H, ^1H^\uparrow, ^2H, ^3He$
 unpolarized: $^1H, ^2H, ^3He, ^4He, ^{14}N, ^{20}Ne, ^{84}Kr, ^{131}Xe$

● tracking: $\delta p/p \sim 2\%$, $\delta\Theta < 0.6$ mrad, 40-220 mrad




● PID: RICH, TRD, Preshower, Calorimeter
lepton-hadron separation > 98%

Reconstruction of Λ events




Quasi-real photoproduction

$Q^2 < 0.05 \text{ GeV}^2$ for 80% of events
 $<\nu> = 15.6 \text{ GeV}$



Background suppression:
 Cherenkov information +
 vertex cuts

1995-2000 data
 (all targets except Xe)



$$N_{\Lambda} \approx 250k$$

$$N_{\bar{\Lambda}} \approx 50k$$



Parity violating decay $\Lambda \rightarrow \pi^- p$:
 p preferentially emitted along Λ spin


$$\frac{dN}{d\Omega_p} = \frac{dN_0}{d\Omega_p} (1 + \alpha P_\Lambda \cos \theta_p) \quad (\text{in } \Lambda \text{ CMS})$$

$\alpha = 0.642 \pm 0.013$




Unpolarised beam and target ($P_B P_T = 0.0000 \pm 0.0005$):
 Spontaneous polarisation is directed along \hat{n}

$$\vec{P}_\Lambda = P_\Lambda \cdot \hat{n}, \quad \hat{n} = \frac{\vec{p}_e \times \vec{p}_\Lambda}{|\vec{p}_e \times \vec{p}_\Lambda|}$$




Extraction of Λ polarisation

Formalism is based on
up/down mirror (geometrical) symmetry of the
detector



$$\langle \cos \theta \rangle_0^{up} = - \langle \cos \theta \rangle_0^{down}$$




$$P_\Lambda = \frac{\langle \cos \theta_p \rangle}{\alpha \langle \cos^2 \theta_p \rangle} = \frac{\frac{1}{N_\Lambda} \sum_{i=1}^{N_\Lambda} \cos \theta_p}{\alpha \frac{1}{N_\Lambda} \sum_{i=1}^{N_\Lambda} \cos^2 \theta_p}$$

Kinematical dependences of P_Λ

Phys.Rev.D76 (2007) 092008

1995-2000 data (all targets except Xe)



$$\zeta = (E_\Lambda + p_{z\Lambda}) / (E_e + p_e)$$

ζ and x_F are correlated
 $\zeta > 0.25 \leftrightarrow x_F > 0$

Λ : $P_n = 0.078 \pm 0.006_{\text{stat.}} \pm 0.012_{\text{syst.}}$

$\bar{\Lambda}$: $P_n = -0.025 \pm 0.015_{\text{stat.}} \pm 0.018_{\text{syst.}}$

● $P_n(\Lambda)$ is positive

● Opposite sign compared to
pion and proton beams





● Same sign as for K^- ($\bar{u}s$) and
 Σ^- ($dd\bar{s}$) beams

● Origin: s -quark content of γ ?

Kinematical dependences of P_n

1995-2000 data (all targets except Xe)

Phys.Rev.D76 (2007) 092008




$$\zeta = (E_\Lambda + p_{z\Lambda}) / (E_e + p_e)$$


$P_n(\Lambda)$ increases with increasing p_T

$P_n(\Lambda)$ is larger for low ξ (target fragm.) than for high ξ (current fragmentation)

A dependence in pA collisions




Experiment @ FNAL
 $p A \rightarrow \Lambda X$
 (targets Cu, Pb, Be)
 $p_{\text{beam}} = 400 \text{ GeV}$



Experiment @ BNL
 $p A \rightarrow \Lambda X$
 (targets H, D, Be)
 $p_{\text{beam}} = 28 \text{ GeV}$

A dependence of Λ polarisation



1995-2005 data; $N_{\Lambda} \approx 385k$
 (50 % more D + 25k Kr, 17k Xe)




$P_{\Lambda}(^1H) \gg P_{\Lambda}(^2H)$



P_{Λ} compatible with zero
for large A

→ $P_{\Lambda}(n) \ll P_{\Lambda}(p) ?$

A/Z dependence of Λ polarisation



$P_\Lambda(n) \ll P_\Lambda(p)$ not sufficient
to explain vanishing P_Λ for
large A

Additional nuclear medium
effects required for
explanation,
 P_Λ destroyed by FSI ?




● Transverse Λ polarisation observed in quasi-real photoproduction

● $P_n(\Lambda)$ is positive. Same sign as for $K^- (us)$ and $\Sigma^- (dds)$ beams. Origin: s -quark content of γ ?

● $P_\Lambda(^1H) \gg P_\Lambda(^2H) \rightarrow P_n(\Lambda)$ for neutrons substantially smaller than for protons ?

● Nuclear medium effects:
 $P_n(\Lambda)$ appears to vanish for large A (A/Z)

A/Z dependence of Λ polarisation



A dependence of Λ polarisation

